



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 33/574, 33/577, C07K 16/30, A61K 39/395, 47/48, C12Q 1/68, G01N 33/543		A2	(11) International Publication Number: WO 98/37418
			(43) International Publication Date: 27 August 1998 (27.08.98)
(21) International Application Number: PCT/US98/03690		(81) Designated States: AL, AM, AT, AU, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 25 February 1998 (25.02.98)		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(30) Priority Data:			
08/806,596 25 February 1997 (25.02.97) US 08/904,809 1 August 1997 (01.08.97) US Not furnished 9 February 1998 (09.02.98) US			
(71) Applicant: CORIXA CORPORATION [US/US]; Suite 200, 1124 Columbia Street, Seattle, WA 98104 (US).			
(72) Inventors: XU, Jiangchun; 15805 Southeast 43rd Place, Bellevue, WA 98006 (US). DILLON, Davin, C.; 21607 N.E. 24th Street, Redmond, WA 98053 (US).			
(74) Agents: MAKI, David, J. et al.; Seed and Berry LLP, 6300 Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-7092 (US).			

(54) Title: COMPOUNDS FOR IMMUNODIAGNOSIS OF PROSTATE CANCER AND METHODS FOR THEIR USE

(57) Abstract

Compounds and methods for diagnosing prostate cancer are provided. The inventive compounds include polypeptides containing at least a portion of a prostate tumor protein. The inventive polypeptides may be used to generate antibodies useful for the diagnosis and monitoring of prostate cancer. Nucleic acid sequences for preparing probes, primers, and polypeptides are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

COMPOUNDS FOR IMMUNODIAGNOSIS OF PROSTATE CANCER AND METHODS FOR THEIR USE

TECHNICAL FIELD

The present invention relates generally to the treatment and monitoring of prostate cancer. The invention is more particularly related to polypeptides comprising at least a portion of a prostate protein. Such polypeptides may be used for the production of compounds, such as antibodies, useful for diagnosing and monitoring the progression of prostate cancer, and possibly other tumor types, in a patient.

BACKGROUND OF THE INVENTION

Prostate cancer is the most common form of cancer among males, with an estimated incidence of 30% in men over the age of 50. Overwhelming clinical evidence shows that human prostate cancer has the propensity to metastasize to bone, and the disease appears to progress inevitably from androgen dependent to androgen refractory status, leading to increased patient mortality. This prevalent disease is currently the second leading cause of cancer death among men in the U.S.

In spite of considerable research into diagnosis and therapy of the disease, prostate cancer remains difficult to detect and to treat. Commonly, treatment is based on surgery and/or radiation therapy, but these methods are ineffective in a significant percentage of cases. Two previously identified prostate specific proteins - prostate specific antigen (PSA) and prostatic acid phosphatase (PAP) - have limited diagnostic and therapeutic potential. For example, PSA levels do not always correlate well with the presence of prostate cancer, being positive in a percentage of non-prostate cancer cases, including benign prostatic hyperplasia (BPH). Furthermore, PSA measurements correlate with prostate volume, and do not indicate the level of metastasis.

Accordingly, there remains a need in the art for improved and diagnostic methods for prostate cancer.

SUMMARY OF THE INVENTION

The present invention provides methods for immunodiagnosis of prostate cancer, together with kits for use in such methods. Polypeptides are disclosed which comprise at least an immunogenic portion of a prostate tumor protein or a variant of said protein that differs only in conservative substitutions and/or modifications, wherein the prostate tumor protein comprises an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177, 179-224 and variants thereof. Such polypeptides may be usefully employed in the diagnosis and monitoring of prostate cancer.

In one specific aspect of the present invention, methods are provided for detecting prostate cancer in a patient, comprising: (a) contacting a biological sample obtained from a patient with a binding agent that is capable of binding to one of the above polypeptides; and (b) detecting in the sample a protein or polypeptide that binds to the binding agent. In preferred embodiments, the binding agent is an antibody, most preferably a monoclonal antibody.

In related aspects, methods are provided for monitoring the progression of prostate cancer in a patient, comprising: (a) contacting a biological sample obtained from a patient with a binding agent that is capable of binding to one of the above polypeptides; (b) determining in the sample an amount of a protein or polypeptide that binds to the binding agent; (c) repeating steps (a) and (b); and comparing the amounts of polypeptide detected in steps (b) and (c).

Within related aspects, the present invention provides antibodies, preferably monoclonal antibodies, that bind to the inventive polypeptides, as well as diagnostic kits comprising such antibodies, and methods of using such antibodies to inhibit the development of prostate cancer.

The present invention further provides methods for detecting prostate cancer comprising: (a) obtaining a biological sample from a patient; (b) contacting the sample with a first and a second oligonucleotide primer in a polymerase chain reaction, at least one of the oligonucleotide primers being specific for a DNA molecule that encodes one of the above polypeptides; and (c) detecting in the sample a DNA sequence that amplifies in the presence of the first and second oligonucleotide primers. In a preferred embodiment, at least one of the

generate antibodies that detect the presence or absence of prostate cancer. Such modified sequences may be prepared and tested using, for example, the representative procedures described herein.

As used herein, a "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. In general, the following groups of amino acids represent conservative changes: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his.

Variants may also, or alternatively, contain other modifications, including the deletion or addition of amino acids that have minimal influence on the antigenic properties, secondary structure and hydrophobic nature of the polypeptide. For example, a polypeptide may be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (*e.g.*, poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

A nucleotide "variant" is a sequence that differs from the recited nucleotide sequence in having one or more nucleotide deletions, substitutions or additions. Such modifications may be readily introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis as taught, for example, by Adelman et al. (*DNA*, 2:183, 1983). Nucleotide variants may be naturally occurring allelic variants, or non-naturally occurring variants. Variant nucleotide sequences preferably exhibit at least about 70%, more preferably at least about 80% and most preferably at least about 90% identity to the recited sequence. Such variant nucleotide sequences will generally hybridize to the recite nucleotide sequence under stringent conditions. As used herein, "stringent conditions" refers to prewashing in a solution of 6X SSC, 0.2% SDS; hybridizing at 65 °C, 6X SSC, 0.2% SDS overnight; followed by two washes of 30 minutes each in 1X SSC, 0.1% SDS at 65 °C and two washes of 30 minutes each in 0.2X SSC, 0.1% SDS at 65 °C.

"Polypeptides" as used herein also include combination, or fusion, polypeptides. A "combination polypeptide" is a polypeptide comprising at least one of the above immunogenic portions and one or more additional immunogenic prostate tumor-specific sequences, which are joined via a peptide linkage into a single amino acid chain. The sequences may be joined directly (*i.e.*, with no intervening amino acids) or may be joined by way of a linked sequence (*e.g.*, Gly-Cys-Gly) that does not significantly diminish the immunogenic properties of the component polypeptides.

The prostate tumor proteins of the present invention, and DNA molecules encoding such proteins, may be isolated from prostate tumor tissue using any of a variety of methods well known in the art. DNA sequences corresponding to a gene (of a portion thereof) encoding one of the inventive prostate tumor proteins may be isolated from a prostate tumor cDNA library using a subtraction technique as described in detail below. Examples of such DNA sequences are provided in SEQ ID Nos: 1-107, 109-111, 115-171, 173-175, 177 and 179-224. Partial DNA sequences thus obtained may be used to design oligonucleotide primers for the amplification of full-length DNA sequences in a polymerase chain reaction (PCR), using techniques well known in the art (see, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY, 1989). Once a DNA sequence encoding a polypeptide is obtained, any of the above modifications may be readily introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis as taught, for example, by Adelman et al. (*DNA*, 2:183, 1983).

The prostate tumor polypeptides disclosed herein may also be generated by synthetic or recombinant means. Synthetic polypeptides having fewer than about 100 amino acids, and generally fewer than about 50 amino acids, may be generated using techniques well known to those of ordinary skill in the art. For example, such polypeptides may be synthesized using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain (see, for example, Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963). Equipment for automated synthesis of polypeptides is commercially available from

oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA molecule having a partial sequence selected from the group consisting of SEQ ID Nos: 2-3, 5-107, 109-11, 115-171, 173-175, 177 and 179-224.

In a further aspect, the present invention provides a method for detecting prostate cancer in a patient comprising: (a) obtaining a biological sample from the patient; (b) contacting the sample with an oligonucleotide probe specific for a DNA molecule that encodes one of the above polypeptides; and (c) detecting in the sample a DNA sequence that hybridizes to the oligonucleotide probe. Preferably, the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA molecule having a partial sequence selected from the group consisting of SEQ ID Nos: 2-3, 5-107, 109-11, 115-171, 173-175, 177 and 179-224.

In related aspects, diagnostic kits comprising the above oligonucleotide probes or primers are provided.

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for the immunodiagnosis and monitoring of prostate cancer. The inventive compositions are generally polypeptides that comprise at least a portion of a prostate tumor protein. Also included within the present invention are molecules (such as an antibody or fragment thereof) that bind to the inventive polypeptides. Such molecules are referred to herein as "binding agents."

In particular, the subject invention discloses polypeptides comprising at least a portion of a human prostate tumor protein, or a variant thereof such a protein, wherein the prostate tumor protein includes an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-11, 115-171, 173-175, 177, 179-224, the complements of said nucleotide sequences and variants thereof. As used herein, the term "polypeptide"

encompasses amino acid chains of any length, including full length proteins, wherein the amino acid residues are linked by covalent peptide bonds. Thus, a polypeptide comprising a portion of one of the above prostate proteins may consist entirely of the portion, or the portion may be present within a larger polypeptide that contains additional sequences. The additional sequences may be derived from the native protein or may be heterologous, and such sequences may be immunoreactive and/or antigenic.

As used herein, an "immunogenic portion" of a human prostate tumor protein is a portion that is capable of eliciting an immune response in a patient inflicted with prostate cancer and as such binds to antibodies present within sera from a prostate cancer patient. Immunogenic portions of the proteins described herein may thus be identified in antibody binding assays. Such assays may generally be performed using any of a variety of means known to those of ordinary skill in the art, as described, for example, in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1988. For example, a polypeptide may be immobilized on a solid support (as described below) and contacted with patient sera to allow binding of antibodies within the sera to the immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, ¹²⁵I-labeled Protein A. Alternatively, a polypeptide may be used to generate monoclonal and polyclonal antibodies for use in detection of the polypeptide in blood or other fluids of prostate cancer patients.

The compositions and methods of the present invention also encompass variants of the above polypeptides and DNA molecules. A polypeptide "variant," as used herein, is a polypeptide that differs from the recited polypeptide only in conservative substitutions and/or modifications, such that the therapeutic, antigenic and/or immunogenic properties of the polypeptide are retained. Polypeptide variants preferably exhibit at least about 70%, more preferably at least about 90% and most preferably at least about 95% identity to the identified polypeptides. For prostate tumor polypeptides with immunoreactive properties, variants may, alternatively, be identified by modifying the amino acid sequence of one of the above polypeptides, and evaluating the immunoreactivity of the modified polypeptide. For prostate tumor polypeptides useful for the generation of diagnostic binding agents, a variant may be identified by evaluating a modified polypeptide for the ability to

suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, CA), and may be operated according to the manufacturer's instructions.

Alternatively, any of the above polypeptides may be produced recombinantly by inserting a DNA sequence that encodes the polypeptide into an expression vector and expressing the protein in an appropriate host. Any of a variety of expression vectors known to those of ordinary skill in the art may be employed to express recombinant polypeptides of this invention. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast and higher eukaryotic cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line, such as CHO cells. The DNA sequences expressed in this manner may encode naturally occurring polypeptides, portions of naturally occurring polypeptides, or other variants thereof.

In general, regardless of the method of preparation, the polypeptides disclosed herein are prepared in substantially pure form (*i.e.*, the polypeptides are homogenous as determined by amino acid composition and primary sequence analysis). Preferably, the polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. In certain embodiments, described in more detail below, the substantially pure polypeptides are incorporated into pharmaceutical compositions or vaccines for use in one or more of the methods disclosed herein.

In a related aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, a polypeptide of the present invention and a known prostate antigen, together with variants of such fusion proteins. The fusion proteins of the present invention may also include a linker peptide between the first and second polypeptides.

A DNA sequence encoding a fusion protein of the present invention is constructed using known recombinant DNA techniques to assemble separate DNA sequences encoding the first and second polypeptides into an appropriate expression vector. The 3' end of a DNA sequence encoding the first polypeptide is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide so that the reading frames

of the sequences are in phase to permit mRNA translation of the two DNA sequences into a single fusion protein that retains the biological activity of both the first and the second polypeptides.

A peptide linker sequence may be employed to separate the first and the second polypeptides by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may be from 1 to about 50 amino acids in length. Peptide sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

The ligated DNA sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of DNA are located only 5' to the DNA sequence encoding the first polypeptides. Similarly, stop codons require to end translation and transcription termination signals are only present 3' to the DNA sequence encoding the second polypeptide.

Polypeptides and/or fusion proteins of the present invention may be used to generate binding agents, such as antibodies or fragments thereof, that are capable of detecting metastatic human prostate tumors. Binding agents of the present invention may generally be prepared using methods known to those of ordinary skill in the art, including the representative procedures described herein. Binding agents are capable of differentiating between patients with and without prostate cancer, using the representative assays described

herein. In other words, antibodies or other binding agents raised against a prostate tumor protein, or a suitable portion thereof, will generate a signal indicating the presence of primary or metastatic prostate cancer in at least about 20% of patients afflicted with the disease, and will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without primary or metastatic prostate cancer. Suitable portions of such prostate tumor proteins are portions that are able to generate a binding agent that indicates the presence of primary or metastatic prostate cancer in substantially all (*i.e.*, at least about 80%, and preferably at least about 90%) of the patients for which prostate cancer would be indicated using the full length protein, and that indicate the absence of prostate cancer in substantially all of those samples that would be negative when tested with full length protein. The representative assays described below, such as the two-antibody sandwich assay, may generally be employed for evaluating the ability of a binding agent to detect metastatic human prostate tumors.

The ability of a polypeptide and/or fusion protein prepared as described herein to generate antibodies capable of detecting primary or metastatic human prostate tumors may generally be evaluated by raising one or more antibodies against the polypeptide (using, for example, a representative method described herein) and determining the ability of such antibodies to detect such tumors in patients. This determination may be made by assaying biological samples from patients with and without primary or metastatic prostate cancer for the presence of a polypeptide that binds to the generated antibodies. Such test assays may be performed, for example, using a representative procedure described below. Polypeptides that generate antibodies capable of detecting at least 20% of primary or metastatic prostate tumors by such procedures are considered to be useful in assays for detecting primary or metastatic human prostate tumors. Polypeptide specific antibodies may be used alone or in combination to improve sensitivity.

Polypeptides and/or fusion proteins capable of detecting primary or metastatic human prostate tumors may be used as markers for diagnosing prostate cancer or for monitoring disease progression in patients. In one embodiment, prostate cancer in a patient may be diagnosed by evaluating a biological sample obtained from the patient for the level of

one or more of the above polypeptides, relative to a predetermined cut-off value. As used herein, suitable "biological samples" include blood, sera, urine and/or prostate secretions.

The level of one or more of the above polypeptides may be evaluated using any binding agent specific for the polypeptide(s). A "binding agent," in the context of this invention, is any agent (such as a compound or a cell) that binds to a polypeptide as described above. As used herein, "binding" refers to a noncovalent association between two separate molecules (each of which may be free (*i.e.*, in solution) or present on the surface of a cell or a solid support), such that a "complex" is formed. Such a complex may be free or immobilized (either covalently or noncovalently) on a support material. The ability to bind may generally be evaluated by determining a binding constant for the formation of the complex. The binding constant is the value obtained when the concentration of the complex is divided by the product of the component concentrations. In general, two compounds are said to "bind" in the context of the present invention when the binding constant for complex formation exceeds about 10^3 L/mol. The binding constant may be determined using methods well known to those of ordinary skill in the art.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome with or without a peptide component, an RNA molecule or a peptide. In a preferred embodiment, the binding partner is an antibody, or a fragment thereof. Such antibodies may be polyclonal, or monoclonal. In addition, the antibodies may be single chain, chimeric, CDR-grafted or humanized. Antibodies may be prepared by the methods described herein and by other methods well known to those of skill in the art.

There are a variety of assay formats known to those of ordinary skill in the art for using a binding partner to detect polypeptide markers in a sample. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In a preferred embodiment, the assay involves the use of binding partner immobilized on a solid support to bind to and remove the polypeptide from the remainder of the sample. The bound polypeptide may then be detected using a second binding partner that contains a reporter group. Suitable second binding partners include antibodies that bind to the binding partner polypeptide complex. Alternatively, a competitive assay may be utilized, in which a

polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding partner after incubation of the binding partner with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding partner is indicative of the reactivity of the sample with the immobilized binding partner.

The solid support may be any material known to those of ordinary skill in the art to which the antigen may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the antigen and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10 μ g, and preferably about 100 ng to about 1 μ g, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (*see, e.g.*, Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is a two-antibody sandwich assay. This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that polypeptides within the sample are allowed to bind to the immobilized antibody. Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a second antibody (containing a reporter group) capable of binding to a different site on the polypeptide is added. The amount of second antibody that remains bound to the solid support is then determined using a method appropriate for the specific reporter group.

More specifically, once the antibody is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20™ (Sigma Chemical Co., St. Louis, MO). The immobilized antibody is then incubated with the sample, and polypeptide is allowed to bind to the antibody. The sample may be diluted with a suitable diluent, such as phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is that period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with prostate cancer. Preferably, the contact time is sufficient to achieve a level of binding that is at least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. The second antibody, which contains a reporter group, may then be added to the solid support. Preferred reporter groups include enzymes (such as horseradish peroxidase), substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. The conjugation of antibody to reporter group may be achieved using standard methods known to those of ordinary skill in the art.

The second antibody is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound polypeptide. An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound second antibody is then removed and bound second antibody is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of prostate cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value is the average mean signal obtained when the immobilized antibody is incubated with samples from patients without prostate cancer. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for prostate cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, p. 106-7. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (*i.e.*, sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (*i.e.*, the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for prostate cancer.

In a related embodiment, the assay is performed in a flow-through or strip test format, wherein the antibody is immobilized on a membrane, such as nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized antibody as the sample passes through the membrane. A second, labeled antibody then binds to the antibody-polypeptide complex as a solution containing the second antibody flows through the membrane. The detection of bound second antibody may then be performed as described above. In the strip test format, one end of the membrane to which antibody is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing second antibody and to the area of immobilized antibody. Concentration of second antibody at the area of immobilized antibody indicates the presence of prostate cancer. Typically, the concentration of second antibody at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of antibody immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of polypeptide that would be sufficient to generate a positive signal in the two-antibody sandwich assay, in the format discussed above. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1 μ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use with the antigens or antibodies of the present invention. The above descriptions are intended to be exemplary only.

In another embodiment, the above polypeptides may be used as markers for the progression of prostate cancer. In this embodiment, assays as described above for the diagnosis of prostate cancer may be performed over time, and the change in the level of reactive polypeptide(s) evaluated. For example, the assays may be performed every 24-72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, prostate cancer is progressing in those patients in whom the level of polypeptide detected by the binding agent increases over time. In contrast, prostate cancer is not progressing when the level of reactive polypeptide either remains constant or decreases with time.

Antibodies for use in the above methods may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In one such technique, an immunogen comprising the antigenic polypeptide is initially injected into any of a wide variety of mammals (e.g., mice, rats, rabbits, sheep and goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for the antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (i.e., reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield,

such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

Monoclonal antibodies of the present invention may also be used as therapeutic reagents, to diminish or eliminate prostate tumors. The antibodies may be used on their own (for instance, to inhibit metastases) or coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides, differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed antiviral protein.

A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the

catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spitler), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers which provide multiple sites for attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.). A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing

nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

A variety of routes of administration for the antibodies and immunoconjugates may be used. Typically, administration will be intravenous, intramuscular, subcutaneous or in the bed of a resected tumor. It will be evident that the precise dose of the antibody/immunoconjugate will vary depending upon the antibody used, the antigen density on the tumor, and the rate of clearance of the antibody.

Diagnostic reagents of the present invention may also comprise DNA sequences encoding one or more of the above polypeptides, or one or more portions thereof. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify prostate tumor-specific cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for a DNA molecule encoding a prostate tumor protein of the present invention. The presence of the amplified cDNA is then detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes specific for a DNA molecule encoding a prostate tumor protein of the present invention may be used in a hybridization assay to detect the presence of an inventive polypeptide in a biological sample.

As used herein, the term "oligonucleotide primer/probe specific for a DNA molecule" means an oligonucleotide sequence that has at least about 80%, preferably at least about 90% and more preferably at least about 95%, identity to the DNA molecule in question. Oligonucleotide primers and/or probes which may be usefully employed in the inventive diagnostic methods preferably have at least about 10-40 nucleotides. In a preferred embodiment, the oligonucleotide primers comprise at least about 10 contiguous nucleotides of a DNA molecule having a sequence selected from SEQ ID Nos: 1-107, 109-111, 115-171, 173-175, 177 and 179-224. Preferably, oligonucleotide probes for use in the inventive diagnostic methods comprise at least about 15 contiguous oligonucleotides of a DNA molecule having a sequence provided in SEQ ID Nos: 1-107, 109-111, 115-171, 173-175, 177 and 179-224. Techniques for both PCR based assays and hybridization assays are well known in the art (see, for example, Mullis et al. *Ibid*; Ehrlich, *Ibid*). Primers or probes may

thus be used to detect prostate tumor-specific sequences in biological samples, including blood, semen, prostate tissue and/or prostate tumor tissue.

Polypeptides of the present invention that comprise an immunogenic portion of a prostate tumor protein may also be used for immunotherapy of prostate cancer, wherein the polypeptide stimulates the patient's own immune response to prostate tumor cells. In further aspects, the present invention provides methods for using one or more of the immunoreactive polypeptides encoded by a DNA molecule having a sequence provided in SEQ ID NO: 1-107, 109-111, 115-171, 173-175, 177 and 179-224 (or DNA encoding such polypeptides) for immunotherapy of prostate cancer in a patient. As used herein, a "patient" refers to any warm-blooded animal, preferably a human. A patient may be afflicted with a disease, or may be free of detectable disease. Accordingly, the above immunoreactive polypeptides may be used to treat prostate cancer or to inhibit the development of prostate cancer. The polypeptides may be administered either prior to or following surgical removal of primary tumors and/or treatment by administration of radiotherapy and conventional chemotherapeutic drugs.

In these aspects, the polypeptide is generally present within a pharmaceutical composition and/or a vaccine. Pharmaceutical compositions may comprise one or more polypeptides, each of which may contain one or more of the above sequences (or variants thereof), and a physiologically acceptable carrier. The vaccines may comprise one or more of such polypeptides and a non-specific immune response enhancer, such as an adjuvant, biodegradable microsphere (*e.g.*, polylactic galactide) or a liposome (into which the polypeptide is incorporated). Pharmaceutical compositions and vaccines may also contain other epitopes of prostate tumor antigens, either incorporated into a combination polypeptide (*i.e.*, a single polypeptide that contains multiple epitopes) or present within a separate polypeptide.

Alternatively, a pharmaceutical composition or vaccine may contain DNA encoding one or more of the above polypeptides, such that the polypeptide is generated *in situ*. In such pharmaceutical compositions and vaccines, the DNA may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacteria and viral expression systems. Appropriate nucleic acid

expression systems contain the necessary DNA sequences for expression in the patient (such as a suitable promoter). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an epitope of a prostate cell antigen on its cell surface. In a preferred embodiment, the DNA may be introduced using a viral expression system (e.g., vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective), replication competent virus. Suitable systems are disclosed, for example, in Fisher-Hoch et al., *PNAS* 86:317-321, 1989; Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld et al., *Science* 252:431-434, 1991; Kolls et al., *PNAS* 91:215-219, 1994; Kass-Eisler et al., *PNAS* 90:11498-11502, 1993; Guzman et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993. Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. The DNA may also be "naked," as described, for example, in published PCT application WO 90/11092, and Ulmer et al., *Science* 259:1745-1749, 1993, reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

Routes and frequency of administration, as well as dosage, will vary from individual to individual and may parallel those currently being used in immunotherapy of other diseases. In general, the pharmaceutical compositions and vaccines may be administered by injection (e.g., intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (e.g., by aspiration) or orally. Between 1 and 10 doses may be administered over a 3-24 week period. Preferably, 4 doses are administered, at an interval of 3 months, and booster administrations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of polypeptide or DNA that is effective to raise an immune response (cellular and/or humoral) against prostate tumor cells in a treated patient. A suitable immune response is at least 10-50% above the basal (i.e., untreated) level. In general, the amount of polypeptide present in a dose (or produced *in situ* by the DNA in a dose) ranges from about 1 pg to about 100 mg per kg of host, typically from

about 10 pg to about 1 mg, and preferably from about 100 pg to about 1 μ g. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.01 mL to about 5 mL.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a lipid, a wax and/or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and/or magnesium carbonate, may be employed. Biodegradable microspheres (e.g., polylactic glycolide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

Any of a variety of non-specific immune response enhancers may be employed in the vaccines of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a nonspecific stimulator of immune response, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis*. Such adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI) and Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ).

Polypeptides disclosed herein may also be employed in *ex vivo* treatment of prostate cancer. For example, cells of the immune system, such as T cells, may be isolated from the peripheral blood of a patient, using a commercially available cell separation system, such as CellPro Incorporated's (Bothell, WA) CEPRATE™ system (see U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). The separated cells are stimulated with one or more of the immunoreactive polypeptides contained within a delivery vehicle, such as a microsphere, to provide antigen-specific T cells. The population of tumor antigen-specific T cells is then expanded using standard techniques and the cells are administered back to the patient.

The following Examples are offered by way of illustration and not by way of limitation.

EXAMPLES

EXAMPLE 1

ISOLATION AND CHARACTERIZATION OF PROSTATE TUMOR POLYPEPTIDES

This Example describes the isolation of prostate tumor polypeptides from a prostate tumor cDNA library.

A human prostate tumor cDNA expression library was constructed from prostate tumor poly A⁺ RNA using a Superscript Plasmid System for cDNA Synthesis and Plasmid Cloning kit (BRL Life Technologies, Gaithersburg, MD 20897) following the manufacturer's protocol. Specifically, prostate tumor tissues were homogenized with polytron (Kinematica, Switzerland) and total RNA was extracted using Trizol reagent (BRL Life Technologies) as directed by the manufacturer. The poly A⁺ RNA was then purified using a Qiagen oligotex spin column mRNA purification kit (Qiagen, Santa Clarita, CA 91355) according to the manufacturer's protocol. First-strand cDNA was synthesized using the NotI/Oligo-dT18 primer. Double-stranded cDNA was synthesized, ligated with EcoRI/BAXI adaptors (Invitrogen, San Diego, CA) and digested with NotI. Following size fractionation with Chroma Spin-1000 columns (Clontech, Palo Alto, CA 94303), the cDNA was ligated into the EcoRI/NotI site of pCDNA3.1 (Invitrogen) and transformed into ElectroMax *E. coli* DH10B cells (BRL Life Technologies) by electroporation.

Using the same procedure, a normal human pancreas cDNA expression library was prepared from a pool of six tissue specimens (Clontech). The cDNA libraries were characterized by determining the number of independent colonies, the percentage of clones that carried insert, the average insert size and by sequence analysis. The prostate tumor library contained 1.64×10^7 independent colonies, with 70% of clones having an insert and the average insert size being 1745 base pairs. The normal pancreas cDNA library contained 3.3×10^6 independent colonies, with 69% of clones having inserts and the average insert size

being 1120 base pairs. For both libraries, sequence analysis showed that the majority of clones had a full length cDNA sequence and were synthesized from mRNA, with minimal rRNA and mitochondrial DNA contamination.

cDNA library subtraction was performed using the above prostate tumor and normal pancreas cDNA libraries, as described by Hara et al. (*Blood*, 84:189-199, 1994) with some modifications. Specifically, a prostate tumor-specific subtracted cDNA library was generated as follows. Normal pancreas cDNA library (70 µg) was digested with EcoRI, NotI, and SfuI, followed by a filling-in reaction with DNA polymerase Klenow fragment. After phenol-chloroform extraction and ethanol precipitation, the DNA was dissolved in 100 µl of H₂O, heat-denatured and mixed with 100 µl (100 µg) of Photoprobe biotin (Vector Laboratories, Burlingame, CA). As recommended by the manufacturer, the resulting mixture was irradiated with a 270 W sunlamp on ice for 20 minutes. Additional Photoprobe biotin (50 µl) was added and the biotinylation reaction was repeated. After extraction with butanol five times, the DNA was ethanol-precipitated and dissolved in 23 µl H₂O to form the driver DNA.

To form the tracer DNA, 10 µg prostate tumor cDNA library was digested with BamHI and XhoI, phenol chloroform extracted and passed through Chroma spin-400 columns (Clontech). Following ethanol precipitation, the tracer DNA was dissolved in 5 µl H₂O. Tracer DNA was mixed with 15 µl driver DNA and 20 µl of 2 x hybridization buffer (1.5 M NaCl/10 mM EDTA/50 mM HEPES pH 7.5/0.2% sodium dodecyl sulfate), overlaid with mineral oil, and heat-denatured completely. The sample was immediately transferred into a 68 °C water bath and incubated for 20 hours (long hybridization [LH]). The reaction mixture was then subjected to a streptavidin treatment followed by phenol/chloroform extraction. This process was repeated three more times. Subtracted DNA was precipitated, dissolved in 12 µl H₂O, mixed with 8 µl driver DNA and 20 µl of 2 x hybridization buffer, and subjected to a hybridization at 68 °C for 2 hours (short hybridization [SH]). After removal of biotinylated double-stranded DNA, subtracted cDNA was ligated into BamHI/XhoI site of chloramphenicol resistant pBCSK⁺ (Stratagene, La Jolla, CA 92037) and transformed into ElectroMax *E. coli* DH10B cells by electroporation to generate a prostate tumor specific subtracted cDNA library (prostate subtraction 1).

To analyze the subtracted cDNA library, plasmid DNA was prepared from 100 independent clones, randomly picked from the subtracted prostate tumor specific library and grouped based on insert size. Representative cDNA clones were further characterized by DNA sequencing with a Perkin Elmer/Applied Biosystems Division Automated Sequencer Model 373A (Foster City, CA). Six cDNA clones, hereinafter referred to as F1-13, F1-12, F1-16, H1-1, H1-9 and H1-4, were shown to be abundant in the subtracted prostate-specific cDNA library. The determined 3' and 5' cDNA sequences for F1-12 are provided in SEQ ID NO: 2 and 3, respectively, with determined 3' cDNA sequences for F1-13, F1-16, H1-1, H1-9 and H1-4 being provided in SEQ ID No: 1 and 4-7, respectively.

The cDNA sequences for the isolated clones were compared to known sequences in the gene bank using the EMBL and GenBank databases (release 96). Four of the prostate tumor cDNA clones, F1-13, F1-16, H1-1, and H1-4, were determined to encode the following previously identified proteins: prostate specific antigen (PSA), human glandular kallikrein, human tumor expression enhanced gene, and mitochondria cytochrome C oxidase subunit II. H1-9 was found to be identical to a previously identified human autonomously replicating sequence. No significant homologies to the cDNA sequence for F1-12 were found.

Subsequent studies led to the isolation of a full-length cDNA sequence for F1-12. This sequence is provided in SEQ ID NO: 107, with the corresponding predicted amino acid sequence being provided in SEQ ID NO: 108.

To clone less abundant prostate tumor specific genes, cDNA library subtraction was performed by subtracting the prostate tumor cDNA library described above with the normal pancreas cDNA library and with the three most abundant genes in the previously subtracted prostate tumor specific cDNA library: human glandular kallikrein, prostate specific antigen (PSA), and mitochondria cytochrome C oxidase subunit II. Specifically, 1 μ g each of human glandular kallikrein, PSA and mitochondria cytochrome C oxidase subunit II cDNAs in pCDNA3.1 were added to the driver DNA and subtraction was performed as described above to provide a second subtracted cDNA library hereinafter referred to as the "subtracted prostate tumor specific cDNA library with spike".

Twenty-two cDNA clones were isolated from the subtracted prostate tumor specific cDNA library with spike. The determined 3' and 5' cDNA sequences for the clones referred to as J1-17, L1-12, N1-1862, J1-13, J1-19, J1-25, J1-24, K1-58, K1-63, L1-4 and L1-14 are provided in SEQ ID Nos: 8-9, 10-11, 12-13, 14-15, 16-17, 18-19, 20-21, 22-23, 24-25, 26-27 and 28-29, respectively. The determined 3' cDNA sequences for the clones referred to as J1-12, J1-16, J1-21, K1-48, K1-55, L1-2, L1-6, N1-1858, N1-1860, N1-1861, N1-1864 are provided in SEQ ID Nos: 30-40, respectively. Comparison of these sequences with those in the gene bank as described above, revealed no significant homologies to three of the five most abundant DNA species, (J1-17, L1-12 and N1-1862; SEQ ID Nos: 8-9, 10-11 and 12-13, respectively). Of the remaining two most abundant species, one (J1-12; SEQ ID NO:30) was found to be identical to the previously identified human pulmonary surfactant-associated protein, and the other (K1-48; SEQ ID NO:33) was determined to have some homology to *R. norvegicus* mRNA for 2-arylpropionyl-CoA epimerase. Of the 17 less abundant cDNA clones isolated from the subtracted prostate tumor specific cDNA library with spike, four (J1-16, K1-55, L1-6 and N1-1864; SEQ ID Nos:31, 34, 36 and 40, respectively) were found to be identical to previously identified sequences, two (J1-21 and N1-1860; SEQ ID Nos: 32 and 38, respectively) were found to show some homology to non-human sequences, and two (L1-2 and N1-1861; SEQ ID Nos: 35 and 39, respectively) were found to show some homology to known human sequences. No significant homologies were found to the polypeptides J1-13, J1-19, J1-24, J1-25, K1-58, K1-63, L1-4, L1-14 (SEQ ID Nos: 14-15, 16-17, 20-21, 18-19, 22-23, 24-25, 26-27, 28-29, respectively).

Subsequent studies led to the isolation of full length cDNA sequences for J1-17, L1-12 and N1-1862 (SEQ ID NOS: 109-111, respectively). The corresponding predicted amino acid sequences are provided in SEQ ID NOS: 112-114.

In a further experiment, four additional clones were identified by subtracting a prostate tumor cDNA library with normal prostate cDNA prepared from a pool of three normal prostate poly A+ RNA (prostate subtraction 2). The determined cDNA sequences for these clones, hereinafter referred to as U1-3064, U1-3065, V1-3692 and 1A-3905, are provided in SEQ ID NO: 69-72, respectively. Comparison of the determined sequences with those in the gene bank revealed no significant homologies to U1-3065.

A second subtraction with spike (prostate subtraction spike 2) was performed by subtracting a prostate tumor specific cDNA library with spike with normal pancreas cDNA library and further spiked with PSA, J1-17, pulmonary surfactant-associated protein, mitochondrial DNA, cytochrome c oxidase subunit II, N1-1862, autonomously replicating sequence, L1-12 and tumor expression enhanced gene. Four additional clones, hereinafter referred to as V1-3686, R1-2330, 1B-3976 and V1-3679, were isolated. The determined cDNA sequences for these clones are provided in SEQ ID NO:73-76, respectively. Comparison of these sequences with those in the gene bank revealed no significant homologies to V1-3686 and R1-2330.

Further analysis of the three prostate subtractions described above (prostate subtraction 2, subtracted prostate tumor specific cDNA library with spike, and prostate subtraction spike 2) resulted in the identification of sixteen additional clones, referred to as 1G-4736, 1G-4738, 1G-4741, 1G-4744, 1G-4734, 1H-4774, 1H-4781, 1H-4785, 1H-4787, 1H-4796, 1I-4810, 1I-4811, 1J-4876, 1K-4884 and 1K-4896. The determined cDNA sequences for these clones are provided in SEQ ID NOS: 77-92, respectively. Comparison of these sequences with those in the gene bank as described above, revealed no significant homologies to 1G-4741, 1G-4734, 1I-4807, 1J-4876 and 1K-4896 (SEQ ID NOS: 79, 81, 87, 90 and 92, respectively). Further analysis of the isolated clones led to the determination of extended cDNA sequences for 1G-4736, 1G-4738, 1G-4741, 1G-4744, 1H-4774, 1H-4781, 1H-4785, 1H-4787, 1H-4796, 1I-4807, 1J-4876, 1K-4884 and 1K-4896, provided in SEQ ID NOS: 179-188 and 191-193, respectively, and to the determination of additional partial cDNA sequences for 1I-4810 and 1I-4811, provided in SEQ ID NOS: 189 and 190, respectively.

An additional subtraction was performed by subtracting a normal prostate cDNA library with normal pancreas cDNA (prostate subtraction 3). This led to the identification of six additional clones referred to as 1G-4761, 1G-4762, 1H-4766, 1H-4770, 1H-4771 and 1H-4772 (SEQ ID NOS: 93-98). Comparison of these sequences with those in the gene bank revealed no significant homologies to 1G-4761 and 1H-4771 (SEQ ID NOS: 93 and 97, respectively). Further analysis of the isolated clones led to the determination of extended cDNA sequences for 1G-4761, 1G-4762, 1H-4766 and 1H-4772 provided in SEQ

ID NOS: 194-196 and 199, respectively, and to the determination of additional partial cDNA sequences for 1H-4770 and 1H-4771, provided in SEQ ID NOS: 197 and 198, respectively.

Subtraction of a prostate tumor cDNA library, prepared from a pool of polyA-RNA from three prostate cancer patients, with a normal pancreas cDNA library (prostate subtraction 4) led to the identification of eight clones, referred to as 1D-4297, 1D-4309, 1D.1-4278, 1D-4288, 1D-4283, 1D-4304, 1D-4296 and 1D-4280 (SEQ ID NOS: 99-107). These sequences were compared to those in the gene bank as described above. No significant homologies were found to 1D-4283 and 1D-4304 (SEQ ID NOS: 103 and 104, respectively). Further analysis of the isolated clones led to the determination of extended cDNA sequences for 1D-4309, 1D.1-4278, 1D-4288, 1D-4283, 1D-4304, 1D-4296 and 1D-4280, provided in SEQ ID NOS: 200-206, respectively.

cDNA clones isolated in prostate subtraction 1 and prostate subtraction 2, described above, were colony PCR amplified and their mRNA expression levels in prostate tumor, normal prostate and in various other normal tissues were determined using microarray technology (Synteni, Palo Alto, CA). Briefly, the PCR amplification products were dotted onto slides in an array format, with each product occupying a unique location in the array. mRNA was extracted from the tissue sample to be tested, reverse transcribed, and fluorescent-labeled cDNA probes were generated. The microarrays were probed with the labeled cDNA probes, the slides scanned and fluorescence intensity was measured. This intensity correlates with the hybridization intensity. Two novel clones (referred to as P509S and P510S) were found to be over-expressed in prostate tumor and normal prostate and expressed at low levels in all other normal tissues tested (liver, pancreas, skin, bone marrow, brain, breast, adrenal gland, bladder, testes, salivary gland, large intestine, kidney, ovary, lung, spinal cord, skeletal muscle and colon). The determined cDNA sequences for P509S and P510S are provided in SEQ ID NO: 223 and 224, respectively. Comparison of these sequences with those in the gene bank as described above, revealed some homology to previously identified ESTs.

EXAMPLE 2

DETERMINATION OF TISSUE SPECIFICITY OF PROSTATE TUMOR POLYPEPTIDES

Using gene specific primers, mRNA expression levels for the representative prostate tumor polypeptides F1-16, H1-1, J1-17, L1-12, F1-12 and N1-1862 were examined in a variety of normal and tumor tissues using RT-PCR.

Briefly, total RNA was extracted from a variety of normal and tumor tissues using Trizol reagent as described above. First strand synthesis was carried out using 1-2 μ g of total RNA with SuperScript II reverse transcriptase (BRL Life Technologies) at 42 °C for one hour. The cDNA was then amplified by PCR with gene-specific primers. To ensure the semi-quantitative nature of the RT-PCR, β -actin was used as an internal control for each of the tissues examined. First, serial dilutions of the first strand cDNAs were prepared and RT-PCR assays were performed using β -actin specific primers. A dilution was then chosen that enabled the linear range amplification of the β -actin template and which was sensitive enough to reflect the differences in the initial copy numbers. Using these conditions, the β -actin levels were determined for each reverse transcription reaction from each tissue. DNA contamination was minimized by DNase treatment and by assuring a negative PCR result when using first strand cDNA that was prepared without adding reverse transcriptase.

mRNA Expression levels were examined in four different types of tumor tissue (prostate tumor from 2 patients, breast tumor from 3 patients, colon tumor, lung tumor), and sixteen different normal tissues, including prostate, colon, kidney, liver, lung, ovary, pancreas, skeletal muscle, skin, stomach, testes, bone marrow and brain. F1-16 was found to be expressed at high levels in prostate tumor tissue, colon tumor and normal prostate, and at lower levels in normal liver, skin and testes, with expression being undetectable in the other tissues examined. H1-1 was found to be expressed at high levels in prostate tumor, lung tumor, breast tumor, normal prostate, normal colon and normal brain, at much lower levels in normal lung, pancreas, skeletal muscle, skin, small intestine, bone marrow, and was not detected in the other tissues tested. J1-17 and L1-12 appear to be specifically over-expressed in prostate, with both genes being expressed at high levels in prostate tumor and normal prostate but at low to undetectable levels in all the other tissues

examined. N1-1862 was found to be over-expressed in 60% of prostate tumors and detectable in normal colon and kidney. The RT-PCR results thus indicate that F1-16, H1-1, J1-17, N1-1862 and L1-12 are either prostate specific or are expressed at significantly elevated levels in prostate.

Further RT-PCR studies showed that F1-12 is over-expressed in 60% of prostate tumors, detectable in normal kidney but not detectable in all other tissues tested. Similarly, R1-2330 was shown to be over-expressed in 40% of prostate tumors, detectable in normal kidney and liver, but not detectable in all other tissues tested. U1-3064 was found to be over-expressed in 60% of prostate tumors, and also expressed in breast and colon tumors, but was not detectable in normal tissues.

RT-PCR characterization of R1-2330, U1-3064 and 1D-4279 showed that these three antigens are over-expressed in prostate and/or prostate tumors.

Northern analysis with four prostate tumors, two normal prostate samples, two BPH prostates, and normal colon, kidney, liver, lung, pancreas, skeletal muscle, brain, stomach, testes, small intestine and bone marrow, showed that L1-12 is over-expressed in prostate tumors and normal prostate, while being undetectable in other normal tissues tested. J1-17 was detected in two prostate tumors and not in the other tissues tested. N1-1862 was found to be over-expressed in three prostate tumors and to be expressed in normal prostate, colon and kidney, but not in other tissues tested. F1-12 was found to be highly expressed in two prostate tumors and to be undetectable in all other tissues tested.

The micro-array technology described above was used to determine the expression levels of representative antigens described herein in prostate tumor, breast tumor and the following normal tissues: prostate, liver, pancreas, skin, bone marrow, brain, breast, adrenal gland, bladder, testes, salivary gland, large intestine, kidney, ovary, lung, spinal cord, skeletal muscle and colon. L1-12 was found to be over-expressed in normal prostate and prostate tumor, with some expression being detected in normal skeletal muscle. Both J1-12 and F1-12 were found to be over-expressed in prostate tumor, with expression being lower or undetectable in all other tissues tested. N1-1862 was found to be expressed at high levels in prostate tumor and normal prostate, and at low levels in normal large intestine and normal colon, with expression being undetectable in all other tissues tested. R1-2330 was found to

be over-expressed in prostate tumor and normal prostate, and to be expressed at lower levels in all other tissues tested. 1D-4279 was found to be over-expressed in prostate tumor and normal prostate, expressed at lower levels in normal spinal cord, and to be undetectable in all other tissues tested.

Example 3

ISOLATION AND CHARACTERIZATION OF PROSTATE TUMOR POLYPEPTIDES BY PCR-BASED SUBTRACTON

A cDNA subtraction library, containing cDNA from normal prostate subtracted with ten other normal tissue cDNAs (brain, heart, kidney, liver, lung, ovary, placenta, skeletal muscle, spleen and thymus) and then submitted to a first round of PCR amplification, was purchased from Clontech. This library was subjected to a second round of PCR amplification, following the manufacturer's protocol. The resulting cDNA fragments were subcloned into the vector pT7 Blue T-vector (Novagen, Madison, WI) and transformed into XL-1 Blue MRF⁺ *E. coli* (Stratagene). DNA was isolated from independent clones and sequenced using a Perkin Elmer/Applied Biosystems Division Automated Sequencer Model 373A.

Fifty-nine positive clones were sequenced. Comparison of the DNA sequences of these clones with those in the gene bank, as described above, revealed no significant homologies to 25 of these clones, hereinafter referred to as P5, P8, P9, P18, P20, P30, P34, P36, P38, P39, P42, P49, P50, P53, P55, P60, P64, P65, P73, P75, P76, P79, and P84. The determined cDNA sequences for these clones are provided in SEQ ID NO:41-45, 47-52 and 54-65, respectively. P29, P47, P68, P80 and P82 (SEQ ID NO:46, 53 and 66-68, respectively) were found to show some degree of homology to previously identified DNA sequences. To the best of the inventors' knowledge, none of these sequences have been previously shown to be present in prostate.

Further studies using the PCR-based methodology described above resulted in the isolation of more than 180 additional clones, of which 23 clones were found to show no significant

homologies to known sequences. The determined cDNA sequences for these clones are provided in SEQ ID NO: 115-123, 127, 131, 137, 145, 147-151, 153, 156-158 and 160. Twenty-three clones (SEQ ID NO: 124-126, 128-130, 132-136, 138-144, 146, 152, 154, 155 and 159) were found to show some homology to previously identified ESTs. An additional ten clones (SEQ ID NO: 161-170) were found to have some degree of homology to known genes. An additional clone, referred to as P703, was found to have five splice variants. The determined DNA sequence for the variants referred to as DE1, DE13 and DE14 are provided in SEQ ID NOS: 171, 175 and 177, respectively, with the corresponding predicted amino acid sequences being provided in SEQ ID NO: 172, 176 and 178, respectively. The DNA sequences for the splice variants referred to as DE2 and DE6 are provided in SEQ ID NOS: 173 and 174, respectively.

mRNA Expression levels for representative clones in tumor tissues (prostate (n=5), breast (n=2), colon and lung) normal tissues (prostate (n=5), colon, kidney, liver, lung (n=2), ovary (n=2), skeletal muscle, skin, stomach, small intestine and brain), and activated and non-activated PBMC was determined by RT-PCT as described above. Expression was examined in one sample of each tissue type unless otherwise indicated.

P9 was found to be highly expressed in normal prostate and prostate tumor compared to all normal tissues tested except for normal colon which showed comparable expression. P20 was found to be highly expressed in normal prostate and prostate tumor, compared to all twelve normal tissues tested. A modest increase in expression of P20 in breast tumor (n=2), colon tumor and lung tumor was seen compared to all normal tissues except lung (1 of 2). Increased expression of P18 was found in normal prostate, prostate tumor and breast tumor compared to other normal tissues except lung and stomach. A modest increase in expression of P5 was observed in normal prostate compared to most other normal tissues. However, some elevated expression was seen in normal lung and PBMC. Elevated expression of P5 was also observed in prostate tumors (2 of 5), breast tumor and one lung tumor sample. For P30, similar expression levels were seen in normal prostate and prostate tumor, compared to six of twelve other normal tissues tested. Increased expression was seen in breast tumors, one lung tumor sample and one colon tumor sample, and also in normal PBMC. P29 was found to be over-expressed in prostate tumor (5 of 5) and normal prostate (5

of 5) compared to the majority of normal tissues. However, substantial expression of P29 was observed in normal colon and normal lung (2 of 2). P80 was found to be over-expressed in prostate tumor (5 of 5) and normal prostate (5 of 5) compared to all other normal tissues tested, with increased expression also being seen in colon tumor.

Further studies using the above methodology resulted in the isolation of twelve additional clones, hereinafter referred to as 10-d8, 10-h10, 11-c8, 7-g6, 8-b5, 8-b6, 8-d4, 8-d9, 8-g3, 8-h11, g-f12 and g-f3. The determined DNA sequences for 10-d8, 10-h10, 11-c8, 8-d4, 8-d9, 8-h11, g-f12 and g-f3 are provided in SEQ ID NO: 207, 208, 209, 216, 217, 220, 221 and 222, respectively. The determined forward and reverse DNA sequences for 7-g6, 8-b5, 8-b6 and 8-g3 are provided in SEQ ID NO: 210 and 211; 212 and 213; 214 and 215; and 218 and 219, respectively. Comparison of these sequences with those in the gene bank revealed no significant homologies to the sequences of 7-g6 and g-f3. The clones 10-d8, 11-c8 and 8-h11 were found to show some homology to previously isolated ESTs, while 10-h10, 8-b5, 8-b6, 8-d4, 8-d9, 8-g3 and g-f12 were found to show some homology to previously identified genes.

EXAMPLE 4

SYNTHESIS OF POLYPEPTIDES

Polypeptides may be synthesized on an Applied Biosystems 430A peptide synthesizer using Fmoc chemistry with HPTU (O-Benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate) activation. A Gly-Cys-Gly sequence may be attached to the amino terminus of the peptide to provide a method of conjugation, binding to an immobilized surface, or labeling of the peptide. Cleavage of the peptides from the solid support may be carried out using the following cleavage mixture: trifluoroacetic acid:ethanedithiol:thioanisole:water:phenol (40:1:2:2:3). After cleaving for 2 hours, the peptides may be precipitated in cold methyl-t-butyl-ether. The peptide pellets may then be dissolved in water containing 0.1% trifluoroacetic acid (TFA) and lyophilized prior to purification by C18 reverse phase HPLC. A gradient of 0%-60% acetonitrile (containing 0.1% TFA) in water (containing 0.1% TFA) may be used to elute the peptides. Following

lyophilization of the pure fractions, the peptides may be characterized using electrospray or other types of mass spectrometry and by amino acid analysis.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for the purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

34

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANTS: Xu, Jiangchun
Dillon, Davin C.

(ii) TITLE OF INVENTION: COMPOUNDS FOR IMMUNODIAGNOSIS OF PROSTATE CANCER
AND METHODS FOR THEIR USE

(iii) NUMBER OF SEQUENCES: 224

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: SEED and BERRY LLP
(B) STREET: 6300 Columbia Center, 701 Fifth Avenue
(C) CITY: Seattle
(D) STATE: WA
(E) COUNTRY: USA
(F) ZIP: 98104

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER
(B) FILING DATE: 23-FEB-1998
(C) CLASSIFICATION:

(vii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Maki, David J.
(B) REGISTRATION NUMBER: 31,392
(C) REFERENCE/DOCKET NUMBER: 210121.426C3

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (206) 622-4900
(B) TELEFAX: (206) 682-6031

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 814 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TTTTTTTTTT TTTTTCACAG TATAACAGCT GTTATTCTCT GTGAGTTCTA CTAGGAAATC

60

ATCAAATCTG	AGGGTTGTCT	GGAGGACTTC	AATACACCTC	CCCCCATAGT	GAATCAGCTT	120
CCAGGGGGTC	CAGTCCCTCT	CCTTACTTCA	TCCCCATCCC	ATGCCAAAGG	AAGACCCTCC	180
CTCCTTGGCT	CACAGCCTTC	TCTAGGCTTC	CCAGTGCCTC	CAGGACAGAG	TGGGTATATG	240
TTTCAGCTCC	ATCCTTGCTG	TGAGTGTCTG	CTGCGTTGTG	CCTCCAGCTT	CTGCTCAGTG	300
CTTCATGGAC	AGTGTCCAGC	ACATGTCACT	CTCCACTCTC	TCAGTGTGGA	TCCACTAGTT	360
CTAGAGCGGC	CGCCACCGCG	GTGGAGCTCC	AGCTTTTGTT	CCCTTTAGTG	AGGGTTAATT	420
GCGCGCTTGG	CGTAATCATG	GTCATAACTG	TTTCCTGTGT	GAAATTGTTA	TCCGCTCACA	480
ATTCCACACA	ACATACGAGC	CGGAAGCATA	AAGTGTAAG	CCTGGGGTGC	CTAATGAGTG	540
ANCTAACTCA	CATTAAATTG	GTTGCGCTCA	CTCNCCGCTT	TCCAGTCNGG	AAAAGTGTCT	600
TGCCAGCTGC	ATTAATGAAT	CGGCCAACGC	NCGGGGAAAA	GCGGTTTGCG	TTTTGGGGGC	660
TCTTCCGCTT	CTCGCTCACT	NANTCCTGCG	CTCGGTCNTT	CGGCTGCGGG	GAACGGTATC	720
ACTCCTCAAA	GGNGGTATTA	CGGTTATCCN	NAAATCNGGG	GATACCCNGG	AAAAAANTTT	780
AACAAAAGGG	CANCAAAGGG	CNGAAACGTA	AAAA			814

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 816 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ACAGAAATGT	TGGATGGTGG	AGCACCTTTC	TATACGACTT	ACAGGACAGC	AGATGGGGAA	60
TTCATGGCTG	TTGGAGCAAT	AGAACCCAG	TTCTACGAGC	TGCTGATCAA	AGGACTTGGA	120
CTAAAGTCTG	ATGAACCTCC	CAATCAGATG	AGCATGGATG	ATTGGCCAGA	AATGAAGAAG	180
AAGTTTGAG	ATGTATTTGC	AAAGAAGACG	AAGGCAGAGT	GGTGTCAAAT	CTTTGACGGC	240
ACAGATGCCT	GTGTGACTCC	GGTTCGACT	TTTGAGGAGG	TTGTTTCATCA	TGATCACAAC	300
AAGGAACGGG	GCTCGTTTAT	CACCACTGAG	GAGCAGGACG	TGAGCCCCCG	CCCTGCACCT	360
CTGCTGTTAA	ACACCCAGC	CATCCCTTCT	TTCAAAAGGG	ATCCACTAGT	TCTAGAAGCG	420
GCCGCCACCG	CGGTGGAGCT	CCAGCTTTTG	TTCCCTTTAG	TGAGGGTTAA	TTGCGCGCTT	480
GGCGTAATCA	TGGTCATAGC	TGTTTCTGTG	GTGAAATTGT	TATCCGCTCA	CAATTCCCCC	540
AACATACGAC	CCGGAACATA	AAGTGTTAAG	CCTGGGGTGC	CTAATGANTG	AGCTAACTCN	600
CATTAAATTG	GTTGCGCTCA	CTGCCCCTT	TCCAGTCGGG	AAAAGTGTCT	TGCCACTGCN	660
TTANTGAATC	NGCCACCCCC	CGGGAAAAGG	CGGTTGCNTT	TTGGGCCTCT	TCCGCTTTCC	720
TCGCTCATTG	ATCCTNGCNC	CCGGTCTTCG	GCTGCGGNGA	ACGGTTCACT	CCTCAAAGGC	780
GGTNTNCCGG	TTATCCCCAA	ACNNGGGATA	CCNGA			816

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 773 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CTTTTGAAAG	AAGGGATGGC	TGGGGTGTTT	AACAGCAGAG	GTGCAGGGCG	GGGGCTCAGC	60
TCCTGCTCCT	CACTGCTGAT	AAACGAGCCC	CGTTCCTTGT	TGTGATCATG	ATGAACAACC	120

TCCTCAAAAG	TCAGAACCGB	AGTCACACAG	GCATCTGTGC	CGTCAAAAGAT	TTGACACCAC	180
TCTTCCTTCG	TCTTCTTTTC	AAATACATCT	GCAAACTTCT	TCTTCATTTG	TGSCCAATCA	240
TCCATGCTCA	TCTGATTGGG	AAGTTCATCA	GACTTTAGTG	CANNTCCTTT	GATCAGCAGC	300
TGCTAGAACT	GGGTTCTAT	TGCTCCAACA	CCCATGAATT	CCCCATCTGC	TGCTCTGTAA	360
GTCTATAGA	AAGGTGCTCC	ACCATCCAAC	ATGTTCTGTG	CTCGAGGGGG	GGCCCGGTAC	420
CTAATTCGCC	CTATANTGAG	TCGTATTACG	CGCGCTCACT	GGCGTCTGTT	TTACAACGTT	480
GTGACTGGGA	AAACCCTGGG	CGTTACCAAC	TTAATCGCCT	TGCAGCACAT	CCCGCTTTGG	540
CCAGCTGGGC	GTAATANCGA	AAAGGCCCGG	ACCGATCGCG	CTTCCAACAG	TTCCGCACCT	600
GAATGGGNAA	ATGGGACCCC	CCTGTTACCG	CGCATTNAAC	CCCCGCNGGG	TTTNGTTGTT	660
ACCCCAACNT	NNACCGCTTA	CACTTTGCCA	GCGCCTTANC	GCCCCGCTCCG	TTTCNCCTTT	720
CTTCCTTGG	TTTCNCNCCN	CTTTCCTCCG	JGGTTTCCCG	ENTCAAACCC	CNA	773

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 823 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCTCCTGAGT	CCTACTGACC	TGTGCTTTCT	GGTGTGGAGT	CCAGGGCTGC	TAGGAAAAGG	60
AATGGGCAGA	CACAGGTGTA	TGCCAATGTT	TCTGAAATGG	GTATAATTTG	GTCCTCTCCT	120
TGGGAACACT	GCCTGTCTCT	GAAGACTTTC	CGCTCACTTT	CAGTGAGGAC	ACACACAAAG	180
AGGTGGGTGA	CCATGTTGTT	TGTGGGGTGC	AGAGATGGGA	GGGGTGGGGC	CCACCCTGGA	240
AGASTGGACA	GTGACACAAG	GTGGACACTC	TCTACAGATC	ACTGAGGATA	AGCTGGAGCC	300
ACAATGCATG	AGGCACACAC	ACAGCAAGGA	TGACNCTGTA	AACATAGCCC	ACGCTGTCCT	360
GNGGGCACTG	GGAAGCCTAN	ATNAGGCCGT	GAGCANAAAG	AAGGGGAGGA	TCCACTAGTI	420
CTANAGCGGC	CGCCACCGCG	GTGGANCTCC	ANCTTTTGTT	CCCTTTAGTG	AGGTTAATT	480
GCGCGCTTGG	CNTAATCATG	GTGATANTCN	TTTCCTGTGT	GAAATTGTTA	TCCGCTCACA	540
ATTCCACACA	ACATACGANC	CGGAAACATA	AANTGTAAAC	CTGGGGTGCC	TAATGANTGA	600
CTAACTCACA	TTAATTGCGT	TGCGCTCACT	GCCCCGCTTT	CAATCNGGAA	ACCTGCTCTG	660
CCNCTTGCA	TNATGAATCN	GCCAAACCCC	GGGGAAAAGC	GTTTGCGTTT	TGGGCGCTCT	720
TCCGCTTCCT	CNCTCANTTA	NTCCCTNCNC	TGGGTCAATT	CGGCTGCNGC	AAACCGGTTG	780
ACCNCCTCCA	AAGGGGTAT	TGCGTTTCC	CCNAATCCCG	GGANANCE		823

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 831 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TTTTTTTTTT	TTTTTACTGA	TAGATGCAAT	TTATTAACT	TTTCACATGT	GATAGCACAT	60
AGTTTTAAAT	GCATCCAAAG	TACTAACAAG	AACTCTAGCA	ATCAAGAATG	GCAGCATGTT	120
ATTTTATAAC	AATCAACAGC	TGTGGCTTTT	AAAATTTGCT	TTTATAAGA	TAATTTATAC	180
TGAAGTAAAT	CTAGCCATGC	TTTTAAAAAA	TGCTTTAGGT	CAGTCCAAGC	TTGGCACTTA	240

ACATTTGGCA	TAAACAATAA	TAAAACAATC	ACAATTTAAT	AAATAACAAA	TACAACATTG	300
TAGGCCATAA	TCATATACAG	TATAAGGAAA	AGCTGGTAGT	GTTGAGTAAG	CAGTTATTAG	360
AATAGAATAC	CTTGGCCTCT	ATGCAAATAT	GTCTAGACAC	TTTGATTAC	TCAGCCCTGA	420
CATTCAGTTT	TCAAAGTAGG	AGACAGGTTT	TACAGTATCA	TTTTACAGTT	TCCAACACAT	480
TGAAAACAAG	TAGAAAATGA	TGAGTTGATT	TTTATTAATG	CATTACATCC	TCAAGAGTTA	540
TCACCAACCC	CTCAGTTATA	AAAAATTTTC	AAGTTATATT	AGTCATATAA	CTTGGTGTGC	600
TTATTTTAAA	TTAGTGCTAA	ATGGATTAAG	TGAAGACAAC	AATGGTCCCC	TAATGTGATT	660
GATATTGGTC	ATTTTACCA	GCTTCTAAAT	CTNAACTTTC	AGGCTTTTGA	ACTGGAACA	720
TGNATNACAG	TGTTCCANAG	TTNCAACCTA	CTGGAACATT	ACAGTGTGCT	TGATTCAAAA	780
TGTTATTTTG	TTAAAAATTA	AATTTTAACC	TGGTGAAAAA	ATAATTTGAA	ATNA	834

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 818 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TTTTTTTTTT	TTTTTTTTTT	AAGACCCTCA	TCAATAGATG	GAGACATACA	GAAATAGTCA	60
AACCACATCT	ACAAAATGCC	AGTATCAGGC	GGCGGCTTCG	AAGCCAAAGT	GATGTTTGGA	120
TGTAAAGTGA	AATATTAGTT	GGCGGATGAA	GCAGATAGTG	AGGAAAGTTG	AGCCAATAAT	180
GACGTGAAGT	CCGTGGAAGC	CTGTGGCTAC	AAAAAATGTT	GAGCCGTAGA	TGCCGTCCGA	240
AATGGTGAAG	GGAGACTCGA	AGTACTCTGA	GGCTTGTAGG	AGGGTAAAAT	AGAGACCCAG	300
TAAAATTGTA	ATAAGCAGTG	CTTGAATTAT	TTGGTTTTCGG	TTGTTTTCTA	TTAGACTATG	360
GTGAGCTCAG	GTGATTGATA	CTCCTGATGC	GAGTAATACG	GATGTGTTTA	GGAGTGGGAC	420
TTCTAGGGGA	TTTAGCGGGG	TGATGCCTGT	TGGGGGCCAG	TGCCCTCCTA	GTTGGGGGGT	480
AGGGGCTAGG	CTGGAGTGTT	AAAAGGCTCA	GAAAAATCCT	GCGAAGAAAA	AAACTTCTGA	540
GGTAATAAAT	AGGATTATCC	CGTATCGAAG	GCCTTTTTGG	ACAGGTGGTG	TGTGGTGGCC	600
TTGGTATGTG	CTTTCTCGTG	TTACATCGCG	CCATCATTGG	TATATGGTTA	GTGTGTGGG	660
TTANTANGGC	CTANTATGAA	GAACTTTTGG	ANTGGAATTA	AATCAATNGC	TTGGCCGGAA	720
GTCATTANGA	NGGCTNAAAA	GGCCCTGTTA	NGGGTCTGGG	CTNGGTTTTA	CCCNACCCAT	780
GGAATNCNCC	CCCCGGACNA	NTGNATCCCT	ATTCTTAA			818

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 817 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TTTTTTTTTT	TTTTTTTTTT	TGGCTCTAGA	GGGGGTAGAG	GGGGTGCTAT	AGGGTAAATA	60
CGGGCCCTAT	TTCAAAGATT	TTTAGGGGAA	TTAATTCTAG	GACGATGGGT	ATGAAACTGT	120
GGTTTGCTCC	ACAGATTTCA	GAGCATTGAC	CGTAGTATAC	CCCCGGTCGT	GTAGCGGTGA	180
AAGTGCTTTG	GTTTAGACGT	CCGGGAATTG	CATCTGTTTT	TAAGCCTAAT	GTGGGGACAG	240
CTCATGAGTG	CAAGACGTCT	TGTGATGTAA	TTATTATACN	AATGGGGGCT	TCAATCGGGA	300

GTACTACTCG	ATTGTCAACG	TCAAGGASTC	GCAGGTCCGC	TGGTTCTAGG	AATAATGGGG	360
GAAATATGTA	GGAATTGAAG	ATTAATCCCG	CGTAGTCGGT	GTTCTCCTAG	GTTCAATAAC	420
ATTGGTGBCG	AATTGATTTG	ATGGTAAGGG	GAGGGATCGT	TGAACTCGTC	TGTTATGTAA	480
AGGATNCCTT	NGGGATGGGA	AGGCNATNAA	GGACTANGGA	TNAATGGCGG	CCANGATATT	540
TCAAACNGTC	TCTANTTCCT	GAAACGTCTG	AAATGTTAAT	AANAATTAAN	TTTNGTTATT	600
GAAATNTTNG	GAAAAGGGCT	TACAGGAATA	GAAACCAAAT	ANGAAAANTA	ATNNTAANGG	660
CNTTATCNTN	AAAGGTNATA	ACCNCTCCTA	TNATCCCAAC	CAATNGNATT	CCCCACNCNN	720
ACNATTGGAT	NCCCCANTTC	CANAAANGGG	CNCCCCCGGG	TGNANNCCNC	CTTTTGTTCG	780
CTTNANTGAN	GGTTATTTCN	CCCTNGCNTT	ATCANCC			817

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 799 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CATTTCCGGG	TTTACTTTCT	AAGGAAAGCC	GAGCGGAAGC	TGCTAACGTG	GGAATCGGTG	60
CATAAGGAGA	ACTTTCTGCT	GGCACGCGCT	AGGACACAAGC	GGGAGAGCGA	CTCCGAGCGT	120
CTGAAGCGCA	CGTCCCAGAA	GGTGGACTTG	GCCTGAAAC	AGCTGGGACA	CATCCGCGAG	180
TACGAACAAC	GCCTGAAAGT	GCTGGAGCGG	GAGGTCCAGC	AGTGTAGCCG	CGTCTCTGGG	240
TGGGTGGGCG	ANGCCTGANC	CGCTCTGCTT	TGCTGCCCCC	ANGTGGGCGG	CCATCCCCCTG	300
ACCTGCCTGG	GTCCAAACAC	TGAGCCCTGC	TGGGCGACTT	CAAGGANAAC	CCCCACANGG	360
GGATTTTGCT	CCTANANTAA	GGCTCATCTG	GGCTTCGGCC	CCCCCACCTG	GTTGGCCTTG	420
TCTTTGANGT	GAGCCCCATG	TCCATCTGGG	CCACTGTCNG	GACCACCTTT	NGGGAGTGTT	480
CTCCTTACAA	CCACANNATG	CCCGGCTCCT	CCCGGAAACC	ANTCCCANCC	TGNGAAGGAT	540
CAAGNCCTGN	ATCCACTNNT	NCTANAACCG	GCCNCCNCCG	CNGTGGAACC	CNCCTTNTGT	600
TCCTTTTCNT	TNAGGGTTAA	TNNGGCTTG	GCCTTNCCAN	NGTCCTNCNC	NTTTTCCNNT	660
GTTNAAATTG	TTANGCNCNC	NCCNNTCCCN	CNNNNNNAN	CCCGACCCNN	ANNTTNNANN	720
NCCTGGGGGT	NCCNNGAT	TGACCCNCC	NCCCTNTANT	TGCTTTNGGG	NNCNNTGCCG	780
CTTTCCCTCT	NGGGANNCC					799

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 801 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ACGCGCTTGAT	CCTCCCAAGG	TGGGACTGCT	TCTGGGAGGA	GCCGGGCATG	CTGTGCTTTG	60
TAAGATGAC	ACTCCCAAAG	GTGGTCTTGA	CASTGGCCCA	GATGGACATG	GGGCTCACCT	120
CAAGGACAAG	GCCACCAAGT	GCGGGGGCGG	AAGCCCAAT	GATCCTTACT	CTATGAGCAA	180
AATCCCTCTG	GGGGGCTTCT	CCTTGAAGTC	CGCCANCAGG	GCTCAGTCTT	TGSACCCANG	240
CAGGTGATCC	GGTTGTNGNC	CAACTGGGGG	CCNCAACGCA	AAANGGCNCA	GGGCTCTNGN	300
CACCCATGCC	ANGATGGGGG	TACACTNCTG	GATCTCCNC	TCCACCACTT	TCATGCGCTG	360

TTCNTACCCG	CGNATNTGTC	CCANCTGTTT	CNGTGCCNAC	TCCANCTTCT	NGGACGTGCG	420
CTACATACGC	CCGGANTCNC	NCTCCCGCTT	TGTCCCTATC	CACGTNCCAN	CAACAAATTT	480
CNCCNTANTG	CACCNATTCC	CACNTTTTNC	AGNTTTCCNC	NNCGNGCTTC	CTTNTAAAAG	540
GGTTGANCCC	CGGAAAATNC	CCCCAAGGGG	GGGGCCCNCC	TACCCAACTN	CCCCCTNATA	600
GCTGAANTCC	CCATNACCNN	GNCTCNAATG	ANCCNTCCNT	TTTAANNACN	TTCTNAACTT	660
GGGAANANCC	CTCGNCCNTN	CCCCCNTTAA	TCCCNCCCTG	CNANGNNCNT	CCCCCNNTCC	720
NCCCNNTNG	GCNTNTNANN	CNAAAAAGGC	CCNNNANCAA	TCTCCTNNCN	CCTCANTTCG	780
CCANCCCTCG	AAATCGGCCN	C				801

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 789 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CAGTCTATNT	GGCCAGTGTG	GCAGCTTTCC	CTGTGGCTGC	CGGTGCCACA	TGCCTGTCCC	60
ACAGTGTGGC	CGTGGTGACA	GCTTCAGCCG	CCCTCACCGG	GTTACCTTC	TCAGCCCTGC	120
AGATCCTGCC	CTACACACTG	GCCTCCCTCT	ACCACCGGGA	GAAGCAGGTG	TTCTGCCCCA	180
AATACCGAGG	GGACACTGGA	GGTGCTAGCA	GTGAGGACAG	CCTGATGACC	AGCTTCCTGC	240
CAGGCCCTAA	GCCTGGAGCT	CCCTTCCCTA	ATGGACACGT	GGGTGCTGGA	GGCAGTGGCC	300
TGCTCCACCC	TCCACCCGCG	CTCTGCGGGG	CCTCTGCCTG	TGATGTCTCC	GTACGTGTGG	360
TGGTGGGTGA	GCCCACCGAN	GCCAGGGTGG	TTCCGGGGCCG	GGGCATCTGC	CTGGACCTCG	420
CCATCCTGGA	TAGTGCTTCC	TGCTGTCCCA	NGTGGCCCCA	TCCCTGTTTA	TGGGCTCCAT	480
TGTCCAGCTC	AGCCAGTCTG	TCACTGCCTA	TATGGTGTCT	GCCGCAGGCC	TGGGTCTGGT	540
CCCATTTACT	TTGCTACACA	GGTANTATTT	GACAAGAACG	ANTTGGCCAA	ATACTCAGCG	600
TTAAAAAATT	CCAGCAACAT	TGGGGGTGGA	AGGCCTGCCT	CACTGGGTCC	AACTCCCCGC	660
TCCTGTTAAC	CCCATGGGGC	TGCCGGCTTG	GCCGCCAATT	TCTGTTGCTG	CCAAANTNAT	720
GTGGCTCTCT	GCTGCCACCT	GTTGCTGGCT	GAAGTGCNTA	CNGCNCANCT	NGGGGGGTNG	780
GGNGTTCCC						789

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 772 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CCCACCCTAC	CCAAATATTA	GACACCAACA	CAGAAAAGCT	AGCAATGGAT	TCCCTTCTAC	60
TTTGTAAAT	AAATAAGTTA	AATATTTAAA	TGCCTGTGTC	TCTGTGATGG	CAACAGAAGG	120
ACCAACAGGC	CACATCCTGA	TAAAAGGTAA	GAGGGGGGTG	GATCAGCAAA	AAGACAGTGC	180
TGTGGGCTGA	GGGGACCTGG	TTCTTGTGTG	TTGCCCTCA	CGACTCTTCC	CCTACAAATA	240
ACTTTCATAT	GTTCAAATCC	CATGGAGGAG	TGTTTCATCC	TAGAAACTCC	CATGCAAGAG	300
CTACATTAAA	CGAAGCTGCA	GGTTAAGGGG	CTTANAGATG	GGAAACCAGG	TGACTGAGTT	360
TATTCAGCTC	CCAAAAACCC	TTCTCTAGGT	GTGTCTCAAC	TAGGAGGCTA	GCTGTTAACC	420

40

```

CTGAGCCTGG GTAATCCACC TGCAGAGTCC CCGCATTCOA GTGCATGGAA CCCCTCTGGC 480
CTCCCTGTAT AAGTCCAGAC TGAACCCCCC TTGGAAGGNC TCCAGTCAGG CAGCCCTANA 540
AACTGGG3AA AAAAGAAAAG GACGCCCCAN CCCCAGCTG TGCANCTACG CACCTCAACA 600
GCACAGG3TG GCAGCAAAAA AACCACCTTA CTTTGGCACA AACAAAAACT NGGGGGGGCA 660
ACCCCGGCAC CCCNANGGGG GTTAACAGGA ANCNNGGNAA CNTGGAACCC AATTNAGGCA 720
GGCCCNCCAC CCCNAATNTT GCTGGGAAAT TTTTCTCTCC CTAAATTNTT TC 772

```

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 751 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```

GCCCCAATTC CAGCTGCCAC ACCACCCACG GTGACTGCAT TAGTTCGGAT GTCATACAAA 60
AGCTGATTGA AGCAACCCCTC TACTTTTTTG TCSTGAGCCT TTTGCTTGGT GCAGGTTTCA 120
TTGGCTGTGT TGGTGACGTT GTCATTGCAA CAGAATGGGG GAAAGGCACT GTTCTCTTTG 180
AAGTANGGTG AGTCCCTCAA ATCCGTATAG TTGCTGAAGC CACAGCACTT GAGCCCTTTC 240
ATGGTGGTGT TCCACACTTG AGTGAAGTCT TCCTGGGAAC CATAATCTTT CTTGATGGCA 300
GGCACTACCA GCAACGTCAG GGAAGTGCTC AGCCATTGTG GTGTACACCA AGGCGACCAC 360
AGCAGCTGCN ACCTCAGCAA TGAAGATGAN GAGGANGATG AAGAAGAACG TCNCGAGGGC 420
ACACTTGCTC TCAGTCTTAN CACCATANCA GCCCNTGAAA ACCAANANCA AAGACCACNA 480
CNCCGGCTGC GATGAAGAAA TNACCCCNCG TTGACAAACT TGCATGGCAC TGGGANCCAC 540
AGTGGCCNA AAAATCTTCA AAAAGGATGC CCCATCNATT GACCCCCCAA ATGCCCACTG 600
CCAACAGGGG CTGCCCCACN CNCNNAACGA TGANCCNATT GNACAAGATC TNCNTGGTCT 660
TNATNAACNT GAACCCTGCN TNGTGGCTCC TGTTCAGGNC CNNGGCCTGA CTTCTNAANN 720
AANGAACTCN GAAGNCCCCA CNGGANANNC G 751

```

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 729 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

```

GAGCCAGGGG TCCCTCTGCC TGCCCACTCA GTGGCAACAC CCGGGAGCTG TTTTGTCTTT 60
TGTGGANCCCT CAGCASTNCC CTCTTTCAGA ACTCANTGCC AAGANCCCTG AACAGGAGCC 120
ACCATGCAGT GCTTCAGCTT CATTAAGACC ATGATGATCC TCTTCAATTT GCTCATCTTT 180
CTGIGTGGTG CAGCCCTGTT GGCAGTGGGC ATCTGGGTGT CAATCGATGG GGCATCCTTT 240
CTGAAGATCT TCGGGCCACT GTGCTCCAGT GCCATGCAGT TTCTCAACGT GGGCTACTTC 300
CTCATCGCAG CCGGCGTTGT GGTCTTAGCT CTAGGTTTCC TGGGCTGCTA TGGTGCTAAG 360
ACTGAGAGCA ACTGTGCCCT CGTGACGTTT TTCTTCATCC TCCTCCTCAT CTTTATTGCT 420
GAGGTTGCAA TGCTGTGGTC GCCTTGGTGT ACACCACAAT GGCTGAGCA TTTCTGACGT 480
TGCTGTAAT GCTGTGCATC AANAAAAGAT TATGGGTTC CAGGAANACT TCACTCAAGT 540
GTTGGAACAC CACCATGAAA GGGCTCAAGT GCTGTGGCTT CNNGCAACTA TACGATTTT 600

```

GAAGANTCAC	CTACTTCAAA	GAAAAANAGTG	CCTTTCCCCC	ATTTCTGTTG	CAATTGACAA	660
ACGTCCCCAA	CACAGCCAAT	TGAAAACCTG	CACCCAACCC	AAANGGGTCC	CCAACCANAA	720
ATTNAAGGG						729

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 816 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TGCTCTTCCT	CAAAGTTGTT	CTTGTTGCCA	TAACAACCAC	CATAGGTAAA	GC3GGCCGAG	60
TGTTGCTGA	AGGGGTTGTA	GTACCAGCGC	GGGATGCTCT	CCTTGACAGG	TCCTGTGTCT	120
GGCAGGTCCA	CGCAGTGCCC	TTTGTCACTG	GGGAAATGGA	TGCGCTGGAG	CTCGTCAAAG	180
CCACTCGTGT	ATTTTTCACA	GGCAGCCTCG	TCCGACGCGT	CGGGJCAGTT	GGGGGTGTCT	240
TCACACTCCA	GGAAACTGTC	NATGCAGCAG	CCATTGCTGC	AGCGGAACTG	GGTGGGCTGA	300
CANGTGCCAG	AGCACACTGG	ATGGCGCCTT	TCCATGNNAN	GGGCCCTGNG	GGAAAGTCCC	360
TGANCCCCAN	ANCTGCCTCT	CAAANGCCCC	ACCTTGACAA	CCCCGACAGG	CTTAATGGA	420
ATCTTCTTCC	CGAAAGGTAG	TTNTTCTTGT	TGCCCAANCC	ANCCCCNTAA	ACAAACTCTT	480
GCANATCTGC	TCCGNGGGGG	TCNTANTACC	ANCGTGGGAA	AAGAACCCCA	GGGNGCGAAC	540
CAANCTTGTT	TGGATNCGAA	GCNATAATCT	NCNTTCTGTC	TTGGTGGACA	GCACCANTNA	600
CTGTNNANCT	TTAGNCCNTG	GTCCCTCNTGG	GTTGNNTTGT	AACCTAATCN	CCNNTCAACT	660
GGGACAAGGT	AANTNGCCNT	CCTTTNAATT	CCCNANCNTN	CCCCCTGGTT	TGGGGTTTTN	720
CNCNCTCCTA	CCCCAGAAAN	NCCGTGTTCC	CCCCCAACTA	GGGGCCNAAA	CCNNTTNTTC	780
CACAACCCTN	CCCCACCCAC	GGGTTGNGNT	GGTTNG			816

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 783 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CCAAGGCCCTG	GGCAGGCATA	NACTTGAAGG	TACAACCCCA	GGAACCCCTG	GT3CTGAAGC	60
ATGTGGAAAA	CACAGATTGG	CGCCTACTGC	GGGGTGACAC	GGATGTCAGG	GTAGAGAGGA	120
AAGACCCAAA	CCAGGTGGAA	CTGTGGGGAC	TCAAGGAANG	CACCTACCTG	TTCCA3CTGA	180
CAGTGACTAG	CTCA3ACCAC	CCAGAGGACA	CGGCCAACGT	CACAGTCACT	GT3CTGTCCA	240
CCAAGCAGAC	AGAAGACTAC	TGCCTCGCAT	CCAACAANGT	GGGTGCTGCG	CGGGGCTCTT	300
TCCCACGCTG	GTACTATGAC	CCCACGGAGC	AGATCTGCAA	GAGTTTCGTT	TATGGAG3CT	360
GCTTGGGCAA	CAAGAACAAC	TACCTTCGGG	AAGAAGAGTG	CATTCTANCC	TGTCNGGGTG	420
TGCAAGGTGG	GCCTTTGANA	NGCANCTCTG	GGGCTCANGC	GACTTTCCCC	CAGGGCCCCCT	480
CCATGGAAAG	GCGCCATCCA	NTGTTCTCTG	GCACCTGTCA	GCCCCCCCA3	TTCCGCTGCA	540
NCAATGGCTG	CTGCATCNAC	ANTTTCCTNG	AATTGTGACA	ACACCCCCCA	NTGCCCCCAA	600
CCCTCCCAAC	AAAGCTTCCC	TGTTNAAAAA	TACNCCANTT	GGCTTTTNAC	AAACNCCCGG	660
CNCCTCCNTT	TTCCCCNNTN	AACAAAGGGC	NCTNGCNTTT	GAACTGCCCN	AACCCNGGAA	720

TCTNCCNNGG AAAAAANTNCC CCCCCCTGGTT CCTNNAANCC CCTCCNCNAA ANCTNCCCCC 780
 CCC 783

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 801 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GCCCCAATTC	CAGCTGCCAC	ACCACCCACT	GTGACTGCAT	TAGTTGGGAT	ETCATACAAA	50
AGCTGATTGA	ASCAACCCCT	TACITTTTGG	TGCTGAGCC	TTTGCTTGGT	GCAGGTTTCA	120
TTGGCTGTGT	TGGTGAAGTT	GTCATTGCAA	CAGAATGGGG	GAAAGGCACT	TTTCTCTTTG	180
AAGTAGGGTG	AGTCTTCAAA	ATCCGTATAG	TTGGTGAAGC	CACAGCACTT	GAGCCCTTTC	240
ATGGTGGTGT	TCCACACTTG	AGTGAAGTCT	TCCTGGGAAC	CATAATCTTT	CTTGATGGCA	300
GGCACTACCA	GCAACGTCAG	GAAGTGCTCA	GCCATTGTGG	TGTACACCAA	GGCGACCACA	360
GCAGCTGCAA	CCTCAGCAAT	GAAGATGAGG	AGGAGGATGA	AGAAGAACGT	CNCGAGGGCA	420
CACTTGCTCT	CCGTCTTAGC	ACCATAGCAG	CCCANGAAAC	CAAGAGCAAA	GACCACAACG	480
CCNGCTGCCA	ATGAAAGAAA	NTACCCACGT	TGACAAACTG	CATGGCCACT	GGACGACAGT	540
TGGCCCGAAN	ATCTTCAGAA	AAGGGATGCC	CCATCGATTG	AACACCCANA	TGCCCCACTGC	600
CNACAGGGCT	GCNCCNCCN	GAAAGAATGA	GCCATTGAAG	AAGGATCNTC	NTGGTCTTAA	660
TGAACTGAAA	CCNTGCATGG	TGGCCCCCTG	TCAGGGCTCT	TGGCACTGAA	TTCTGANAAA	720
AAGGAACNGC	NTNAGCCCCC	CCAAANGANA	AAACACCCCC	GGGTGTTGCC	CTGAATTGGC	780
GGCCAAGGAN	CCCTGCCCCN	G				801

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 740 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GTGAGAGCCA	GGCGTCCCTC	TGCCTGCCCA	CTCACTGCCA	ACACCCGGGA	GCTGTTTTGT	50
ECTTTGTGGA	GCCTCAGCAG	TTCCCTCTTT	CAGAACTCAC	TGCCAAGAGC	CCTGAACAGG	120
AGCCACCATG	CAGTGCTTCA	GCTTCATTAA	GACCATGATG	ATCCTCTTCA	ATTTGCTCAT	180
CTTTCTGTGT	GGTGCAGCCC	TGTTGGCAGT	GGGCATCTGG	GTGTCAATCG	ATGGGGGCATC	240
CTTTCTGAAG	ATCTTCGGGC	CACTGTCGTC	CAGTGCCATG	CAGTTTGTCA	ACGTGGGGCTA	300
CTTCCTCATC	GCAGCCGGCG	TTGTGGTGCT	TGCTCTTGGT	TTCCTGGGGT	GCTATGGTGC	360
TAAGACGGAG	AGCAAGTGTG	CCCTCGTGAC	GTCTTCCTTC	ATCTCTCTCC	TCATCTTCAT	420
TGCTGAAGTT	GCAGCTGCTG	TGGTGGCCTT	GGTGTACACC	ACAATGGCTG	AACCATCTCT	480
GAGGTTGCTG	GTANTGCTTG	CCATCAANAA	AGATTATGGG	TTCCTCAGGAA	AAATTCACTC	540
AANTNTGGAA	CACCNCCATG	AAAAGGGCTC	CAATTTCTGN	TGGCTTCCCC	AACTATACCG	600
GAATTTTGAA	AGANTCNCCC	TACTTCCAAA	AAAAAANANT	TGCCTTTNCC	CCCTTCTGT	660
TGCAATGAAA	ACNTCCCAAN	ACNGCCAATN	AAAACCTGCC	CNNNCAAAAA	GGNTCNCAAA	720
CAAAAAAANT	NNAAGGTTN					740

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 802 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

```

CCGCTGGTTG CGCTGGTCCA GNGNAGCCAC GAAGCACGTC AGCATAACACA GCCTCAATCA      60
CAAGGTCTTC CAGCTGCCGC ACATTACGCA GGGCAAGAGC CTCCAGCAAC ACTGCATATG      120
GGATACACTT TACTTTAGCA GCCAGGGTGA CAACTGAGAG GTGTGGAAGC TTATTCTTCT      180
GAGCCTCTGT TAGTGGAGGA AGATTCCGGG CTTCAGCTAA GTAGTCAGCC TATGTCCCAT      240
AAGCAAACAC TGTGAGCAGC CGGAAGGTAG AGGCAAAGTC ACTCTCAGCC AGCTCTCTAA      300
CATTGGGCGT GTCCAGCAGT TCTCCAAACA CGTAGACACC AGNGGCCTCC AGCACCTGAT      360
GGATCACTCT GGCACCGCT GCGCCCTTGG CCGACTTGGC TAGGAGCAGA AATTGCTCCT      420
GGTTCTGCCC TGTCACCTTC ACTTCCGCAC TCATCACTGC ACTGAGTGTG GGGGACTTGG      480
GCTCAGGATG TCCAGAGACG TGGTTCGGCC CCGTCNCTTA ATGACACCGN CCANNCAACC      540
GTCGGCTCCC GCCGANTGNG TTCGTCGTNC CTGGGTCAGG GTCTGCTGGC CNCTACTTGC      600
AANCTTCGTC NGGCCCCATGG AATTCACCNC ACCGGAAC TN GTANGATCCA CTNNTTCTAT      660
AACCGGNEGC CACCGC NNT GGAACCTCAC TTTNTTNC TTTACTTGAG GGTAAAGGTC      720
ACCTTNNCG TTACCTTGGT CCAAACNTN CCGTGTGTG ANATNGTNAA TCNGGNCCNA      780
TNCCANCCNC ATANGAAGCC NG                                     802

```

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 731 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

```

CNAAGCTTCC AGGTNACGGG CCGCNAANCC TGACCCNAGG TANCANAANG CAGNCNGCGG      60
GAGCCCACCG TCACGNGGNG GNGTCTTTAT NGGAGGGGGC GGAGCCACAT CNCTGSACT      120
CNTGACCCCA ACTCCCNCT NCNCANTGUA GTGATGACTG CAGAACTGAA GGTNAOCTGG      180
CAGGAACCAA GANCAANNC TGCTCCNNTC CAAGTCGGCN NAGGGGGCGG GGCTGGGCAN      240
GCNCATCCNT CNAAGTGCTG AAAGCCCCNN CCGTGTCTACT TGTGTTGGAGA ACNGCNNGA      300
CATGCCCCAGN GTTANATAAC NGGCGAGAGG TNANTTTGCC TCTCCCTTCC GGCTGGGCAN      360
CGNGTNTGCT TAGNGGACAT AACCTGACTA CTTAACTGAA CCGNNGAATC TNCNCCCT      420
CCACTAAGCT CAGAACA AAACTTCGACA CCACTCANTT GTCACCTGNC TGCTCAAGTA      480
AAGTGTAACC CATNCCCAAT GTNTGCTNGA NGCTCTGNCC TGCNTTANGT TCGGTCTCTG      540
GAAGACCTAT CAATTNAAG TATGTTTCTG ACTGCCCTCT GCTCCCTGNA ACAANCNACC      600
CNNC NNTCA AGGGGGG GNC GGCCTCCAAT CCCCCCAACC NTNAATTNAN TTTANCCCN      660
CCCCNGGDC CGGCCTTTTA CNANCNTCN NNACNGGGNA AAACCNNGC TTTNCCCAAC      720
NNAATCCNCC T                                     731

```

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 754 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

```

TTTTTTTTTT TTTTTTTTTT TAAAAACCCC CTCATTNAA TGNAAACTTC CGAAATTGTC      50
CAACCCCCTC NTCCAAATNN CCNTTTCGGG GNGGGGGTTC CAAACCCAAN TTANNTTTGG      100
ANNTTAAATT AAATNTTNNNT TGGNGGNMNA ANCCNAATGT NANGAAAGTT NAACCCANTA      150
TNANCTTNAA TNCCTGGAAA CCGNGINGNTT CCAAAAATNT TTAACCCCTA ANTCCCTCCG      200
AAATN3TTNA NGGAAAACCC AANTTCTCNT AAGGTTGTTT GAAGGNTNAA TNAAAANCCC      250
NNCCAAATGT TTTTNGCCAC CCCIGAATTA ATTGGNTTCC GNTGTTTTCC NTTAAAANAA      300
GGNNANCCCC GGTANTNAA TCCCCCNCNN CCCAATTATA CCGANTTTTT TTNGAATTGG      350
GANCCNCGG GAATTAACGG GGNNNNTCC TTTGGGGGG CNGGNNCCCC CCCCNTCCGG      400
CGTTN3GGNC AGGNONNAAT TGTTTAAGGG TCCGAAAAAT CCTCCNAGA AAAAAANCTC      450
CCAGGNTGAG NNTNGGGTTT NCCCCCCCCC CANGGCCCT CTGNNANAGT TGGGGTTTGG      500
GGGGCTGGG ATTTTNTTTT CCTNTTNC TCCCCCCCCC CNGGGGANAG AGGTTNGNGT      550
TTTGNTCNCN GGGCCNCNCN AAGANCTTTN CCGANTTNAN TTAAATCCNT GCCTNGGCGA      600
AGTCCNTTGN AGGNTAAAN GGGCCCTNN CGGG                                     754

```

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 755 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

```

ATCANCOCAT GACCCCNAA C NNGGGACCNC TCANCCGGNC NNWNCACCNC CGGCCNATCA      50
NNGTNAGNNC ACTNONTTN NATCACNCCC CNCCNACTAC GCCCNCMANC CNACGCNCTA      100
NNCANATNCC ACTGANNGCG CGANGTNGAN NGAGAAANCT NATACCANAG NCACCANACN      150
CCAGCTGTCC NANAANGCCT NNNATACNGG NNNATCCAAT NTGNANCCTC CNAAGTATTN      200
NNCNCANAT GATTTTCCTN ANCCGATTAC CCNTNCCCCC TANCCCCCTC CCCCCAACNA      250
CGAAGGCNCT CGNCCNAAGG NNGCGNCNCC CCGCTAGNTC CCCNNCAAGT CNCNCNCCTA      300
AACTCANCCN NATACNCGC TTCNTGAGTA TCACTCCCCG AATCTCACCC TACTCAACTC      350
AAAAANATCN GATACAAAAT AATNCAAGCC TGNTTATNAC ACTNTGACTG GGTCTCTATT      400
TTAGNGGTCC NTNAANCNTC CTAATACTTC CAGTCTNCCT TCNCCAATTT CCNAANGGCT      450
CTTTGNGACA GCATNTTTTG GTTCCCNNTT GGGTCTTAN NGAAATGCCC TTCNTNGAAC      500
GGGCTCNTCT TTTCTTCGG TTANCCTGGN TTCNNCCGGC CAGTTATTAT TTCCNTTTT      550
AAATTCTNCC NTNTTNTT TGGCNTTCNA AACCCCGGC CTTGAAAACG GCCCCCTGGT      600
AAAAGGTTGT TTTGANAATA TTTTGTGTTT GTTCC                                     755

```

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 849 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

TTTTTTTTTT	TTTTTANGTG	TNGTCGTGCA	GGTAGAGGCT	TACTACAANT	GTGAANACGT	60
ACGCTNGGAN	TAANGCGACC	CGANTTCTAG	GANNNCNCCCT	AAAATCANAC	TGTGAAGATN	120
ATCCTGNMNA	CGGAANGGTC	ACCGGNNGAT	NNTGCTAGGG	TGNCCNCTCC	CANNNCNTTN	180
CATAACTCNG	NGGCCCTGCC	CACCACCTTC	GGCGGCCCCNG	NGNCCGGGGCC	CGGGTCATTN	240
GNMTTAACCN	CACTNNGCNA	NCGGTTTCCN	NCCCCNNCNG	ACCCNNGGCGA	TCCGGGGTNC	300
TCTGTCTTCC	CCTGNAGNCN	ANAAANTGGG	CCNCGGNCCC	CTTTACCCCT	NNACAAGCCA	360
CNGCCNTCTA	NCCNCNGCCC	CCCCCTCCANT	NNGGGGGACT	GCCNANNGCT	CCGTTNCTNG	420
NNACCCCNNN	GGGTNCCTCG	GTTGTGANT	CNACCGNANG	CCANGGATTC	CNAAGGAAGG	480
TGCGTTNTTG	GGCCCTACCC	TTCGCTNCGG	NNCACCTTC	CCGACNANGA	NCCGCTCCCG	540
CNCNNCGMNG	CCTCNCCTCG	CAACACCCGC	NCTCNCNGT	NCGGNNNCCC	CCCCACCCGC	600
NCCCTCNCNC	NGNCGNANCN	CTCCNCCNCC	GTCTCANNCA	CCACCCCGCC	CCGCCAGGCC	660
NTCANCCACN	GGMNGACNNG	NAGCNCNNTC	GCNCCGCGCN	GCGNCNCCCT	CGCCNCNGAA	720
CTNCNTCNGG	CCANTMNCGC	TCAANCCNNA	CNAAACGCCG	CTGCGCGGCC	CGNAGCGNCC	780
NCCTCCNCGA	GTCCTCCCGN	CTTCCNACCC	ANGNNTTCCN	CGAGGACACN	NNACCCCGCC	840
NNCANGCGG						849

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 872 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GCGCAAACCTA	TACTTCGCTC	GNACTCGTGC	GCCTCGCTNC	TCTTTTCCTC	CGCAACCATG	60
TCTGACNANC	CCGATTNGGC	NGATATCNAN	AAGNTCGANC	AGTCCAAACT	GANTAACACA	120
CACACNCNAN	AGANAAATCC	NCTGCCTTCC	ANAGTANACN	ATTGAACNNG	AGAACCANGC	180
NGGCGAATCG	TAATNAGGCG	TGCGCCGCCA	ATNTGTCNCC	GTTTATTNTN	CCAGCNTCNC	240
CTNCCNACCC	TACNTCTTCN	NAGCTGTENN	ACCCCTNGTN	CGNACCCCCC	NAGGTCGGGA	300
TGGGTTTNN	NNTGACCGNG	CNNCCCCCTC	CCCCNTCCAT	NACGANCCNC	CCGCACCACC	360
NANNGCNCSC	NCCCCGNCT	CTTCGCCNCC	CTGTCTNTN	CCCCGTNGC	CTGGCNCNGN	420
ACCGCATGGA	CCCTCGCCNN	CTNCNNGAAA	NCGNANACGT	CCGGGTGNN	ANNANCGCTG	480
TGGGNNGCG	TCTGCNCCGC	GTTCTTCCN	NCNNCTTCCA	CCATCTTCNT	TACNGGGTCT	540
CCNCGCCNTC	TCNNNCACNC	CCTGGGACGC	TNTCCTNTGC	CCCCCTTNAC	TCCCCCCCCTT	600
CGNCGTGNCC	CGNCCCCACC	NTCATTNCA	NACGNTCTTC	ACAANNNCCT	GGNTNNTCC	660
CNANCNGNCN	GTCANCCNAG	GGAAGGGNGG	GGNNCCNNTG	NTTGACGTTG	NGGNGANGTC	720
CGAANANTCC	TCNCCNTCAN	CNCTACCCCT	CGGGCGNNCT	CTCNGTTNCC	AACTTANCAA	780
NTCTCCCCCG	NGNGCNCNTC	TCAGCCTCNC	CCNCCCCNCT	CTCTGCANTG	TNCTCTGCTC	840
TNACCNTTAC	GANTNTTCGN	CNCCCTCTTT	CC			872

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

46

- (A) LENGTH: 815 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

```

GCATGCAAGC TTGAGIATTC TATAGNGTCA CCTAAATANC TTGGCNTAAT CATGGTCNTA      50
NCTGNCTTCC TGTGTCAAAAT GTATACNAAN TANATATGAA TCTNATNTGA CAAGANNGTA      100
TCNTNCATTA GTAACAANTG TNNTGTCCAT CCTGTCNGAN CANATTCCCA TNNATTNCGN      150
CGCATTNCN CNANCANTATN TAATNGGGAA NTCNNNTNNN NCACCNNCAT CTATCNTNCC      200
GCNCCCTGAC TGGNAGAGAT GGATNANTTC TNNTNTGACC NACATGTTCA TCTTGGATTN      250
AANANCCCCC CGCNGNCCAC CGGTTNGNNG CNAGCCNNTC CCAAGACCTC CTGTGGAGGT      300
AACCTGGGTC AGANNCATCA AACNTGGGAA ACCCGCANNCC ANGTNNAAGT NGNNNCANAN      350
GATCCCGTCC AGNNTTNAAC ATCCCTTCNC AGCGCCCCCT TTNGTGCCTT ANAGNGNAGC      400
GTGTCCNANC CNCTCAACAT GANACGCGCC AGNCCANCCG CAATTNGGCA CAATGTCGNC      450
GAACCCCTTA GGGGGANTNA TNCAAANCCC CAGGATTGTC CNCNCANGAA ATCCNCANC      500
CCNCCCTAC CCNCTTTTGG GACNGTGACC AANTCCCGGA GTNCCAGTCC GGCCNGNCTC      550
CCCCACCGGT NNCCNTGGGG GGGTGAANCT CNGNNTCANC CNGNCGAGGN NTCGNAAGGA      600
ACCGGNTCTN GGNCBAANNG ANCNNTCNGA AGNGCCNCNT CGTATAACCC CCCCNCNCCA      650
NCCNACNGNT AGNTCCCCC CNGGGTNCGG AANGG      700
      750
      800
      815
  
```

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 775 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

```

CCGAGATGTC TCGCTCCGTG GCCTTAGCTG TGCTCGCGCT ACTCTCTCTT TCTGGCCTGG      60
AGGCTATCCA GCGTACTCCA AAGATTCAGG TTTACTCAGC TCATCCAGCA GAGAATGGAA      120
AGTCAAATTT CCTGAATTGC TATGTGTCTG GGTTCATCC ATCCGACATT GAANTTGACT      180
TACTGAAGAA TGGANAGAGA ATTGAAAAAG TGGAGCATTC AGACTTGTCT TTCAGCAAGG      240
ACTGGTCTTT CTATCTCNTG TACTACACTG AATTCACCCC CACTGAAAAA GATGACTATC      300
CCTGCCGTGT GAACCATGTG ACTTTGTGAC AGCCCAAGAT AGTTAAGTGG GATCGAGACA      360
TGTAAGCAGN CNNCATGGAA GTTTGAAGAT GCCGCATTTG GATTGGATGA ATTCCAAATT      420
CTGCTTGCTT GNTTTTAAAT ANTGATATGC NTATACACCC TACCCTTTAT GNCCCCAAAT      480
TGTAGGGGTT ACATNANTGT TCNCNTNGGA CATGATCTTC CTTTATAANT CCNCCNTTCG      540
AATTGCCCCG CCCCCNGTTN NGAATGTTTC CNNAACCACG GTTGGCTCCC CCAGGTCTCC      600
TCTTACGGAA GGGCCTGGGC CNCTTTNCAA GGTGGGGGGA ACCNAAAATT TCNCTTNTGC      660
CCNCCNCCA CNNTCTTGNG NNCNCANTTT GGAACCCCTC CNATTCCCCC TGGCCTCNNA      720
NCCTTNCTA ANAAAACTT AAANCGTNGC NAAANNTTTN ACTTCCCCC TTACC      775
  
```

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 820 base pairs
 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

ANATTANTAC	AGTGTAATCT	TTCCCCAGAG	GTGTGTANAG	GGAACGGGGC	CTAGAGGCAT	60
CCCANAGATA	NCTTATANCA	ACAGTGCTTT	GACCAAGAGC	TGCTGGGCAC	ATTTCCCTGCA	120
GAAAAGGTGG	CGGTCCCCAT	CACTCCTCCT	CTCCCATAGC	CATCCCAGAG	GGGTGAGTAG	180
CCATCANGCC	TTCGGTGGGA	GGGAGTCANG	GAAACAACAN	ACCACAGAGC	ANACAGACCA	240
NTGATGACCA	TGGGCGGGAG	CGAGCCTCTT	CCCTGNACCG	GGGTGGCANA	NGANAGCCTA	300
NCTGAGGGGT	CACACTATAA	ACGTTAACGA	CCNAGATNAN	CACCTGCTTC	AAGTGCACCC	360
TTCTACCTG	ACNACCAGNG	ACCNNAACT	GCNGCCTGGG	GACAGCNCTG	GGANCAGCTA	420
ACNNAGCACT	CACCTGCCCC	CCCATGGCCG	TNCGCNTCCC	TGGTCCTGNC	AAGGGAAGCT	480
CCCTGTTGGA	ATTNCGGGGA	NACCAAGGGA	NCCCCCTCCT	CCANCTGTGA	AGGAAAAANN	540
GATGGAATTT	TNCCCTTCCG	GCCNNTCCCC	TCTTCCTTTA	CACGCCCCCT	NNTACTCNTC	600
TCCCTCTNTT	NTCCTGNCNC	ACTTTTNACC	CCNNNATTTT	CCTTNATTGA	TCGGANNCTN	660
GANATTCCAC	TNNCGCCTNC	CNTCNATCNG	NAANACNAAA	NACTNTCTNA	CCCGGGGGAT	720
GGGNCCCTCG	NTCATCCTCT	CTTTTTCNCT	ACCNCCNNTT	CTTTGCCTCT	CCTTINGATCA	780
TCCAACCNTC	GNTGGCCNTN	CCCCCCCNNN	TCCTTTNCCC			820

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 818 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

TCTGGGTGAT	GGCCTCTTCC	TCCTCAGGGA	CCTCTGACTG	CTCTGGGCCA	AAGAATCTCT	60
TGTTTCTTCT	CCGAGCCCCA	GGCAGCGGTG	ATTCAGCCCT	GCCCCAACCTG	ATTCTGATGA	120
CTGCGGATGC	TGTGACGGAC	CCAAGGGGCA	AATAGGGTCC	CAGGGTCCAG	GGAGGGGGCGC	180
CTGCTGAGCA	CTTCCGCCCC	TCACCCTGCC	CAGCCCCCTGC	CATGAGCTCT	GGGCTGGGTC	240
TCCGCCTCCA	GGGTTCTGCT	CTTCCANGCA	NGCCANCAAG	TGGCGCTGGG	CCACACTGGC	300
TTCTTCCTGC	CCCNTCCCTG	GCTCTGANTC	TCTGTCTTCC	TGTCCTGTGC	ANGCNCCTTG	360
GATCTCAGTT	TCCCTCNCCT	ANNGAACTCT	GTTTCTGANN	TCTTCANTTA	ACTNTGANTT	420
TATNACCNAN	TGGNCTGTNC	TGTCNNACTT	TAATGGGGCCN	GACCGGCTAA	TCCCTCCCTC	480
NCTCCCTTCC	ANTTCNRRNA	ACCNGCCTTNC	CNTCNTCTCC	CCNTANCCCG	CCNGGGGAANC	540
CTCCTTTGCC	CTNACCANGG	GCCNNNACCG	CCCNTNNCTN	GGGGGGCCNNG	GTNNCTNCNC	600
CTCNTNNCCC	CNCTCNCNNT	TNCCTCGTCC	CNNCNCNCGCN	NNGCANNTTC	NCGTCCCNNT	660
TNNCTCTTCN	NGTNTCGNAA	NGNTCNCNTN	TNNNNNGNCH	NGNTNNTNCH	TCCCTCTCNC	720
CNNNTGNANG	TNNTTNWNNC	NCGNNCCCCC	NNNNCNCNNN	NGGNNNTNNT	TCTNCCNCGC	780
CCCNCCCCC	NGNATTAAGG	CCTCCNNTCT	CCGGCCNC			818

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 731 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

AGGAAGGGCG	GAGGGATATT	GTANGGGATT	GAGGGATAGG	AGNATAANGG	GGGAGGTGTG	60
TCCCAACATG	ANGGTGNGGT	TCTCTTTTGA	ANGAGGGTTG	NGTTTTTAN	CCNGGTGGGT	120
GATTNAACCC	CATTGTATGG	AGNNAAGGN	TTTNAGGGAT	TTTTCGGCTC	TTATCAGTAT	180
NTANATTCCCT	G'TNAATCGGA	AAATNATNTT	TCNNCNGGAA	AATNTTGCTC	CCATCCGNAA	240
ATTNCTCCCC	GGTAGTGCAT	NTTNGGGGGN	CNGCCANGTT	TCCCAGGCTC	CTANAATCGT	300
ACTAAAGNTT	NAAGTGGGAN	TNCAAATGAA	AACCTNNCAC	AGAGNATCCN	TACCCGACTG	360
TNNNTTNCCT	TCGCCCNTNG	ACTCTGCNNG	AGCCCAATAC	CCNNGNGNAT	GTNCCCNGN	420
NNNGCGNCNC	TGAAANNNNC	TCGNGGCTNN	GANCATCANG	GGGTTTCGCA	TCAAAAAGCNV	480
CGTTTTNCAT	NAAGGCACTT	TNGCCTCATC	CAACCNCTNG	CCCTCNCCA	TTTNGCGCTC	540
NGGTTNCCT	ACGCTNNTNG	CNCCTNNNTN	GANATTTTNC	CCGCTNGGG	NAANCCTCCT	600
GNAATGGTGA	GGGNCCTTNC	TTTNACCNV	GNGGTNTACT	AATCNNTTNC	ACGCTNCTT	660
TCTCNACCCC	CCCCCTTTTT	CAATCCCAN	GGCNAATGGG	GTCTCCCTCN	CGANGGGGGG	720
NNNCCCANNC	C					731

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 822 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

ACTAGTCCAG	TGTGGTGGAA	TTCCATTGTG	TTGGGGNCNC	TTCTATGANT	ANNTNTTAGAT	60
CGCTCANACC	TCACANCCTC	CCNACNANGC	CTATAANGAA	MANNAATAGA	NCTGTNCNNT	120
ATNTNTACNC	TCATANNCCT	CNNNACCCAC	TCCCTCTTAA	CCCNCTACTGT	GCCTATNGCN	180
TNNCTANTCT	NTGCCGCCCTN	CNANCCACCN	GTGGGGCCNAC	CNCNNGNATT	CTCNATCTCC	240
TCNCCATNTN	GCCTANANTA	NGTNCATACC	CTATACCTAC	NCCAATGCTA	NNNCTAANCN	300
TCCATNANTT	ANNNTAACTA	CCACTGACNT	NGACTTTCNC	ATNANCTCCT	AATTTGAATC	360
TACTCTGACT	CCCACNGCCT	ANNNATTAGC	ANCNTCCCCC	NACNATNTCT	CAACCAAATC	420
NTCAACAACC	TATCTANCTG	TTCNCCAACC	NTTNCTCCG	ATCCCCNNAC	AACCCCCCTC	480
CCAAATACCC	NCCACCTGAC	NCCTAACCCN	CACCATCCCC	GCAAGCCNAN	GGNCATTTAN	540
CCACTGGAAT	CACNATNGGA	NAAAAAAAC	CCNAACTCTC	TANCNCNNAT	CTCCCTAANA	600
AATNCTCCTN	NAATTTACTN	NCANTNCCAT	CAANCCACN	TGAAACNNAA	CCCCGTTTTT	660
TANATCCCTT	CTTTCGAAAA	CCNACCCTTT	ANNNCCCAAC	CTTTNGGGCC	CCCCCNCTNC	720
CCNAATGAAG	GNCNCCCAAT	CNANGAAACG	NCCNTGAAAA	ANCNAGGCNA	ANANNNTCCG	780
CANATCCTAT	CCCTTANTTN	GGGGNCCCTT	NCCCNGGGCC	CC		822

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 787 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

```

CGGCCCGCCTG CTCTGGCACA TGCCTCCTGA ATGGCATCAA AAGTGATGGA CTGCCCATTG      60
CTAGAGAAGA CCTTCTCTCC TACTGTCAAT ATGGAGCCCT GCAGACTGAG GGCTCCCCTT      120
GTCTGCAGGA TTGATGTCT GAAGTCGTGG AGTGTGGCTT GGAGCTCCTC ATCTACATNA      180
GCTGGAAGCC CTGGAGGGCC TCTCTGCCA GCCTCCCCCT TCTCTCCACG CTCTCCANGG      240
ACACCAGGGG CTCCAGGCAG CCCATTATTC CCAGNANGAC ATGGTGTTC TCCACGCGGA      300
CCCATGGGGC CTGNAAGGCC AGGGTCTCCT TTGACACCAT CTCTCCCGTC CTGCCTGGCA      360
GGCCGTGGGA TCCACTANTT CTANAACGGN CGCCACCNCG GTGGGAGCTC CAGCTTTTGT      420
TCCCNNTAAT GAAGGTAAAT TGCNCGCTTG GCGTAATCAT NGGTCANAAC TTTTCCTGT      480
GTGAAATTGT TTNTCCCCTC NCNATTCCNC NCNACATACN AACCCGGAAN CATAAAGTGT      540
TAAAGCCTGG GGTNGCCTN NNGAATNAAC TNAACTCAAT TAATTGCGTT GGCTCATGGC      600
CCGCTTTCCN TTCNGGAAA CTGTCTCCC CTGCNTTNNNT GAATCGGCCA CCCCCNGGG      660
AAAAGCGGTT TGCNTTTTNG GGGGNTCCTT CCNCTTCCCC CCTCNCTAAN CCTNCGCCT      720
CGGTCGTTNC NGGTNGCGGG GAANGGGNAT NNNCTCCNC NAAGGGGGNG AGNNNGNTAT      780
CCCCAA

```

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 799 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

```

TTTTTTTTTT TTTTTTTGGC GATGCTACTG TTTAATTGCA GGAGGTGGGG GTGTGTGTAC      60
CATGTACCAG GGCTATTAGA AGCAAGAAGG AAGGAGGGAG GGCAGAGCGC CCTGCTGAGC      120
AACAAAGGAC TCCTGCAGCC TTCTCTGTCT GTCTCTTGGC GCAGGCACAT GGGGAGGCCT      180
CCCGCAGGGT GGGGGCCACC AGTCCAGGGG TGGGAGCACT ACANGGGGTG GGAGTGGGTG      240
GTGGCTGGTN CNAATGGCCT GNCACANATC CCTACGATTC TTGACACCTG GATTTACCA      300
GGGGACCTTC TGTTCTCCCA NGGNAACTTC NTNNATCTCN AAAGAACACA ACTGTTTCTT      360
CNGCANTTCT GGCTGTTTAT GGAAAGCACA GGTGTCCNAT TTNGGCTGGG ACTTGGTACA      420
TATGGTTCCG GCCCACCTCT CCNCTCNAAN AAGTAATTCA CCCCCCCCN CCNCTNTTGT      480
CCTGGGCCCT TAANTACCCA CACCGGAAC CANTTANTTA TTCATCTTNG GNTGGGCTTG      540
NTNATCNCCN CCTGAANGCG CCAAGTTGAA AGGCCACGCC GTNCCCNCTC CCCATAGNAN      600
NTTTTNNCNT CANCTAATGC CCCCCCNGGC AACNATCCAA TCCCCCCCCN TGGGGGCCCC      660
AGCCCANGGC CCCCNGCTCG GGNNCCNGN CNCGNANTCC CCAGGNTCTC CCANTCNGNC      720
CCNNNGCNCC CCCGCACGCA GAACANAAGG NTNGAGCCNC CGCANNNNNN NGGTNNCNA      780
CTCGCCCCCC CCNCCGNG

```

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 789 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	60
TTTTNNCNA3	GGCAGGTTTA	TTGACAACCT	CNCGGGACAC	AANCAGGCTG	GGGACAGGAC	120
GGCAATAGGC	TCCGGCG3CG	GCGGCGGCGG	CCCTACCTGC	GSTACCAAAT	NTGCAG3CTC	180
CGCTCCGCT	TGATNTTCCT	CTGCAGCTGC	AGGATGCCNT	AAAACAGGGC	CTCGGC3NTN	240
GGTGG3CACG	CTGGGATTTN	AATTTCCACG	GGCACAATGC	GGTCGCANCC	GGTCAC3ACC	300
NATTA3GAAT	AGTGGTNTTA	CCCNCCNCCG	TTGGCNCAC	GGCCNTGGAA	ACCACTTNTC	360
GCGGCTCCGG	CATCTGGTCT	TAAACCTTGC	AAACNCTGGG	GGCCTCTTTT	TGGTTANTNT	420
NCCNG3CACA	ATCATNACTC	AGACTGGCNC	GGGCTGGCCC	CAAAAAANCN	CCCCAAAACC	480
GGNCCATGTC	TTNNCGGGGT	TGCTGCNATN	TNCATCACCT	GGCGGGCNCA	NCAGGNCAAC	540
CCAAAACCTC	TTGNG3CCCN	CAAAAAANCT	CCGGGGGGNC	CCAGTTTCAA	CAAAGTCATC	600
GGCCTTGGCC	CCCAAATCCT	CCCCCGGNTT	NCTGGGTTTG	GGAACCCACG	CCTCTNNCTT	660
TGGNN3GCAA	GNTGGNTCCC	CCTTCGGGCG	CCCGGTGGGC	GGNNCTCTAA	NGAAAAACNC	720
NTCCTNNNCA	GCATCC3CCC	NNGNNACGNT	TANCAANGNA	TCCCTTTTTT	TANAAAAGGG	780
CCCCC3CNC						793

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 793 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

GACAGAACAT	GTTGGATGGT	GGAGCACCTT	TCTATACGAC	TTACAGGACA	GCAGATGGGG	60
AATTCATGGC	TGTTGGAGCA	ATANAACCCC	AGTTCTACGA	GCTGCTGATC	AAAGGACTTG	120
GAATAAAGTC	TGATGAACTT	CCCAATCAGA	TGAGCATGGA	TGATTGGCCA	GAAATGAANA	180
AGAAGTTTGC	AGATGTATTT	GCAAAGAAGA	CGAAGGCAGA	GTGGTGTCAA	ATCTTTGACG	240
GCACAGATGC	GTGTGTGACT	CCGGTTCCTG	CTTTTGAGGA	GGTTGTTTAT	CATGATCACA	300
ACAANGAACG	GGGCTCGTTT	ATCACCANTG	AGGAGCAGGA	CGTGAGCCCC	CGCCCTGCAC	360
CTCTGCTGTT	AAACACCCCA	GCCATCCCTT	CTTTCAAAAG	GGATCCACTA	CTTCTAGAGC	420
GGNCGCCACC	GCGGTGGAGC	TCCAGCTTTT	GTTCCCTTTA	GTGAGGGTTA	ATTGCGCGCT	480
TGGCGTAATC	ATGGTCATAN	CTGTTTCCTG	TGTGAAATTG	TTATCCGCTC	ACAAATCCAC	540
ACAACATACG	ANCCGGAAGC	ATNAAATTTT	AAAGCCTGGN	GGTNGCCTAA	TGANTGAACT	600
NACTCACATT	AATTGGCTTT	GCGCTCACTG	CCCGCTTTCC	AGTCCCGAAA	ACCTGTCCTT	660
GCCAGCTGCC	NTTAATGAAT	CNGGCCACCC	CCCGGGGAAA	AGGCNGTTTG	CTTNTTGGGG	720
CGCNCTTCCC	GCTTTCTCGC	TTCCTGAANT	CCTTCCCCCC	GCTCTTTCGG	CTTGCGGCNA	780
ACGGTATCNA	CCT					793

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 756 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GGCGCGACCG	GCATGTACGA	GCAACTCAAG	GGCGAGTGGA	ACCGTAAAAG	CCCCAATCTT	60
ANCAAGTGCG	GGGAANAGCT	GGGTGACTC	AAGCTAGTTC	TTCTGGAGCT	CAACTTCTTG	120
CCAACCACAG	GGACCAAGCT	GACCAAACAG	CAGCTAATTC	TGGCCCGTGA	CATACTGGAG	180
ATCGGGGGCC	AATGGAGCAT	CCTACGCAAN	GACATCCCCCT	CCTTCGAGCG	CTACATGGCC	240
CAGCTCAAAT	GCTACTACTT	TGATTACAAN	GAGCAGCTCC	CCGAGTCAGC	CTATATGCAC	300
CAGCTCTTGG	GCCTCAACCT	CCTCTTCCTG	CTGTCCCAGA	ACCGGGTGGC	TGANTNCCAC	360
ACGGANTTGG	ANCGGCTGCC	TGCCCAANGA	CATACANACC	AATGTCTACA	TCNACCACCA	420
GTGTCCTGGA	GCAATACTGA	TGGANGGCAG	CTACCNCAA	GTNTTCCTGG	CCNAGGGTAA	480
CATCCCCCGC	CGAGAGCTAC	ACCTTCTTCA	TTGACATCCT	GCTCGACACT	ATCAGGGATG	540
AAAATCGCNG	GGTTGCTCCA	GAAAGGCTNC	AANAANATCC	TTTTCNCTGA	AGGCCCCCGG	600
ATNCTAGT	NCTAGAATCG	GCCCCCATC	GCGGTGGANC	CTCCAACCTT	TCGTTNCCCT	660
TTACTGAGGG	TTNATTGCCG	CCCTTGGCGT	TATCATGGTC	ACNCCNGTTN	CCTGTGTTGA	720
AATTNTTAAC	CCCCACAAT	TCCACGCCNA	CATTNG			756

(2) INFORMATION FOR SEQ ID NO:35:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 834 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(11) MOLECULE TYPE: cDNA

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:35:

GGGGATCTCT	ANATCNACCT	GNATGCATGG	TTGTGGGTGT	GGTCGCTGTC	GATGAANATG	60
AACAGGATCT	TGCCCTTGAA	GCTCTCGGCT	GCTGTNTTTA	AGTTGCTCAG	TCTGCCGTCA	120
TAGTCAGACA	CNCTCTTGGG	CAAAAAACAN	CAGGATNTGA	GTCTTGATTT	CACCTCCAAT	180
AATCTTCNGG	GCTGTCTGCT	CGGTGAACTC	GATGACNANG	GGCAGCTGGT	TGTGTNTGAT	240
AAANTCCANC	ANGTTCTCCT	TGGTGACCTC	CCCTTCAAAG	TTGTTCCGGC	CTTCATCAAA	300
CTTCTNNAAN	ANGANNANCC	CANCTTTGTC	GAGCTGGNAT	TTGGANAACA	CGTCACTGTT	360
GGAAACTGAT	CCCAAATGGT	ATGTCAATCCA	TCGCCTCTGC	TGCCTGCAAA	AAACTTGCTT	420
GGCNCAAATC	CGACTCCCCN	TCCTTGAAAG	AAGCCNATCA	CACCCCCCTC	CCTGGACTCC	480
NNCAANGACT	CTNCCGCTNC	CCCNTCCNNG	CAGGGTTGGT	GGCANNCCGG	GCCCNTGCGC	540
TTCTTCAGCC	AGTTCACNAT	NTTCATCAGC	CCCTCTGCCA	GCTGTTNTAT	TCCTTGGGGG	600
GGAANCCGTC	TCTCCCTTCC	TGAANNAACT	TTGACCGTNG	GAATAGCCGC	GCNTCNCCNT	660
ACNTNCTGGG	CCGGGTTCAA	ANTCCCTCCN	TTGNCNNTCN	CCTCGGGCCA	TTCTGGATTT	720
NCCNAACTTT	TTCTTCCCC	CNCCCCNCGG	NGTTTGGNTT	TTTCATNGGG	CCCCAACTCT	780
GCTNTTGGCC	ANTCCCCCTGG	GGGCTNTAN	CNCCCCCTNT	GGTCCCNNG	GGCC	834

(2) INFORMATION FOR SEQ ID NO:36:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 814 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(11) MOLECULE TYPE: cDNA

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:36:

CGGNGCGTTT	CGNGCCGCGG	CGCGTTTCCA	TGACNAAGGC	TCCCTTCANG	TTAAATACNN	60
CCTAGNAAAC	ATTAATGGGT	TGCTCTACTA	ATACATCATA	CNAACCAGTA	AGCCTGCCCC	120
NAACGCCAAC	TCAGGCCATT	CCTACGAAAAG	GAAGAAAAGGC	TGGTCTCTCC	ACCCCTGTGA	180
GGAAAGGCCT	GCCTTGTAAG	ACACCACAAT	NCGGCTGAAT	CTNAAGTCTT	GTCTTTTACT	240
AATGGAAAAA	AAAAATAAAC	AANAGGTTTT	GTCTCATGG	CTGCCCACCG	CAGCCTGGCA	300
CTAAAAACAC	CCAGCGCTCA	CTTCTGCTTG	GANAAATATT	CTTGCTCTT	TTGGACATCA	360
GGCTTGATGG	TATCACTGCC	ACNTTTCAC	CCAGCTGGGC	NCCCTTCCCC	CATNTTTGTC	420
ANTGANCTGG	AAGSCCTGAA	NCTTAGTCTC	CAAAAGTCTC	NGCCACAAG	ACCGGCCACC	480
AGGGGANGTC	NTTNCAGTG	GATCTGCCAA	ANANTACCCN	TATCATCNNT	GAATAAAAAAG	540
GCCCTGAAC	GANATGCTTC	CANCANCTT	TAAGACCCAT	AATCCTNGAA	CCATGGTGCC	600
CTTCCGGTCT	GATCCNAAAG	GAATGTTCCCT	GGGTCCGANT	CCCTCCTTTG	TTNCTTACGT	660
TGTNTTGGAC	CCNTGCTNGN	ATNACCCAAAN	TGANATCCCC	NGAAGCACCC	TNCCCTGGC	720
ATTTGANTTT	CNTAAATTCT	CTGCCCTACN	NCTGAAAGCA	CNATTCCTN	GGCNCNNAAN	780
GGNGAACTCA	AGAAGGTCTN	NGAAAAACCA	CNCN			840

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 750 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

GCATGCTGCT	CTTCCTCAAA	GTTGTTCTTG	TTGCCATAAC	AACCACCATA	GGTAAAGCGG	60
GCGCAGTGTT	CGCTGAAGGG	GTTGTAGTAC	CAGCGCGGGA	TGCTCTCCTT	GCAGAGTCCT	120
GTGTCTGGCA	GGTCCACGCA	ATGCCCTTTG	TCACTGGGGA	AATGGATGCG	CTGGAGCTCG	180
TCNAANCCAC	TCGTGTATTT	TTACANGCA	GCCTCCTCCG	AAGCNTCCGG	GCAGTTGGGG	240
GTGTCTGTCAC	ACTCCACTAA	ACTGTGATN	CANCAGCCCA	TTGCTGCAGC	GGAACTGGGT	300
GGGCTGACAG	GTGCCAGAAC	ACACTGGATN	GGCCTTTCCA	TGGAAGGGCC	TGGGGGAAAT	360
CNCCTNANCC	CAAACCTGCCT	CTCAAAAGGCC	ACCTTGACACA	CCCCGACAGG	CTAGAAATGC	420
ACTCTTCTTC	CCAAAGGTAG	TTGTTCTTGT	TGCCCCAAGCA	NCCTCCANCA	AACCAAAAAC	480
TTGCAAAATC	TGCTCCGTGG	GGGTCATNNN	TACCANGGTT	GGGGAAANAA	ACCCGGCNGN	540
GANCCNCCTT	GTTTGAATGC	NAAGGNAATA	ATCCTCCTGT	CTTGCTTGGG	TGGAANAGCA	600
CAATTGAACT	GTTAACNTTG	GGCCGNGTTC	CNCTNNGGTG	GTCTGAAACT	AATCACCGTC	660
ACTGGAAAAA	GGTANGTGCC	TTCCTTGAAT	TCCCAAANTT	CCCTTNGNTT	TGGGTNTTTT	720
CTCCTCTNCC	CTAAAAATCG	TNTTCCCCCC	CCNTANGGCG			760

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 724 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTAAAAA	CCCCCTCCAT	TGAATGAAAA	60
CTTCCNAAAT	TGTCCAAACC	CCCTNNCCAA	ATNCCGATTT	CCGGGGGGGG	GTTCGAACCC	120

CAAATTAATT	TTGGANTTTA	AATTAAATNT	TNATTNNGGG	AANAANCCAA	ATGTNAAGAA	180
AATTTAAACC	ATTATNAACT	TAAATNCCTN	GAAACCCNTG	GNTTCCAAAA	ATTTTAAACC	240
CTTAAATCCC	TCCGAAATTG	NTAANGGAAA	ACCAAATTCN	CCTAAGGCTN	TTTGAAGGTT	300
NGATTTAAAC	CCCCTTNANT	TNTTTTNACC	CNNGNCTNAA	NTATTTNGNT	TCCGGTGTTT	360
TCCTNTTAAN	CNTNGGTAAC	TCCCGNTAAT	GAAMNNCCCT	AANCCAATTA	AACCGAATTT	420
TTTTTGAATT	GGAAATTCN	NGGGAATTNA	CCGGGGTTTT	TCCCNITTGG	GGGCCATNCC	480
CCCNCTTTCG	GGGTTTGGGN	NTAGGTTGAA	TTTTTNANG	NCCCCAAAAA	NCCCCCAANA	540
AAAAAACTCC	CAAGNNTTAA	TTNGAATNTC	CCCCTTCCCA	GGCCTTTTGG	GAAAGGNGGG	600
TTTNTGGGGG	CCN3GGANTT	CNTTCCCCCN	TTNCCNCCCC	CCCCCNGCT	AAANGGTTAT	660
NGNNTTTGGT	TTTTGGGCCC	CTTNANGGAC	CTTCCGGATN	GAAATTAAAT	CCCCGGGNGC	720
GCCG						724

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 751 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

TTTTTTTTTT	TTTTCTTTG	CTCACATTTA	ATTTTTATTT	TGATTTTTTT	TAATGCTGCA	60
CAACACAATA	TTTATTTTCT	TTGTTTCTTT	TATTTTCAAT	TATTTGTTTG	CTGCTGCTGT	120
TTTATTTATT	TTTACTGAAA	GTGAGAGGGA	ACTTTTGTGG	CCTTTTTTCC	TTTTTCTGTA	180
GGCCGCCTTA	AGCTTTCTAA	ATTTGGAACA	TCTAAGCAAG	CTGAANGGAA	AAGGGGGTTT	240
CGCAAAATCA	CTCGGGGGAA	NGGAAAGGTT	GCTTTGTTAA	TCATGCCCTA	TGGTGGGTGA	300
TTAACTGCTT	GTACAATTAC	NTTTCACTTT	TAATTAATTG	TGCTNAANGC	TTTAATTANA	360
CTTGGGGGTT	CCCTCCCCAN	ACCAACCCCN	CTGACAAAAA	GTGCCNGCCC	TCAAATNATG	420
TCCCGGCNNT	CNTTGAAACA	CACNGCNGAA	NGTTCTCATT	NTCCCCNCNC	CAGGTNAAAA	480
TGAAGGGTTA	CCATNTTTAA	CNCCACCTCC	ACNTGGCINN	GCCTGAATCC	TCNAAAAANC	540
CCCTCAANCN	AATTNCTNNG	CCCCGGTCNC	GCNTNNGTCC	CNCCCGGGCT	CCGGGAANTN	600
CACCCCCNGA	ANNCNNTNNC	NAACNAAATT	CCGAAAATAT	TCCCNNTCNC	TCAATTCCCC	660
CNNAGACTNT	CCTCMNCNAN	CNCAATTTTC	TTTTNNTCAC	GAACNCGNNC	CNNAATATGN	720
NNNCCNCTC	CNCTNGTCCN	NAATCNCCAN	C			751

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 753 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

GTGGTATTTT	CTGTAAGATC	AGGTGTTCTT	CCCTCGTAGG	TTTAGAGGAA	ACACCTCAT	60
AGATGAAAC	CCCCCGAGA	CAGCAGCACT	GCAACTGCCA	AGCAGCCGGG	GTAGGAGGGG	120
CGCCCTATGC	ACAGCTGGGC	CCTTGAGACA	GCAGGGCTTC	GATGTCAGGC	TCGATGTCAA	180
TGGTCTGGAA	GCGGCGGCTG	TACCTGCGTA	GGGGCACACC	GTCAGGGCCC	ACCAGGAACT	240
TCTCAAAGTT	CCAGGCAACN	TCGTTGCGAC	ACACCGGAGA	CCAGGTGATN	AGCTTGGGCT	300

CGGTGATAAN	CGCGGTGSGG	TCGTGCGTGG	GAGGTGGCAG	GGCCTCCCGG	AGGAAGGCNA	360
ATAAAAGGTG	CGTCCCGCA	CGGTTCANCT	CGCACTTCTG	NAANACCATG	ANGTTGGGCT	420
CNAACCCACC	ACGANNCCGG	ACTTCCTTGA	NGGAATTCCG	AAATCTCTTG	GNTCTTGGGC	480
TTCTNCTGAT	GCGCTANCTG	GTTGCCCCNG	ATGCCAANCA	NCCCCAANCC	CGGGGGTCCT	540
AAANCACCCN	CCTCCTCNTT	TCATCTGGGT	TNTTNTCCCG	GGACCNTGGT	TCCTCTCAAG	600
GGANCCDATA	TCTCNACCAW	TACTCACCN	NCCCCCCCC	GNNACCCANC	CTTCTANNGN	660
TTCCCNCCCG	NCCTCTGGCC	CNTCAAANAN	GCTTNCACNA	CCTGGGTCTG	CCTTCCCCCC	720
TNCCCTATCT	GNACCCCN	TTTGTCTCAN	TNT			753

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 341 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

ACTATATCCA	TCACAACAGA	CATGCTTCAT	CCCATAGACT	TCTTGACATA	GCTTCAAAATG	60
AGTGAACCCA	TCCTTGATTT	ATATACATAT	ATGTTCTCAG	TATTTTGGGA	GCCTTTCCAC	120
TTCTTTAAAC	CTTGTTTCATT	ATGAACACTG	AAAATAGGAA	TTTGTGAAGA	GTTAAAAAGT	180
TATAGCTTGT	TTACGTAGTA	AGTTTTTGAA	GTCTACATTC	AATCCAGACA	CTTAGTTGAG	240
TGTTAAACTG	TGATTTTTTAA	AAAATATCAT	TTGAGAATAT	TCTTTCAGAG	GTATTTTCAT	300
TTTTACTTTT	TGATTAAATTG	TGTTTTATAT	ATTAGGGTAG	T		341

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 101 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

ACTTACTGAA	TTTAGTTCTG	TGCTCTTCCT	TATTTAGTGT	TGTATCATAA	ATACTTTGAT	60
GTTTCAAACA	TTCTAAATAA	ATAATTTTCA	GTGGCTTCAT	A		101

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 305 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

ACATCTTTGT TACAGTCTAA GATGTGTTCT TAAATCACCA TTCCTTCCTG GTCCTCACCC	60
TCCAGGGTGG TCTCAGCTG TAATTAGAGC TATTGAGCAG TCTTTACAGC AAATTAAGAT	120
TCAGATGCCT TGCTAAGTCT AGAGTTCTAG AGTTATGTTT CAGAAAGTCT AAGAAACCCA	180
CCTCTTGAGA GGTCACTAAA GAGGACTTAA TATTTTCATAT CTACAAAATG ACCACAGGAT	240
TGGATACAGA ACGAGAGTTA TCCTGGATAA CTCAGAGCTG AGTACCTGCC CGGGGGCCGC	300
TCGAA	305

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 852 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

ACATAAATAT CAGAGAAAAG TAGTCTTTGA AATATTTACG TCCAGGAGTT CTTTGTTTCT	60
GATTATTTGG TGTGTGTTTT GGTITGTGTC CAAAGTATTG GCAGCTTCAG TTTTCATTTT	120
CTCTCCATCC TCGGGCATTG TCCCAAAT TATATACCAG TCTTCGTCCA TCCACACGCT	180
CCAGAAATTC TCTTTTGTAG TAATATCTCA TAGCTCGGCT GAGCTTTTCA TAGGTCATGC	240
TGCTGTTGTT CTTCTTTTTA CCCCATAGCT GAGCCACTGC CTCTGATTTT AAGAACCTGA	300
AGACGCCCTC AGATCGGTCT TCCCATTTTA TTAATCCTGG GTTCTTGTCT GGGTTCAAGA	360
GGATGTCGCG GATGAATTC CATAAGTGAG TCCCTCTCGG GTTGTGCTTT TTGGTGTGGC	420
ACTTGGCAGG GGGGTCTTGC TCCTTTTCA TATCAGGTGA CTCTGCAACA GGAAGGTGAC	480
TGGTGGTGT CATGGAGATC TGAGCCCGGC AGAAAGTTTT GCTGTCCAAC AAATCTACTG	540
TGCTACCATA GTTGGTGTCA TATAAATAGT TCTNGTCTTT CCAGGTGTTT ATGATGGAAG	600
GCTCAGTTTG TTCAGTCTTG ACAATGACAT TGTGTGTGGA CTGGAACAGG TCACTACTGC	660
ACTGGCCGTT CCACTTCAGA TGCTGCAAGT TGCTGTACAC GAGNTGCCCC GCCGTCCCTG	720
CCGCCCCGGT GAACTCCTGC AAACATCATG TGCAAAGGTG CTCGCCGTTG ATGTCGAAC	780
CNTCCAAAGC CATACAATTG GCATCCAGCT GGTGTTGTG CAGGAGGTGA TGGAGCCACT	840
CCCACACCTG GT	852

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 234 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(i) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

ACAACAGACC CTTGCTCGCT AACGACCTCA TGCTCATCAA GTTGGACGAA TCCGTGTCCG	60
AGTCTGACAC CATCCGGAGC ATCAGCATTG CTTCCGAGTG CCCTACCGCG GGGAACTCTT	120
GCCTCGTTTC TGGCTGGGGT CTGCTGGCGA ACGGCAGAAAT GCCTACCGTG CTGCAGTGCG	180
TGAACGTGTC GGTGCTGTCT GAGGAGGTCT GCAGTAAGCT CTATGACCCG CTGT	234

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 590 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

ACTTTTTATT TAAATGTTTA TAAGGCAGAT CTATGAGAAT GATAGAAAAC ATGGTGTGTA	60
ATTTGATAGC AATATTTTGG AGATTACAGA GTTTTAGTAA TTACCAATTA CACAGTAAA	120
AAGAAGATAA TATATTCCAA GCANATACAA AATATCTAAT GAAAGATCAA GGCAGGAAAA	180
TGANTATAAC TAATTGACAA TGGAAAATCA ATTTTAATGT GAATTGCACA TTATCCTTTA	240
AAAGCTTTCA AAANAAANAA TTATTGCAGT CTANTTAATT CAAACAGTGT TAAATGGTAT	300
CAGGATAAAN AACTGAAGGG CANAAAGAAT TAATTTTCAC TTCATGTAAC NCACCCANAT	360
TTACAATGGC TTAAATGCAN GGAAAAAGCA GTGGAAGTAG GGAAGTANTC AAGGCTCTTC	420
TGGTCTCTAA TCTGCCTTAC TCTTTGGGTG TGGCTTTGAT CCTCTGGAGA CAGCTGCCAG	480
GGCTCCTGTT ATATCCACAA TCCCAJCGAC AAGATGAAGG GATGAAAAAG GACACATGCT	540
GCCTTCCTTT GAGGAGACTT CATCTCACTG GCCAACACTC AGTCACATGT	590

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 774 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

ACAAGGGGGC ATAATGAAGG AGTGGGGANA GATTTTAAAG AAGGAAAAAA AACGAGGGCC	60
TGAACAGAAT TTTCTGNAG AACGGGGCTT CAAATAAATT TTCTTGGGA GTTCAAGAC	120

GCTTCACTGC TTGAAACTTA AATGGATGTG GGACANAATT TTCTGTAATG ACCCTGAGGG	180
CATTACAGAC GGGACTCTGC GAGGAAGGAT AAACAGAAAG GGGACAAAGC CTAATCCCCAA	240
AACATCAAAG AAAGGAAGGT GGCCTCATAC CTCCCAGCCT ACACAGTTCT CCAGGGCTCT	300
CCTCATCCCT CGAGGACGAC AGTGGAGGAA CAACTGACCA TGTCCCCAGG CTCTGTGTG	360
CTGGCTCCTG GTCTTCAGCC CCCAGCTCTG GAAGCCCACC CTCTGCTGAT CCTGCGTGGC	420
CCACACTCCT TGAACACACA TCCCCAGGTT ATATTCTCTG ACATGGCTGA ACCTCCTATT	480
CCTACTTCCG AGATGCCTTG CTCCCTGCAG CCTGTCAAAA TCCCCTCAC CCTCCAAACC	540
ACGGCATGGG AAGCCTTTCT GACTTGCCTG ATTACTCCAG CATCTTGGA CAATCCCTGA	600
TTCCCCACTC CTTAGAGGCA AGATAGGGTG GTTAAGAGTA GGGCTGGACC ACTTGGAGCC	660
AGGCTGCTGG CTTCAAATTN TGGCTCATTT ACGAGCTATG GGACCTTGG CAAGTNATCT	720
TCACTTCTAT GGGCNTCATT TTGTTCTACC TGCAAAATGG GGGATAATAA TAGT	774

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 124 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

CANAAATTGA AATTTTATAA AAAGGCATTT TTCTCTTATA TCCATAAAAT GATATAATTT	60
TTGCAANTAT ANAAATGTGT CATAAATTAT AATGTTCTTT AATTACAGCT CAACGCAACT	120
TGGT	124

(2) INFORMATION FOR SEQ ID NO:49:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 147 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

GCCGATGCTA CTATTTTATT GCAGGAGGTG GGGCTGTTTT TATTATTCTC TCAACAGCTT	60
TGTGGCTACA GGTGGTGTCT GACTGCATNA AAAANTTTTT TACGGGTGAT TGCAAAAATT	120
TTAGGGCACC CATATCCCCA GCANTGT	147

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 107 base pairs
- (B) TYPE: nucleic acid

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

ACATTAAATT AATAAAAGGA CTGTTGGGGT TCTGCTAAAA CACATGGCTT GATATATTGC	60
ATGGTTTGAG GTTAGGAGGA GTTAGGCATA TGTTTGGGA GAGGGGT	107

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 204 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

GTCCTAGGAA GTCTAGGGGA CACACGACTC TGGGGTCACG GGGCCGACAC ACTTGCACGG	60
CGGGAAGGAA AGGCAGAGAA GTGACACCGT CAGGGGGAAA TGACAGAAAG GAAAATCAAG	120
GCCTTGCAAG GTCAGAAAGG GGACTCAGGG CTTCACCAC AGCCCTGCC CACTTGGCCA	180
CCTCCCTTTT GGGACCAGCA ATGT	204

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 491 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

ACAAAGATAA CATTTATCTT ATAACAAAAA TTGATAGTT TTAAAGGTTA GTATTGTGTA	60
GGGTATTTTC CAAAAGACTA AAGAGATAAC TCAGGTAAAA AGTTAGAAAT GTATAAAACA	120
CCATCAGACA GGTTTTTAAA AAACAACATA TTACAAAATT AGACAATCAT CCTTAAAAAA	180
AAAACCTCTT GTATCAATTT CTTTGTGTTA AAATGACTGA CTTAANTATT TTTAAATATT	240
TCANAAACAC TTCCTCAAAA ATTTTCAANA TGGTAGCTTC CANATGTNCC CTCAGTCCCA	300
ATGTTGCTCA GATAAATAAA TCTCGTGAGA ACTTACCACC CACCACAAGC TTTCTGGGGC	360
ATGCAACAGT GTCTTTTCTT TNCCTTTTCT TTTTITTTTT TTACAGGCAC AGAAACTONT	420

CAATTTTATT TGGATAACAA AGGGTCTCCA AATTATATTG AAAAATAAAT CCAAGTTAAT 480
ATCACTCTTG T 491

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 484 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

ACATAATTTA GCAGGGCTAA TTACCATAAG ATGCTATTTA TTAANAGGTN TATGATCTGA 60
GTATTAACAG TTGCTGAAGT TTGGTATTTT TATGCAGCAT TTCTTTTGTG CTTTGATAAC 120
ACTACAGAAC CCTTAAGGAC ACTGAAAATT AGTAAGTAAA GTTCAGAAAC ATTAGCTGCT 180
CAATCAAATC TCTACATAAC ACTATAGTAA TTA AACCGTT AAAAAAAGT GTTGAAATCT 240
GCACTAGTAT ANACCGCTCC TGTCAGGATA ANACTGCTTT GGAACAGAAA GGGAAAAANC 300
AGCTTTGANT TTCTTTGTGC TGATANGAGG AAAGGCTGAA TTACCTTGTT GCCTCTCCCT 360
AATGATTGGC AGGTCNGGTA AATNCCAAAA CATATTCCAA CTCAACACTT CTTTTCCNCG 420
TANCTTGANT CTGTGTATTC CAGGANCAAG CCGATGGAAT GGGCCAGCCC NCGGATGTTC 480
CANT 484

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 151 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

ACTAAACCTC GTGCTTGTTGA ACTCCATACA GAAAACGGTG CCATCCCTGA ACACGGCTGG 60
CCACTGGGTA TACTGCTGAC AACCGCAACA AAAAAACAC AAATCCTTGG CACTGGCTAG 120
TCTATGTCCT CTCAAGTGCC TTTTGTGTTG T 151

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 91 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

60

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

```
ACCTGGGCTTG TCTCCGGGTG GTTCCCGGGG CCCCCACGG TCCCCAGAAC GGACACTTTC 60
GCCCTCCACT GGATACTCGA GCCAAACTGG T 91
```

(2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 133 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

```
GGCGGATGTG CGTTGGTTAT ATACAAATAT GTCATTTTAT GTAAGGGACT TGAGTATACT 60
TGGATTTTGT GTATCTGTGG GTTGGGGGGA CGGTCCAGGA ACCAATACCC CATGGATACC 120
AAGGGACAAC TGT 133
```

(2) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 147 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

```
ACTCTGGAGA ACCTGAGCCG CTGCTCCGCC TCTGGGATGA GGTGATGCAN GCNGTGGCGC 60
GACTGGGAGC TGAGCCCTTC CCTTTGCCGC TGCCCTCAGAG GATTGTTGCC GACNTGCANA 120
TCTCANIGGG CTGGATNCAT GCAGGGT 147
```

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 198 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

ACAGGGATAT AGGTTTNAAG TTATTGTNAT TGTAATAATAC ATTGAATTTT CTGTATACTC	60
TGATTACATA CATTTATCCT TTAAAAAAGA TGTAATCTT AATTTTATG CCATCTATTA	120
ATTTACCAAT GACTTACCTT GTAAATGAGA AGTCATGATA GCACTGAATT TTAAC TAGTT	180
TTGACTTCTA AGTTTGGT	198

(2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 330 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

ACAACAAATG GGTGTGAGG AAGTCTTATC AGCAAACTG GTGATGGCTA CTGAAAAGAT	60
CCATTGAAAA TTATCATTA TGATTTTAAA TGACAAGTTA TCAAAACTC ACTCAATTTT	120
CACCTGTGCT AGCTTGCTAA AATGGGAGTT AACTCTAGAG CAAATATAGT ATCTTCTGAA	180
TACAGTCAAT AAATGACAAA GCCAGGGCCT ACAGGTGGTT TCCAGACTTT CCAGACCCAG	240
CAGAAGGAAT CTATTTTATC ACATGGATCT CCGTCTGTGC TCAAAATACC TAATGATATT	300
TTTCGTCTTT ATTGGAAGAGT	330

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 175 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

ACCGTGGGTG CTTTCTACAT TCCTGACGGC TCCTTCACCA ACATCTGGTT CTACTTCGGC	60
GTGCTGGGCT CTTTCTCTT CATCCTCATC CAGCTGGTGC TGCTCATCGA CTTTGCGCAC	120
TCCTGGAACC AGCGGTGGCT GGGCAAGGCC GAGGAGTGCG ATTCCCGTGC CTGGT	175

(2) INFORMATION FOR SEQ ID NO:61:

62

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 154 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

ACCCCACTTT TCCTCCTGTS AGCAGTCTGG ACTTCTCACT GCTACATGAT GAGGGTGAGT	60
GGTGGTTGCT CTCCAACAGT ATCCTCCCTT TTCGGGATCT GCTGAGCCCG ACAGCAGTCC	120
TGGACTGCAC AGCCGCGGGG CTCACATTG CTGT	154

(2) INFORMATION FOR SEQ ID NO:62:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

CGCTCGAGCC CTATAGTGAG TCGTATTAGA	30
----------------------------------	----

(2) INFORMATION FOR SEQ ID NO:63:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 89 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

ACAAGTCATT TCAGCACCTT TTGCTCTTCA AAAGTGACCA TCTTTTATAT TTAATGCTTC	60
CTGTATGAAT AAAAATGGTT ATGTCAAGT	89

(2) INFORMATION FOR SEQ ID NO:64:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 97 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

ACCGGAGTAA CTGAGTCGGG ACGCTGAATC TGAATCCACC AATAAATAAA GGTTCCTGCAG	60
AATCAGTGCA TCCAGGATTG GTCCTTGGAT CTGGGGT	97

(2) INFORMATION FOR SEQ ID NO:65:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 377 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

ACAACAANAA NTCCCTTCTT TAGGCCACTG ATGGAAACCT GGAACCCCTT TTTGATGGCA	60
GCATGGCGTC CTAGGCCTTG ACACAGCGGC TGGGGTTTGG GCTNTCCCAA ACCGCACACC	120
CCAACCCCTGG TCTACCCACA NTTCTGGCTA TGGGCTGTCT CTGCCACTGA ACATCAGGGT	180
TCGGTTCATAA NATGAAATCC CAANGGGGAC AGAGGTCAGT AGAGGAAGCT CAATGAGAAA	240
GGTGCTGTTT GCTCAGCCAG AAAACAGCTG CCTGGCATTG CCCCCTGAAC TATGAACCCC	300
TGGGGGTGAA CTACCCCCAN GAGGAATCAT GCCTGGGCGA TGCAANGGTG CCAACAGGAG	360
GGGCGGGAGG AGCATGT	377

(2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 305 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

ACGCCTTTCC CTCAGAATTC AGGGAAGAGA CTGTGGCCTG CCTTCTCTCCG TTGTTGCGTG	60
--	----

64

AGAAGCCGTC	TGCCCTTCC	CACCATATCC	ACCCTCGCTC	CATCTTTGAA	CTCAAACACG	120
AGGAACTAAC	TGACCCCTGG	TCCTCTCCCC	AGTCCCCAGT	TCACCCTCCA	TCCCTCACCT	180
TCCTCCACTC	TAAGGGATAT	CAACACTGCC	CAGCACAGGG	GCCCTGAATT	TATGTGGTTT	240
TTATATATTT	TTTAATAAGA	TGCACCTTAT	GTCAATTTTT	AATAAAGTCT	GAAGAATTAC	300
TGTTT						305

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 385 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

ACTACACACA	CTCCACTTGC	CCTTGTGAGA	CACTTTGTCC	CAGCACTTIA	GGAATGCTGA	60
GGTCGGACCA	GCCACATCTC	ATGTGCAAGA	TTGCCACGCA	GACATCAGGT	CTGAGAGTTC	120
CCCTTTTAAA	AAAGGGGACT	TGCTTAAAAA	AGAAGTCTAG	CCACGATTGT	GTAGAGCAGC	180
TGTGCTGTGC	TGGAGATTCA	CTTTTGAGAG	AGTTCTCCTC	TGAGACCTGA	TCTTTAGAGG	240
CTGGGCAGTC	TTGCACATGA	GATGGGGCTG	GTCTGATCTC	AGCACTCCTT	AGTCTGCTTG	300
CCTCTCCCAG	GGCCCCAGCC	TGGCCACACC	TGCTTACAGG	GCACTCTCAG	ATGCCCATAC	360
CATAGTTTCT	GTGCTAGTGG	ACCGT				385

(2) INFORMATION FOR SEQ ID NO:68:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 73 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

ACTTAACCAG	ATATATTTTT	ACCCACAGATG	GGGATATTCT	TTGTAAAAAA	TGAAAATAAA	60
GTTTTTTTAA	TGG					73

(2) INFORMATION FOR SEQ ID NO:69:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 536 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69.

ACTAGTCCAG TGTGGTGGAA TTCCATTGTG TTGGGGGCTC TCACCCTCCT CTCCTGCAGC	60
TCCAGCTTTG TGCTCTGCCT CTGAGGAGAC CATGGCCCAG CATCTGAGTA CCCTGCTGCT	120
CCTGCTGGCC ACCCTAGCTG TGGCCCTGGC CTGGAGCCCC AAGGAGGAGG ATAGGATAAT	180
CCCGGGTGGC ATCTATAACG CAGACCTCAA TGATGAGTGG GTACAGCGTG CCCTTCACTT	240
CGCCATCAGC GAGTATAACA AGGCCACCAA AGATGACTAC TACAGACGTC CGCTGCGGGT	300
ACTAAGAGCC AGGCAACAGA CCGTTGGGGG GGTGAATTAC TTCTTCGACG TAGAGGTGGG	360
CCGAACCATA TGTACCAAGT CCCAGCCCCA CTTGGACACC TGTGCCTTCC ATGAACAGCC	420
AGAACTGCAG AAGAAACAGT TGTGCTCTTT CGAGATCTAC GAAGTTCCTT GGGGAGAACA	480
GAANGTCCCT GGGTGAAATC CAGGTGTCAA GAAATCCTAN GGATCTGTTG CCAGGC	536

(2) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 477 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

ATGACCCCTA ACAGGGGCCC TCTCAGCCCT CCTAATGACC TCCGGCCTAG CCATGTGATT	60
TCACTTCCAC TCCATAACGC TCCTCATACT AGGCCTACTA ACCAACACAC TAACCATATA	120
CCAATGATGG CGCGATGTAA CAGGAGAAAG CACATACCAA GGCCACCACA CACCACCTGT	180
CCAAAAAGGC CTTCGATACG GGATAATCCT ATTTATTACC TCAGAAGTTT TTTTCTTCGC	240
AGGGATTTTT CTGAGCCTTT TACCACTCCA GCCTAGCCCC TACCCCCCAA CTAGGAGGGC	300
ACTGGCCCCC AACAGGCATC ACCCCGCTAA ATCCCTTAGA AGTCCCCTC CTAAACACAT	360
CCGTATTACT CGCATCAGGA GTATCAATCA CCTGAGCTCA CCATAGTCTA ATAGAAAACA	420
ACCGAAACCA AATTATTCAA AGCACTGCTT ATTACAATTT TACTGGGTCT CTATTTT	477

(2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 533 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

AGAGCTATAG	GTACAGTGTG	ATCTCAGCTT	TGCAAACACA	TTTTCTACAT	AGATAGTACT	60
AGGTATTAA	AGATATGTAA	AGAAAGAAAT	CACACCATTA	ATAATGGTAA	GATTGGTTTA	120
TGTGATTTTA	GTGGTATTTT	TGGCACCCCT	ATATATGTTT	TCCAAACTTT	CAGCAGTGAT	180
ATTATTTCCA	TAACTTAAAA	AGTGAATTTG	AAAAAGAAAA	TCTCCAGCAA	GCATCTCATT	240
TAAATAAAGG	TTTGTTCATC	TTAAAAATAC	AGCAATATGT	GACTTTTTTA	AAAAGCTGTC	300
AAATAGGTTG	GACCTACTA	ATAATTATTA	GAAATACATT	TAAAAACATC	GAGTACCTCA	360
AGTCAGTTTG	CCTTGAAAAA	TATCAAATAT	AACTCTTAGA	GAAATGTACA	TAAAAGAATG	420
CTTCGTAAAT	TTGGAGTANG	AGGTTCCCTC	CTCAATTTTG	TATTTTAAAA	AAGTACATGG	480
TAAAAAATAA	AATTCACAAC	AGTATATAAG	GCTGTAAAT	GAAGAATTCT	GCC	533

(2) INFORMATION FOR SEQ ID NO:72:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 511 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

TATTACGGAA	AAACACACCA	CATAATTCAA	CTANCAAAGA	ANACTGCTTC	AGGGCGTGTA	60
AAATGAAAGG	CTTCCAGGCA	GTTATCTGAT	TAAAGAACAC	TAAAAGAGGG	ACAAGGCTAA	120
AAGCCGCAGG	ATGTCTACAC	TATANCAGGC	GCTATTTGGG	TTGGCTGGAG	GAGCTGTGGA	180
AAACATGGAN	AGATTGGTGC	TGGANATCGC	CGTGGCTATT	CCTCATTGTT	ATTACANAGT	240
GAGGTTCTCT	GTGTGCCCAC	TGGTTTGAAA	ACCGTTCCTC	AATAATGATA	GAATAGTACA	300
CACATGAGAA	CTGAAATGGC	CCAAACCCAG	AAAGAAAGCC	CAACTAGATC	CTCAGAAATC	360
GCTTCTAGGG	ACAATAACCG	ATGAAGAAAA	GATGGCCTCC	TTGTGCCCCC	GTCTGTTATG	420
ATTTCTCTCC	ATTGCAGCNA	NAAACCCGTT	CTTCTAAGCA	AACNCAGGTG	ATGATGGCNA	480
AAATACACCC	CCTCTTGAAG	NACCNGGAGG	A			511

(2) INFORMATION FOR SEQ ID NO:73:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 499 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

CAGTGGCAGC	ACTGGTGCCA	GTACCACTAC	CAATAACAGT	GCCAGTGCCA	GTGCCAGCAC	60
CAGTGGTGGC	TTCAGTGCTG	GTGCCAGCCT	GACCGCCACT	CTCAGATTTG	GGCTCTTCGC	120
TGGCCTTGGT	GGAGCTGGTG	CCAGCACCAG	TGGCAGCTCT	GGTGCTGTG	GTTTCTCTTA	180
CAAGTGAGAT	TTTAGATATT	GTTAATCCTG	CCAGTCTTTT	TCTTCAAGCC	AGGCTGCATC	240

CTCAGAAACC TACTCAACAC AGCACTCTAG GCAGCCACTA TCAATCAATT GAAGTTGACA	300
CTCTGCATTA AATCTATTTG CCATTTCTGA AAAAAAAAAA AAAAAAAGGG CGGCCGCTCG	360
ANTCTAGAGG GCGCGTTTAA ACCCGCTGAT CAGCCTCGAC TGTGCCCTTCT ANTTGCCAGC	420
CATCTGTTGT TTGCCCCCTCC CCCGNTGCCT TCCTTGACCC TGGAAAGTGC CACTCCCACT	480
GTCCCTTTCCT AANTAAAAT	499

(2) INFORMATION FOR SEQ ID NO:74:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 537 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

TTTCATAGGA GAACACACTG AGGAGATACT TGAAGAATTT GGATTCAGCC GCGAAGAGAT	60
TTATCAGCTT AACTCAGATA AAATCATTGA AAGTAATAAG GTAAAAGCTA GTCTCTAACT	120
TCCAGGCCCA CGGCTCAAGT GAATTTGAAAT ACTGCATTTA CAGTGTAGAG TAACACATAA	180
CATTGTATGC ATGGAAACAT GGAGGAACAG TATTACAGTG TCCTACCACT CTAATCAAGA	240
AAAGAATTAC AGACTCTGAT TCTACAGTGA TGATTGAATT CTAAAAATGG TAATCATTAG	300
GGCTTTTGAT TTATAANACT TTGGGTACTT ATACTAAAT ATGGTAGTTA TACTGCCTTC	360
CAGTTTGCTT GATATATTTG TTGATATTAA GATTCTTGAC TTATATTTTG AATGGGTCTT	420
ACTGAAAAAN GAATGATATA TTCTTGAAGA CATCGATATA CATTATTTTA CACTCTTGAT	480
TCTACAATGT AGAAAATGAA GGAAATGCCC CAAATTGTAT GGTGATAAAA GTCCCGT	537

(2) INFORMATION FOR SEQ ID NO:75:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 467 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

CAAAACAAAT TTTTCAAAAG ATGCAAAATGA TACACTACTG CTGCAGCTCA CAAACACCTC	60
TGCATATTAC ACGTACCTCC TCCTGCTCCT CAAGTAGTGT GGTCTATTTT GCCATCATCA	120
CCTGCTGTCT GCTTAGAAGA ACGGCTTTCT GCTGCAANGG AGAGAAATCA TAACAGACGG	180
TGGCACAAGG AGGCCATCTT TTCCTCATCC GTTATTGTCC CTAGAAGCGT CTTCTGAGGA	240
TCTAGTTGGG CTTTCTTTCT GGSTTTGGGC CATTTCANTT CTCATGTGTG TACTATTCTA	300
TCATTATTGT ATAACGGTTT TCAAACCNGT GGGCACNCAG AGAACCTCAC TCTGTAATAA	360
CAATGAGGAA TAGCCACGGT GATCTCCAGC ACCAAATCTC TCCATGTTNT TCCAGAGCTC	420
CTCCAGCCAA CCCAAATAGC CGCTGCTATN GTGTAGAACA TCCCTGN	467

(2) INFORMATION FOR SEQ ID NO:76:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 400 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

AAGGTGACAG	CATTCCGGGCC	GAGATGTCTC	GCTCCGTGGC	CTTAGCTGTG	CTCGCGCTAC	60
TCTCTCTTTC	TGGCCTGGAG	GCTATCCAGC	GTACTCCAAA	GATTCAGGTT	TACTCACGTC	120
ATCCAGCAGA	GAATGGAAAG	TCAAATTTCC	TGAATTGCTA	TGTGTCTGGG	TTTCATCCAT	180
CCGACATTGA	AGTTGACTTA	CTGAAGAATG	GAGAGAGAAT	TGAAAAAGTG	GAGCATTCAg	240
ACTTGTCTTT	CAGCAAGGAC	TGGTCTTTCT	ATCTCTTGTA	CTACACTGAA	TTCACCCCCA	300
CTGAAAAAGA	TGAGTATGCC	TGCCGTGTGA	ACCATGTGAC	TTTGTCACAG	CCCAAGATNG	360
TTNAGTGGGA	TCGANACATG	TAAGCAGCAN	CATGGGAGGT			400

(2) INFORMATION FOR SEQ ID NO:77:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 248 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

CTGGAGTGCC	TTGGTGTTC	AAGCCCCTGC	AGGAAGCAGA	ATGCACCTTC	TGAGGCACCT	60
CCAGCTGCCC	CGGCGGGGA	TGCGAGGCTC	GGAGCACCCCT	TGCCCCGGCTG	TGATTGCTGC	120
CAGGCACTGT	TCATCTCAGC	TTTTCTGTCC	CTTTGCTCCC	GGCAAGCGCT	TCTGCTGAAA	180
GTTCATATCT	GGAGCCTGAT	GTCTTAACGA	ATAAAGGTCC	CATGCTCCAC	CCGAAAAAAA	240
AAAAAAA						248

(2) INFORMATION FOR SEQ ID NO:78:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 201 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

69

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

ACTAGTCCAG TGTGGTGGAA TTCCATTGTG TTGGGCCCCAA CACAATGGGT ACCTTTAACA	60
TCACCCAGAC CCGCCCTGC CCGTGCCCCA CGCTGCTGCT AACGACAGTA TGATGCTTAC	120
TCTGCTACTC GGAAACTATT TTTATGTAAT TAATGTATGC TTTCTTGTTT ATAAATGCCCT	180
GATTTAAAAA AAAAAAAAAA A	201

(2) INFORMATION FOR SEQ ID NO:79:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 552 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

TCCTTTTGGT AGGTTTTTGA GACAACCCTA GACCTAAACT GTGTCACAGA CTTCTGAATC	60
TTTAGGCAGT GCTAGTAATT TCCTCGTAAT GATTCTGTTA TTACTTTCCT ATTCTTTATT	120
CCTCTTTCTT CTGAAGATTA ATGAAGTTGA AAATTGAGGT GGATAAATAC AAAAAGGTAG	180
TGTGATAGTA TAAGTATCTA AGTGCAGATG AAAGTGTGTT ATATATATCC ATTCAAAATT	240
ATGCAAGTTA GTAATTACTC AGGGTTAACT AAATTACTTT AATATGCTGT TGAACCTACT	300
CTGTTCCCTG GCTAGAAAAA ATTATAAACA GGACTTTGTT AGTTTGGGAA GCCAAATTGA	360
TAATATTCTA TGTCTAAAA GTTGGGCTAT ACATAAANTA TNAAGAAATA TGGAAATTTTA	420
TTCCCAGGAA TATGGGGTTC ATTTATGAAT ANTACCCGGG ANAGAAGTTT TGANTNAAAC	480
CNGTTTGGT TAATACGTTA ATATGTCTN AATNAACAAG GCNTGACTTA TTTCCAAAAA	540
AAAAAAAAAA AA	552

(2) INFORMATION FOR SEQ ID NO:80:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 476 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

ACAGGGATTT GAGATGCTAA GGCCCCAGAG ATCGTTTGAT CCAACCTCT TATTTTCAGA	60
GGGCAAAATG GGGCCTAGAA GTTACAGAGC ATCTAGGTGG TGGCTGGCA CCCCTGGCCT	120
CACACAGACT CCCGAGTAGC TGGGACTACA GGCACACAGT CACTGAAGCA GGCCTGTTT	180
GCAATTCAGG TTGCCACCTC CAACTTAAAC ATCTTCATA TGTGATGTCC TTAGTCACTA	240
AGGTTAAACT TTCCACCCA GAAAAGGCA CTTAGATAAA ATCTTAGAGT ACTTTCATAC	300

```

TUTTTCTAAGT CTTCTTTCAG CCTCACTTTT AGTCTCTCTT GGGGGTTGAT AGGAANTNTC      360
TCTTGGCTTT CTCAATAAAA TCTCTATCCA TCTCATGTTT AATTTCGTAC GCTAAAAAT      420
GCTGAAAAAA TTAAATCTT CTGCTTTTNC TTTAAAAAA AAAAAAAA AAAAAA      486

```

(2) INFORMATION FOR SEQ ID NO:81:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 232 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

```

TTTTTTTTTG TATGCCNTCN CTGTGGNGTT ATTGTTGCTG CCACCCTGGA GGAGCCCAGT      60
TTCTTCTGTA TCTTTCTTTT CTGGGGGATC TTCCTGGCTC TGCCCCCTCCA TTCCCAGCCT      120
CTCATCCCA TCTTGCACTT TTGTAGGGT TGGAGGCGCT TTCCTGGTAG CCCCTCAGAG      180
ACTCAGTCAG CGGGAATAAG TCCTAGGGGT GGGGGGTGTG GCAAGCCGGC CT      232

```

(2) INFORMATION FOR SEQ ID NO:82:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

```

AGGCGGGAGC AGAAGCTAAA GCCAAAGCCC AAGAAGAGTG GCAGTGCCAG CACTGGTGCC      60
AGTACCAGTA CCAATAACAT GCCAGTGCCA GTGCCAGCAC CAGTGGTGGC TTCAGTGCTG      120
GTGCCAGCCT GACCGCCACT CTCACATTTG GGCTCTTCGC TGGCCTTGGT GGAGCTGGTG      180
CCAGCACCAG TGGCAGCTCT GGTGCCTGTG GTTTCTCCTA CAAGTGAGAT TTTAGATATT      240
GTAAATCCTG CCAGTCTTTC TCTTCAAGCC AGGGTGCATC CTCAGAAACC TACTCAACAC      300
AGCACTCTNG GCAGCCACTA TCAATCAATT GAAGTTGACA CTCTGCATTA AATCTATTTG      360
CCATTTCAAA AAAAAAAAAA AAA      383

```

(2) INFORMATION FOR SEQ ID NO:83:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 494 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

ACCGAATTGG GACCGCTGGC TTATAAGCGA TCATGTCCTC CAGTATTACC TCAACGAGCA	60
GGGAGATCGA GTCTATACGC TGAAGAAATT TGACCCGATG GGACAACAGA CCTGCTCAGC	120
CCATCCTGCT CGGTTCTCCC CAGATGACAA ATACTCTCGA CACCGAATCA CCATCAAGAA	180
ACGCTTCAAG GTGCTCATGA CCCAGCAACC GCGCCCTGTC CTCTGAGGCT CCTTAAACTG	240
ATGTCTTTTC TGCCACCTGT TACCCCTCGG AGACTCCGTA ACCAAACTCT TCGGACTGTG	300
AGCCCTGATG CCTTTTGGCC AGCCATACTC TTTGGCNTCC AGTCTCTCGT GGCGATTGAT	360
TATGCTTGTG TGAGGCAATC ATGGTGGCAT CACCCATNAA GGGAACACAT TTGANTTTTT	420
TTTCNCATAT TTTAAATTAC NACCAGAATA NTTCAGAATA AATGAATTGA AAAACTCTTA	480
AAAAAAAAAA AAAA	494

(2) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 380 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

GCTGGTAGCC TATGGCGTGG CCACGGANGG GCTCCTGAGG CACGGGACAG TGACTTCCCA	60
AGTATCCTGC GCCGCGTCTT CTACCGTCCC TACCTGCAGA TCTTCGGGCA GATTCCCCAG	120
GAGGACATGG ACGTGGCCCT CATGGAGCAC AGCAACTGCT CGTCGGAGCC CGGCTTCTGG	180
GCACACCCTC CTGGGGCCCA GGCGGGCACC TCGTCTCTCC AGTATGCCAA CTGGCTGGTG	240
GTGCTGCTCC TCGTCATCTT CCTGCTCGTG GCCAACATCC TGCTGGTCAC TTGCTCATTG	300
CCATGTTTCA TTACACATTC GGCAAAGTAC AGGGCAACAG CNATCTCTAC TGGGAAGGCC	360
AGCGTTNCCG CCTCATCCGG	380

(2) INFORMATION FOR SEQ ID NO:85:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 481 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

GASTTAGCTC	CTCCACAAAC	TTGATGAGGT	CGTCTGCAGT	GGCCTCTGGC	TTCAIACCCG	60
TNCCATCGTC	ATACTGTAGG	TTTGCCACCA	CGTCCTGCAT	CTTGGGGGGG	CTAATATCGA	120
GGAAACCTCT	AATCAAGTCA	CCGTGNATNA	AACCTGTGGC	TGGTTCTGTC	TTCCCTCTCG	180
TGTGAAAGGA	TCTCCAGAA	GAGTGCTCGA	TCTTCCCCAC	ACTTTTGATG	ACTTTATTGA	240
GTCGATTCTG	CATGTCCAGC	AGGAGGTTGT	ACCAGCTCTC	TGACAGTGAG	GTCACCCAGCC	300
CTATCATGCC	NTTGAACGTG	CCGAAGAACA	CCGAGCCTTG	TGTGGGGGGT	GNAGTCTCAC	360
CCAGATTCTG	CATTACCAGA	NAGCCGTGCG	AAAAGANATT	GACAACTCGT	CCAGGNNGAA	420
AAAGAACACC	TCCTGGAAGT	GCTNGCCGCT	CTCGTCCNT	TGGTGGNNGC	GCTNCCCTTT	480
T						491

(2) INFORMATION FOR SEQ ID NO:86:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 472 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

AACATCTTCC	TGTATAATGC	TGIGTAATAT	CGATCCGATN	TTGTCTGCTG	AGAATTCAAT	60
ACTTGGAAAA	GCAACTTNAA	GCCTGGACAC	TGGTATTAAA	ATTCACAATA	TGCAACACTT	120
TAAACAGTGT	GTCAATCTGC	TCCCTTACTT	TGTCATCACC	AGTCTGGGAA	TAAGGGTATG	180
CCCTATTAC	ACCTGTTAAA	AGGGCGCTAA	GCATTTTTGA	TTCAACATCT	TTTTTTTTGA	240
CACAAGTCCG	AAAAAAGCAA	AAGTAAACAG	TTNTTAATTT	GTTAGCCAAT	TCACTTTCTT	300
CATGGGAGAG	AGCCATTTGA	TTTAAAAAGC	AAATTGCATA	ATATTGAGCT	TTGGGAGCTG	360
ATATNTGAGC	GGAAGANTAG	CCCTTCTACT	TCACCAGACA	CAACTCCTTT	CATATTGGGA	420
TGTTNACNAA	AGTTATGTCT	CTTACAGATG	GGATGCTTTT	GTGGCAATTC	TG	472

(2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 413 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

AGAAACGAGT	ATCTCTNAAA	ACAACCTCTC	ATACCTTGTS	GACCTAATTT	TCTCTGCCCTC	60
TGIGTGTTCG	CGCATATTAT	ATAGACAGGC	ACATCTTTTT	TACTTTTSTA	AAAGCTTATG	120
CCCTTTTGGT	ATCTATATCT	GTGAAAGTTT	TAATGATCTG	CCATAATGTC	TTGGGGACCT	180
TTCTCTTCTG	TGTAAATGGT	ACTAGAGAAA	ACACCTATNT	TATGAGTCAA	TCTAGTTNGT	240
TTTATTCGAC	ATGAAGGAAA	TTTCAGATN	ACAACACTNA	CAAACTCTCC	CTTGACTAGG	300
GGGGACAAAG	AAAAGCANAA	GTGAACATNA	GAAACAATTN	CCTGGTGAGA	AATTNCATAA	360

ACAGAAATTG GGTNGTATAT TGAAANANNG CATCATTNAA ACGTTTTTTT TTT

413

(2) INFORMATION FOR SEQ ID NO:88:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 448 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

CGCAGCGGGT	CCTCTCTATC	TAGCTCCAGC	CTCTCGCCTG	CCCCACTCCC	CGCGTCCCGC	60
GTCCTAGCCN	ACCATGGCCG	GGCCCCGTGG	CGCCCCGCTG	CTCCTGCTGG	CCATCCTGGC	120
CGTGGCCCCG	GCCGTGAGCC	CCGCGGCCGG	CTCCAGTCCC	GGCAAGCCGC	CGCGCCTGCT	180
GGGAGGCCCA	TGGACCCCGC	GTGGAAGAAG	AAGGTGTGCG	GCGTGCACTG	GACTTTGCCG	240
TCGGCNANTA	CAACAAACCC	GCAACNACTT	TTACCNAGCN	CGCGCTGCAG	GTTGTGCCGC	300
CCCAANCAAA	TTGTTACTNG	GGGTAANTAA	TTCTTGGAAG	TTGAACCTGG	GCCAAACNNG	360
TTTACCAGAA	CCNAGCCAAT	TNGAACAAAT	NCCCTCCAT	AACAGCCCCC	TTTAAAAAGG	420
GAANCANTCC	TGNTCTTTTC	CAAATTTT				448

(2) INFORMATION FOR SEQ ID NO:89:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 463 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

GAATTTTGTG	CACTGGCCAC	TGTGATGGAA	CCATTGGGCC	AGGATGCTTT	GAGTTTATCA	60
GTAGTGATTC	TGCCAAAGTT	GGTGTGTGTA	CATGAGTATG	TAAAATGTCA	AAAAATTAGC	120
AGAGGTCTAG	GTCTGCATAT	CAGCAGACAG	TTTGTCCGTG	TATTTTGTAG	CCTTGAAGTT	180
CTCAGTGACA	AGTTNNTTCT	GATGCCAAGT	TCTNATCCCA	GTCTTTTACT	CCTTTGCATC	240
TTTNATGTTN	AGACTTGCCT	CTNTNAAATT	GCTTTTGTNT	TCTGCAGGTA	CTATCTGTGG	300
TTTAACAAAA	TAGAANNACT	TCTCTGCTTN	GAANATTTGA	ATATCTTACA	TCTNAAAATN	360
AATTCTCTCC	CCATANNAAA	ACCCANGCCC	TTGGGANAAT	TTGAAAAANG	GNTCCTTCNN	420
AATTCNNANA	ANTTCAGNTN	TCATACAACA	NAACNGGANC	CCC		463

(2) INFORMATION FOR SEQ ID NO:90:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 400 base pairs
- (B) TYPE: nucleic acid

74

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

AGGGATTGAA	GGTCTNTTNT	ACTGTCCGAC	TGTTCAACCA	CCAACTCTAC	AAGTTGCTGT	60
CTTCCACTCA	CTGTCTGTAA	GCNTNTTAAC	CCAGACTGTA	TCTTCATAAA	TAGAACAAAT	120
TCTTCACCA	TCACATCTTC	TAGGACCTTT	TTGGATTAC	TTACTATAAG	CTCTTCCACT	180
TCCTTTGTTA	AGACTTCATC	TGGTAAAGTC	TTAAGTTTGT	TAGAAAGGAA	TTTAATTGCT	240
CGTTCTCTAA	CAATGTCCCTC	TCCTTGAAGT	ATTGTGGCTGA	ACAACCCACC	TNAAGTCCCT	300
TTGTGCATCC	ATTTTAAATA	TACTTAATAG	GGCATTGGTN	CAC TAGGTTA	AATTCTGCAA	360
GAGTCATCTG	TCTGCAAAAG	TTGCGTTAGT	ATATCTGCCA			400

(2) INFORMATION FOR SEQ ID NO:91:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 480 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

GAGCTCGGAT	CCAATAATCT	TTGTCTGAGG	GCAGCACACA	TATNCAGTGC	CATGGNAACT	60
GGTCTACCCC	ACATGGGAGC	AGCATGCCGT	AGNTATATAA	GGTCATTCCC	TGAGTCAGAC	120
ATGCCTCTTT	GACTACCGTG	TGCCAGTGCT	GGTGATTCTC	ACACACCTCC	NNCCGCTCTT	180
TGTGGAAAAA	CTGGCACTTG	NCTGGAAC TA	GCAAGACATC	ACTTACAAAT	TCACCCACGA	240
GACACTTGAA	AGGTGTAACA	AAGCGACTCT	TGCATTGCTT	TTTGTCCCTC	CGGCACCACT	300
TGTCAATACT	AACCCGCTGG	TTTGCCTCCA	TCACATTGTG	GATCTGTAGC	TCTGGATACA	360
TCTCCTGACA	GTA CTGAAGA	ACTTCTTCTT	TTGTTTCAAA	AGCAACTCTT	GGTGCCTGTT	420
NGATCAGGTT	CCCATTTCCC	AGTCCGAATG	TTCACATGGC	ATATNTTACT	TCCCACAAAA	480

(2) INFORMATION FOR SEQ ID NO:92:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 477 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

ATACAGCCCA	NATCCCACCA	CGAAGATGCC	CTTCTTGA	CTACTGAGT	GAGAACCTGA	TGCGGTCACT	60
GGTCCCGCTG	TAGCCCCAGC	GACTCTCCAC	CTGCTGGAAG	CGGTTGATGC	TGCACTCCTT		120
CCCACGCAGG	CAGCAGCGGG	CCCGGTCAAT	GAAGTCCACT	CGTGGCTTGG	GGTTGACGGT		180
TAANTGCAGG	AAGAGGCTGA	CCACCTCGCG	GTCCACCAGG	ATGCCCCGACT	GTGCGGGGACC		240
TGCAGCGAAA	CTCCTCGATG	CTCATGAGCG	CGAAGCGAAT	GAGCCCCAGG	GCCTTGCCCA		300
GAACCTTCCG	CCTGTTCTCT	GGCGTCACCT	GCAGCTGCTG	CCGCTNACAC	TGCGCCTCGG		360
ACCAGCGGAC	AAACGGCGTT	GAACAGCCGC	ACCTCACGGA	TGCCCCANTGT	GTGCGGCTCC		420
AGGAACGGCN	CCAGCGTGTC	CAGGTCAATG	TGGTGAANC	CTCCGCGGGT	AATGGCG		477

(2) INFORMATION FOR SEQ ID NO:93:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 377 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

GAACGGCTGG	ACCTTGCTC	GCATTGTGCT	GCTGGCAGGA	ATACCTTGGC	AAGCAGCTCC	60
AGTCCGAGCA	GCCCCAGACC	GCTGCCGCCC	GAAGCTAAGC	CTGCCTCTGG	CCTTCCCCCTC	120
CGCCTCAATG	CAGAACCANT	AGTGGGAGCA	CTGTGTTTAG	AGTTAAGAGT	GAACACTGTN	180
TGATTTTACT	TGGGAATTTT	CTCTGTTATA	TAGCTTTTCC	CAATGCTAAT	TTCCAAACAA	240
CAACAACAAA	ATAACATGTT	TGCCTGTTNA	GTTGTATAAA	AGTANGTGAT	TCTGTATNTA	300
AAGAAAATAT	TACTGTTACA	TATACTGCTT	GCAANTTCTG	TATTTATTGG	TNCTCTGGAA	360
ATAAATATAT	TATTAAA					377

(2) INFORMATION FOR SEQ ID NO:94:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 495 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

CCCTTTGAGG	GGTTAGGGTC	CAGTTCCTCAG	TGGAAGAAAC	AGGCCAGGAG	AANTGCGTGC	60
CGAGCTGANG	CAGATTTCCC	ACAGTGACCC	CAGAGCCCTG	CGCTATAGTC	TCTGACCCCT	120
CCAAGGAAAG	ACCACCTTCT	GGGGACATGG	GCTGGAGGGC	AGGACCTAGA	GGCACCAAGG	180
GAAGGCCCCA	TTCCGGGGCT	GTTCCCCGAG	GAGGAAGGGA	AGGGGCTCTG	TGTGCCCCCT	240
AAGAGGAANA	GGCCCTGANT	CCTGGGATCA	NACACCCCTT	CAGTGTATC	CCCACACAAA	300

TGCAAGCTCA	CCAAGGTCCC	CTCTCAGTCC	CTTCCTTACA	CCCTGAACGG	NCACTGGCCC	360
ACACCCACCC	AGANCANCCA	CCCGCCATGG	GGAATGTNCT	CAAGGAATCG	CNGGGCAACG	420
TGGACTCTNG	TCCCNNAAGG	GGGCAGAATC	TCCAATAGAN	GGANNGAACC	CTTGCTNANA	480
AAAAAAAAANA	AAAAA					495

(2) INFORMATION FOR SEQ ID NO:95:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 472 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

GGTTACTTGG	TTTCATTGCC	ACCACTTAGT	GGATGTCATT	TAGAACCATT	TTGTCTGCTC	60
CCTCTGGAAG	CCTTGCGCAG	AGCGGACTTT	GTAATTGTTG	GAGAATAACT	GCTGAATTTT	120
TAGCTGTTTT	GAGTTGATTC	GCACCACTGC	ACCACAACCTC	AATATGAAAA	CTATTTNACT	180
TATTTATTAT	CTTGTGAAAA	GTATACAATG	AAAATTTTGT	TCATACTGTA	TTTATCAAGT	240
ATGATGAAAA	GCAATAGATA	TATATTCTTT	TATTATGTTN	AATTATGATT	GCCATTATTA	300
ATCGGCAAAA	TGTGGAGTGT	ATGTTCTTTT	CACAGTAATA	TATGCCTTTT	GTAACCTTAC	360
TTGGTTATTT	TATTGTAAAT	GAATTACAAA	ATTCTTAATT	TAAGAAAATG	GTANGTTATA	420
TTTANTTCAN	TAATTTCTTT	CCTTGTTTAC	GTTAATTTTG	AAAAGAATGC	AT	472

(2) INFORMATION FOR SEQ ID NO:96:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 476 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

CTGAAGCATT	TCTTCAAAC	TNTCTACTTT	TGTCATTGAT	ACCTGTAGTA	AGTTGACAAT	60
GTGGTGAAAT	TTCAAAATTA	TATGTAACCT	CTACTAGTTT	TACTTTCTCC	CCCAAGTCTT	120
TTTTAACTCA	TGATTTTAC	ACACACAATC	CAGAACTTAT	TATATAGCCT	CTAAGTCTTT	180
ATTCTTCACA	GTAGATGATG	AAAGAGTCCT	CCAGTGTCTT	GNGCANAATG	TTCTAGNTAT	240
AGCTGGATAC	ATACNGTGGG	AGTTCATAAA	ACTCATACCT	CAGTGGGACT	NAACCAAAAT	300
TGTGTTAGTC	TCAATTCCCTA	CCACACTGAG	GGAGCCTCCC	AAATCACTAT	ATTCTTATCT	360
GCAGGTACTC	CTCCAGAAAA	ACNGACAGGG	CAGGCTTGCA	TGAAAAAGTN	ACATCTGCGT	420
TACAAAGTCT	ATCTTCCTCA	NANGTCTGTN	AAGGAACAAT	TTAATCTTCT	AGCTTT	476

(2) INFORMATION FOR SEQ ID NO:97:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 479 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

ACTCTTTCTA ATGCTGATAT GATCTTGAGT ATAAGAATGC ATATGTCACT AGAATGGATA	60
AAATAATGCT GCAAACCTAA TGTTCCTTATG CAAAATGGAA CGCTAATGAA ACACAGCTTA	120
CAATCGCAAA TCAAAACTCA CAAGTGCTCA TCTGTTGTAG ATTTAGTGTA ATAAGACTTA	180
GATTGTGCTC CTTCGGATAT GATTGTTTCT CANATCTTGG GCAATNTTCC TTAGTCAAAT	240
CAGGCTACTA GAATTCTGTT ATTGGATATN TGAGAGCATG AAATTTTTAA NAATACACTT	300
GTGATTATNA AATTAATCAC AAATTTCACT TATACCTGCT ATCAGCAGCT AGAAAAACAT	360
NTNNTTTTTA NATCAAAGTA TTTTGTGTTT GGAANTGTNN AAATGAAATC TGAATGTGGG	420
TTCNATCTTA TTTTTTCCCN GACNACTANT TNCTTTTTTA GGGNCTATTC TGANCCATC	479

(2) INFORMATION FOR SEQ ID NO:98:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 461 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

AGTGACTTGT CCTCCAACAA AACCCTTGA TCAAGTTTGT GGCCTGACA ATCAGACCTA	60
TGCTAGTTCC TGTCATCTAT TCGCTACTAA ATGCAGACTG GAGGGGACCA AAAAGGGGCA	120
TCAACTCCAG CTGGATTATT TTGGAGCCTG CAAATCTATT CCTACTTGTA CGGACTTTGA	180
AGTGATTCAG TTTCTCTAC GGATGAGAGA CTGGCTCAAG AATATCCTCA TGCAGCTTTA	240
TGAAGCCACT CTGAACACGC TGGTTATCTA GATGAGAACA GAGAAATAAA GTCAGAAAAT	300
TTACCTGGAG AAAAGAGGCT TTGGCTGGGG ACCATCCCAT TGAACCTTCT CTTAAGCACT	360
TTAAGAAAAA CTACCACATG TTGTGTATCC TGGTGCCGGC CGTTTATGAA CTGACCACCC	420
TTTGGAATAA TCTTGACGCT CCTGAACCTG CTCCTCTGCG A	461

(2) INFORMATION FOR SEQ ID NO:99:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 171 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

78

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

GTGGCCGCGC GCAGTGTGTTT CCTGTACCG CAGGGCCCCC TCCCTTCCCC AGGCGTCCCT	60
CGGCGCCTCT GCGGCCCCGA GGAGGAGCGG CTGCGGGGTG GGGGGAGTGT GACCCACCT	120
CGGTGAGAAA AGCCTTCTCT AGCGATCTGA GAGCGGTGCC TTGGGGGTAC C	171

(2) INFORMATION FOR SEQ ID NO:100:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 269 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

CGGCCGCAAG TGCAACTCCA GCTGGGGCCC TGCGGACGAA GATTCTGCCA GCAGTTGGTC	60
CGACTGCGAC GACGGCGGCG GCGACAGTCG CAGGTGCAGC GCGGGCGCCT GGGGTCTTGC	120
AAGGCTGAGC TGACCGCGCA GAGGTGCTGT CACGTCCGAC GACCTTGACG CCGTCGGGGA	180
CAGCCGGAAC AGAGCCCGGT GAAGCGGGAG GCCTCGGGGA GCCCCTCGGG AAGGGCGGCC	240
CGAGAGATAC GCAGGTGCAG GTGGCCGCC	269

(2) INFORMATION FOR SEQ ID NO:101:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 405 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

TTTTTTTTTT TTTTGAATC TACTGCCAGC ACAGCAGGTC AGCAACAAGT TTATTTTGCA	60
GCTAGCAAGG TAACAGGGTA GGGCATGGTT ACAATGTCAG GTCAACTTCC TTTGTCTGTG	120
TTGATTGGTT TGTCTTTATG GGGGCGGGGT GGGGTAGGGG AAACGAAGCA AATAACATG	180
AGTGGGTGCA CCTCCCTGT AGAACCTGGT TACAAAGCTT GGGGCAGTTC ACCTGGTCTG	240
TGACCGTCAT TTTCTTGACA TCAATGTTAT TAGAAGTCAG GATATCTTTT AGAGAGTCCA	300
CTGTTCTGGA GGGAGATTAG GGTTCCTTGG CAAATCCAAC AAAATCCACT GAAAAAGTTG	360
GATGATCAGT ACGAATACCG AGGCATATTC TCATATCGGT GGCCA	405

(2) INFORMATION FOR SEQ ID NO:102:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 470 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	60
GGCACHTAAT	CCATTTTAT	TTCAAAATGT	CTACAAATTT	AATCCCATTA	TACGGTATTT	120
TCAAAATCTA	AATTATTCAA	ATTAGCCAAA	TCCTTACCAA	ATAATACCCA	AAAATCAAAA	180
ATATACTTCT	TTCAGCAAAC	TTGTTACATA	AATTAAGGAA	ATATATACGG	CTGGTGTTTT	240
CAAAGTACAA	TTATCTTAAC	ACTGCAAACA	TTTTAAGGAA	CTAAAATAAA	AAAAAACACT	300
CCGCAAAGGT	TAAAGGGAAC	AACAAATTCT	TTTACAACAC	CATTATAAAA	ATCATATCTC	360
AAATCTTAGG	GGAATATATA	CTTCACACGG	GATCTTAACT	TTTACTCACT	TTGTTTATTT	420
TTTTAAACCA	TTGTTTGGGC	CCAACACAAT	GGAATCCCCC	CTGGACTAGT		470

(2) INFORMATION FOR SEQ ID NO:103:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 581 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:103:

TTTTTTTTTT	TTTTTTTTTT	CCCCCTCTT	ATAAAAAACA	AGTTACCATT	TTATTTTACT	60
TACACATATT	TATTTTATAA	TTGGTATTAG	ATATTCAAAA	GGCAGCTTTT	AAAATCAAAC	120
TAAATGGAAA	CTGCCTTAGA	TACATAATTC	TTAGGAATTA	GCTTAAAATC	TGCCTAAAGT	180
GAAAATCTTC	TCTAGCTCTT	TTGACTGTAA	ATTTTTGACT	CTTGTAAGAC	ATCCAAATTC	240
ATTTTTCTTG	TCTTTAAAAT	TATCTAATCT	TTCCATTTTT	TCCCTATTCC	AAGTCAATTT	300
GCTTCTCTAG	CCTCATTTCC	TAGCTCTTAT	CTACTATTAG	TAAGTGGCTT	TTTTCTCTAA	360
AGGGAAAACA	GGAAGAGAAA	TGGCACACAA	AACAAACATT	TTATATTCAT	ATTTCTACCT	420
ACGTTAATAA	AATAGCATTT	TGTGAAGCCA	GCTCAAAAGA	AGGCTTAGAT	CCTTTTATGT	480
CCATTTTAGT	CACTAAACGA	TATCAAAGTG	CCAGAATGCA	AAAGCTTTCT	GAACATTTAT	540
TCAAAAGCTA	ATATAAGATA	TTTCACATAC	TCATCTTTCT	G		581

(2) INFORMATION FOR SEQ ID NO:104:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 578 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

80

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:104:

TTTTTTTTTT	TTTTTTTTTT	TTTTCTCTT	CTTTTTTTTT	GAAATGAGGA	TCGAGTTTTT	60
CACTCTCTAG	ATAGGGCATG	AAGAAAACCT	ATCTTTCCAG	CTTTAAAATA	ACAATCAAAT	120
CTCTTAIGCT	ATATCATATT	TTAAGTTAAA	CTAATGAGTC	ACTGGCTTAT	CTTCTCCTGA	180
AGGAAATCTG	TTCATTCTTC	TCATTTCATAT	AGTTATATCA	AGTACTACCT	TGCATATTGA	240
GAGGTTTTTC	TTCTCTATTT	ACACATATAT	TTCCATGTGA	ATTTGTATCA	AACCTTTATT	300
TTCATGCAAA	CTAGAAAATA	ATCTTTCTTT	TGCATAAGAG	AAGAGAACAA	TATAGCATTAT	360
CAAACTGCT	CAAATTGTTT	GTAAAGTTAT	CCATTATAAT	TAGTTGGCAG	GAGCTAATAC	420
AAATCACATT	TACGACAGCA	ATAATAAAAC	TGAAGTACCA	GTTAAATATC	CAAAATAATT	480
AAAGGAACAT	TTTTAGCCTG	GGTATAATTA	GCTAATTCAC	TTTACAAGCA	TTTATTAGAA	540
TGAATTCACA	TGTTATTATT	CCTAGCCCAA	CACAATGG			578

(2) INFORMATION FOR SEQ ID NO:105:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 538 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:105:

TTTTTTTTTT	TTTTTCAGTA	ATAATCAGAA	CAATATTTAT	TTTTATATTT	AAAATTCATA	60
GAAAAGTGCC	TTACATTTAA	TAAAAGTTTG	TTTCTCAAAG	TGATCAGAGG	AATTAGATAT	120
GTCTTGAACA	CCAATATTAA	TTTGAGGAAA	ATACACCAAA	ATACATTAAG	TAAATTATTT	180
AAGATCATAG	AGCTTGTAAG	TGAAAAGATA	AAATTTGACC	TCAGAAACTC	TGAGCATTAA	240
AAATCCACTA	TTAGCAAATA	AATTACTATG	GACTTCTTGC	TTTAATTTTG	TGATGAATAT	300
GGGGTGTCAC	TGGTAAACCA	ACACATTCTG	AAGGATACAT	TACTTAGTGA	TAGATTCTTA	360
TGTACTTTGC	TAATACGTGG	ATATGAGTTG	ACAAGTTTCT	CTTTCTTCAA	TCTTTTAAGG	420
GGCGAGAAAT	GAGGAAGAAA	AGAAAAGGAT	TACGCATACT	GTTCTTTCTA	TGGAAGGATT	480
AGATATGTTT	CCTTTGCCAA	TATTAAAAAA	ATAATAATGT	TTACTACTAG	TGAAACCC	538

(2) INFORMATION FOR SEQ ID NO:106:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 473 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Homo sapiens*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:106:

TTTTTTTTTT	TTTTTTAGTC	AAGTTTCTAT	TTTTATTATA	ATTAAAGTCT	TGGTCATTTC	60
ATTTATTAGC	TCTGCAACTT	ACATATTTAA	ATTAAAGAAA	CGTTTTAGAC	AACTGTACAA	120
TTTATAAATG	TAAGGTGCCA	TTATTGAGTA	ATATTTCCT	CCAAGAGTGG	ATGTGTCCCT	180
TCTCCCACCA	ACTAATGAAC	AGCAACATTA	GTTTAATTTT	ATTAGTAGAT	ATACACTGCT	240
GCAAACGCTA	ATTCTCTTCT	CCATCCCCAT	GTGATATTGT	GTATATGTGT	GAGTTGGTAG	300
AATGCATCAC	AATCTACAAT	CAACAGCAAC	ATGAAGCTAG	GCTGGGCTTT	CGGTGAAAAT	360
AGACTGTGTC	TGTCTGAATC	AAATGATCTG	ACCTATCCTC	GGTGGCAAGA	ACTCTTCGAA	420
CCGCTTCCTC	AAAGGCGCTG	CCACATTTGT	GGCTCTTTGC	ACTTGTTC	AAA	473

(2) INFORMATION FOR SEQ ID NO:107:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1621 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Homo sapiens*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:107:

CGCCATGGCA	CTGCAGGGCA	TCTCGGTCAT	GGAGCTGTCC	GGCCTGGCCC	CGGGCCCGTT	60
CTGTGCTATG	GTCCTGGCTG	ACTTCGGGGC	GCGTGTGGTA	CGCGTGGACC	GGCCCGGCTC	120
CCGCTACGAC	GTGAGCCGCT	TGGGCCGGGG	CAAGCGCTCG	CTAGTGCTGG	ACCTGAAGCA	180
GCCGCGGGGA	GCCGCCGTGC	TGCGGCGTCT	GTGCAAGCGG	TGGATGTGC	TGCTGGAGCC	240
CTTCCGCCGC	GGTGTCTATG	AGAAACTCCA	GCTGGGCCCC	GAGATTCTGC	AGCGGGAAAA	300
TCCAAGGCTT	ATTTATGCCA	GGCTGAGTGG	ATTTGGCCAG	TCAGGAAGCT	TCTGCCGGTT	360
AGCTGGCCAC	GATATCAACT	ATTTGGCTTT	GTCAGGTGTT	CTCTCAAAAA	TTGGCAGAAG	420
TGGTGAGAAT	CCGTATGCCC	CGCTGAATCT	CCTGGCTGAC	TTTGCTGGTG	GTGGCCTTAT	480
GTGTGCACTG	GGCATTATAA	TGGCTCTTTT	TGACCGCACA	CGCACTGACA	AGGGTCAGGT	540
CATTGATGCA	AATATGGTGG	AAGGAACAGC	ATATTTAAGT	TCTTTTCTGT	GGAAAACTCA	600
GAAATCGAGT	CTGTGGGAAG	CACCTCGAGG	ACAGAACATG	TTGGATGGTG	GAGCACCTTT	660
CTATACGACT	TACAGGACAG	CAGATGGGGA	ATTCATGGCT	GTTGGAGCAA	TAGAACCCCC	720
GTTCTACGAG	CTGCTGATCA	AAGGACTTGG	ACTAAAGTCT	GATGAACTTC	CCAATCAGAT	780
GAGCATGGAT	GATTGGCCAG	AAATGAAGAA	GAAGTTTGCA	GATGTATTTG	CAAAGAAGAC	840
GAAGGCAGAG	TGGTGTCAAA	TCTTTGACGG	CACAGATGCC	TGTGTGACTC	CGGTTCTCAC	900
TTTTGAGGAG	GTTGTTTCATC	ATGATCACAA	CAAGGAACGG	GGCTCGTTTA	TCACCAGTGA	960
GGAGCAGGAC	GTGAGCCCCC	GCCCTGCACC	TCTGCTGTTA	AACACCCCA	CCATCCCTTC	1020
TTTCAAAAGG	GATCCTTTCA	TAGGAGAACA	CACTGAGGAG	ATACTTGAAG	AATTTGGATT	1080
CAGCCGCGAA	GAGATTTATC	AGCTTAACTC	AGATAAAATC	ATTGAAAGTA	ATAAGGTAAA	1140
AGCTAGTCTC	TAACCTCCAG	GCCCACGGCT	CAAGTGAATT	TGAATACTGC	ATTTACAGTG	1200
TAGAGTAACA	CATAACATTG	TATGCATGGA	AAATGAGGAG	AACAGTATTA	CAGTGTCTTA	1260
CCACTCTAAT	CAAGAAAAGA	ATTACAGACT	CTGATTCTAC	AGTGATGATT	GAATTCTAAA	1320
AATGGTTATC	ATTAGGGCTT	TGATTTTATA	AAACTTTGGG	TACTTATACT	AAATTATGST	1380
AGTTATTCTG	CCTTCCAGTT	TGCTTGATAT	ATTTGTTGAT	ATTAAGATTC	TTGACTTATA	1440
TTTTGAATGG	GTTCTAGTGA	AAAAGGAATG	ATATATTCTT	GAAGACATCG	ATATACATTT	1500
ATTTACACTC	TTGATTCTAC	AATGTAGAAA	ATGAGGAAAT	GCCACAAATT	GTATGGTGAT	1560

AAAAAGTCACG TGAACACAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 1620
A 1621

(i) INFORMATION FOR SEQ ID NO:108:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 382 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:108:

Met	Ala	Leu	Gln	Gly	Ile	Ser	Val	Met	Glu	Leu	Ser	Gly	Leu	Ala	Pro	1	5	10	15
Gly	Pro	Phe	Cys	Ala	Met	Val	Leu	Ala	Asp	Phe	Gly	Ala	Arg	Val	Val	20	25	30	
Arg	Val	Asp	Arg	Pro	Gly	Ser	Arg	Tyr	Asp	Val	Ser	Arg	Leu	Gly	Arg	35	40	45	
Gly	Lys	Arg	Ser	Leu	Val	Leu	Asp	Leu	Lys	Gln	Pro	Arg	Gly	Ala	Ala	50	55	60	
Val	Leu	Arg	Arg	Leu	Cys	Lys	Arg	Ser	Asp	Val	Leu	Leu	Glu	Pro	Phe	65	70	75	80
Arg	Arg	Gly	Val	Met	Glu	Lys	Leu	Gln	Leu	Gly	Pro	Glu	Ile	Leu	Gln	85	90	95	
Arg	Glu	Asn	Pro	Arg	Leu	Ile	Tyr	Ala	Arg	Leu	Ser	Gly	Phe	Gly	Gln	100	105	110	
Ser	Gly	Ser	Phe	Cys	Arg	Leu	Ala	Gly	His	Asp	Ile	Asn	Tyr	Leu	Ala	115	120	125	
Leu	Ser	Gly	Val	Leu	Ser	Lys	Ile	Gly	Arg	Ser	Gly	Glu	Asn	Pro	Tyr	130	135	140	
Ala	Pro	Leu	Asn	Leu	Leu	Ala	Asp	Phe	Ala	Gly	Gly	Gly	Leu	Met	Cys	145	150	155	160
Ala	Leu	Gly	Ile	Ile	Met	Ala	Leu	Phe	Asp	Arg	Thr	Arg	Thr	Asp	Lys	165	170	175	
Gly	Gln	Val	Ile	Asp	Ala	Asn	Met	Val	Glu	Gly	Thr	Ala	Tyr	Leu	Ser	180	185	190	
Ser	Phe	Leu	Trp	Lys	Thr	Gln	Lys	Ser	Ser	Leu	Trp	Glu	Ala	Pro	Arg	195	200	205	
Gly	Gln	Asn	Met	Leu	Asp	Gly	Gly	Ala	Pro	Phe	Tyr	Thr	Thr	Tyr	Arg	210	215	220	
Thr	Ala	Asp	Gly	Glu	Phe	Met	Ala	Val	Gly	Ala	Ile	Glu	Pro	Gln	Phe	225	230	235	240
Tyr	Glu	Leu	Leu	Ile	Lys	Gly	Leu	Gly	Leu	Lys	Ser	Asp	Glu	Leu	Pro	245	250	255	
Asn	Gln	Met	Ser	Met	Asp	Asp	Trp	Pro	Glu	Met	Lys	Lys	Lys	Phe	Ala	260	265	270	
Asp	Val	Phe	Ala	Lys	Lys	Thr	Lys	Ala	Glu	Trp	Cys	Gln	Ile	Phe	Asp	275	280	285	
Gly	Thr	Asp	Ala	Cys	Val	Thr	Pro	Val	Leu	Thr	Phe	Glu	Glu	Val	Val				

290		295		300
His His Asp His Asn Lys Glu Arg Gly Ser Phe Ile Thr Ser Glu Glu				
305		310		315
Gln Asp Val Ser Pro Arg Pro Ala Pro Leu Leu Leu Asn Thr Pro Ala				320
		325		330
Ile Pro Ser Phe Lys Arg Asp Pro Phe Ile Gly Glu His Thr Glu Glu				335
		340		345
Ile Leu Glu Glu Phe Gly Phe Ser Arg Glu Glu Ile Tyr Gln Leu Asn				350
		355		360
Ser Asp Lys Ile Ile Glu Ser Asn Lys Val Lys Ala Ser Leu				365
370		375		380

(2) INFORMATION FOR SEQ ID NO:109:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1524 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:109:

GGCACCAGGC	TGCGCCAGGG	CCTGAGCGGA	GGCGGGGGCA	GCCTCGCCAG	CGGGGGCCCC	60
GGGCCTGGCC	ATGCCTCACT	GAGCCAGCGC	CTGCGCTCT	ACCTCGCCGA	CAGCTGGAAC	120
CAGTGCGACC	TAGTGGCTCT	CACCTGCTTC	CTCCTGGGCG	TGGGCTGCCG	GCTGACCCCC	180
GGTTTGTACC	ACCTGGGCGG	CACGTCTCTC	TGCATCGACT	TCATGGTTTT	CACGGTGCGG	240
CTGCTTCACA	TCTTCACGGT	CAACAAACAG	CTGGGGCCCCA	AGATCGTCAT	CGTGAGCAAG	300
ATGATGAAGG	ACGTGTTCTT	CTTCCTCTTC	TTCTCTGGCG	TGTGGCTGGT	AGCCTATGGC	360
GTGGCCACGG	AGGGGCTCCT	GAGGCCACGG	GACAGTGACT	TCCCAAGTAT	CCTGCGCCGC	420
GTCTTCTACC	GTCCCTACCT	GCAGATCTTC	GGGCAGATTC	CCCAGGAGGA	CATGGACGTG	480
GCCCTCATGG	AGCACAGCAA	CTGCTCGTCG	GAGCCCGGCT	TCTGGGCACA	CCCTCCTGGG	540
GCCCAGGCGG	GCACCTGCGT	CTCCAGTAT	GCCAACTGGC	TGGTGGTGCT	GCTCCTCGTC	600
ATCTTCCTGC	TGCTGGCCAA	CATCCTGCTG	GTCAACTTGC	TCATTGCCAT	GTTCAAGTAC	660
ACATTGGGCA	AAGTACAGGG	CAACAGCGAT	CTCTACTGGA	AGGCGCAGCG	TTACCGCCTC	720
ATCCGGGAAT	TCCACTCTCG	GCCCCGCGTG	GCCCCGCGCT	TTATCGTCAT	CTCCCACTTG	780
CGCTCCTGC	TCAGGCAATT	GTGCAGGCGA	CCCCGGAGCC	CCCAGCCGTC	CTCCCCGGCC	840
CTCCAGCATT	TCCGGGTTTA	CCTTTCTAAG	GAAGCCGAGC	GGAAGCTGCT	AACGTGGGAA	900
TCCGTGCATA	AGGAGAACTT	TCTGCTGGCA	CGCGCTAGGG	ACAAGCGGGA	GAGCGACTCC	960
GAGCGTCTGA	AGCGCACGTC	CCAGAAGGTG	GACTTGGCAC	TGAAACAGCT	GGGACACATC	1020
CGCGAGTACG	AACAGCGCCT	GAAAGTGCTG	GAGCGGGAGG	TCCAGCAGTG	TACCCGCGTC	1080
CTGGGGTGGG	TGGCCGAGGC	CTTGAGCCGC	TCTGCTTCTG	TGCCCCCAGG	TGGGCGGCCA	1140
CCCCCTGACC	TGCCTGGGTC	CAAAGACTGA	GCCCTGCTGG	CGGACTTCAA	GGAGAAAGCC	1200
CCACAGGGGA	TTTTGCTCCT	AGAGTAAGGC	TCATCTGGGC	CTCGGCCCCC	GCACCTGGTG	1260
GCCTTGTCCT	TGAGGTGAGC	CCCATGTCCA	TCTGGGCCAC	TGTCAGGACC	ACCTTTGGGA	1320
GTGTCATCCT	TACAAACCAC	AGCATGCCCC	GCTCTCTCCA	GAACCAGTCC	CAGCCTGGGA	1380
GGATCAAGGC	CTGGATCCCC	GGCCGTTATC	CATCTGGAGG	CTGCAGGGTC	CTTGGGGTAA	1440
CAGGGACCAC	AGACCCCTCA	CCACTCACAG	ATTCTCACA	CTGGGGAAAT	AAAGCCATTT	1500
CAGAGGAAAA	AAAAAAAAAA	AAAA				1524

(2) INFORMATION FOR SEQ ID NO:110:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3410 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:110:

GGGAACCAGC	CTGCACGCCG	TGGCTCCGGG	TGACAGCCGC	GCGCCTCGGC	CAGGATCTGA	60
GTGATGAGAC	GTGTCCCCAC	TGAGGTGCCC	CACAGCAGCA	GGTGTGAGC	ATGGGCTGAG	120
AAGCTGGACC	GGCACCAAAG	GGCTGGCAGA	AATGGGCGCC	TGGCTGATTG	CTAGGCAGTT	180
GGCGGCAGCA	AGGAGGAGAG	GGCGCAGCTT	CTGGAGCAGA	GGCGAGACGA	AGCAGTTCTG	240
GAGTGCCTGA	ACGGCCCCCT	GAGCCCTACC	CGCCTGGCCC	ACTATGGTCC	AGAGGCTGTG	300
GGTGAGCCGC	CTGCTGCGGC	ACCGGAAAGC	CCAGCTCTTG	CTGGTCAACC	TGCTAACCTT	360
TGGCCTGGAG	GTGTGTTTGG	CCGCAGGCAT	CACCTATGTG	CCGCCTCTGC	TGCTGGAAGT	420
GGGGCTAGAG	GAGAAGTTCA	TGACCATGGT	GCTGGGCATT	GGTCCAGTGC	TGGGCCTGGT	480
CTGTGTCCCG	CTCCTAGGCT	CAGCCAGTGA	CCACTGGCGT	GGACGCTATG	GCCGCCGCCG	540
GCCCTTCATC	TGGGCACTGT	CCTTGGGCAT	CCTGCTGAGC	CTCTTTCTCA	TCCCAAGGGC	600
CGGCTGGCTA	GCAGGGCTGC	TGTGCCCGGA	TCCCAGGCCC	CTGGAGCTGG	CACTGCTCAT	660
CCTGGGCGTG	GGGCTGCTGG	ACTTCTGTGG	CCAGGTGTGC	TTCCTCCAC	TGGAGGCCCT	720
GCTCTCTGAC	CTCTTCCGGG	ACCCGGACCA	CTGTGCGCCAG	GCCTACTCTG	TCTATGCCTT	780
CATGATCAGT	CTTGGGGGCT	GCCTGGGCTA	CCTCCTGCCT	GCCATTGACT	GGGACACCAG	840
TGCCCTGGCC	CCCTACCTGG	GCACCCAGGA	GGAGTGCCTC	TTTGGCCTGC	TCACCCTCAT	900
CTTCTCACC	TGCGTAGCAG	CCACACTGCT	GGTGGCTGAG	GAGGCAGCGC	TGGGCCCCAC	960
CGAGCCAGCA	GAAGGGCTGT	CGGCCCCCTC	CTTGTGCGCC	CACTGCTGTC	CATGCCGGGC	1020
CCGCTTGGCT	TTCCGGAACC	TGGGCGCCCT	GCTTCCCCCG	CTGCACCAGC	TGTGCTGCCG	1080
CATGCCCCGC	ACCCTGCGCC	GGCTCTTCGT	GGCTGAGCTG	TGCAGCTGGA	TGGCACTCAT	1140
GACCTTCACG	CTGTTTTACA	CGGATTTCTG	GGGCGAGGGG	CTGTACCAGG	GCGTGCCCAG	1200
AGCTGAGCCG	GGCACCGAGG	CCCGGAGACA	CTATGATGAA	GGCGTTCGGA	TGGGCAGCCT	1260
GGGGCTGTTT	CTGCAGTGCG	CCATCTCCCT	GGTCTTCTCT	CTGGTCATGG	ATGGGCTGGT	1320
GCAGCGATTG	GGCACTCGAG	CAGTCTATTT	GGCCAGTGTG	GCAGCTTTCC	CTGTGGCTGC	1380
CGGTGCCACA	TGCCCTGTCC	ACAGTGTGGC	CGTGGTGACA	GCTTCAGCCG	CCCTCACCCG	1440
GTTACCTTC	TCAGCCCTGC	AGATCCTGCC	CTACACACTG	GCCTCCCTCT	ACCACCGGGA	1500
GAAGCAGGTG	TTCCTGCCCC	AATACCGAGG	GGACACTGGA	GGTGCTAGCA	GTGAGGACAG	1560
CCTGATGACC	AGCTTCTCTG	CAGGCCCTAA	GCCTGGAGCT	CCCTTCCCTA	ATGGACACGT	1620
GGGTGCTGGA	GGCAGTGGCC	TGCTCCACCC	TCCACCCCGG	CTCTGCGGGG	CCTCTGCCCTG	1680
TGATGTCTCC	GTACGTGTGG	TGGTGGGTGA	GCCCACCGAG	GCCAGGGTGG	TTCCGGGGCCG	1740
GGGCATCTGC	CTGGACCTCG	CCATCCTGGA	TAGTGCCTTC	CTGCTGTCCC	AGGTGGCCCC	1800
ATCCCTGTTT	ATGGGCTCCA	TTGTCCAGCT	CAGCCAGTCT	GTCACCTGCCT	ATATGGTGTC	1860
TGCCCGAGGC	CTGGGTCTGG	TCGCCATTTA	CTTTGCTACA	CAGGTAGTAT	TTGACAAGAG	1920
CGACTTGGCC	AAATACTCAG	CGTAGAAAAC	TTCCAGCACA	TTGGGGTGGA	GGGCTGCCTT	1980
CACTGGGTCC	CAGCTCCCCG	CTCCTGTTAG	CCCCATGGGG	CTGCCGGGGT	GGCCGCCAGT	2040
TTCTGTTGCT	GCCAAAGTAA	TGTGGCTCTC	TGCTGCCACC	CTGTGCTGCT	GAGGTGCCGT	2100
GCTGCACAGC	TGGGGGCTGG	GGCGTCCCTC	TCCTCTCTCC	CCAGTCTCTA	GGGCTGCCTG	2160
ACTGGAGGCC	TTCGAAGGGG	GTTTCAAGTCT	GGACTTATAC	AGGGAGGCTA	GAAGGGCTCC	2220
ATGCACTGGA	ATGCGGGGAC	TCTGCAGGTG	GATTACCCAG	GCTCAGGGTT	AACAGCTAGC	2280
CTCCTAGTTG	AGACACACCT	AGAGAAGGGT	TTTTGGGAGC	TGAATAAACT	CAGTCACCTG	2340
GTTTCCCATC	TCTAAGCCCC	TTAACCTGCA	GTTTCCTTTA	ATGTAGCTCT	TGCTATGGAG	2400
TTTCTAGGAT	GAAACACTCC	TCCATGGGAT	TTGAACATAT	GACTTATTTG	TAGGGGAAGA	2460

GTCCTGAGGG	GCAACACACA	AGAACCAGGT	CCCCTCAGCC	CACAGCACTG	TCTTTTTGCT	2520
GATCCACCCC	CCTCTTACCT	TTTATCAGGA	TGTGGCCTGT	TGGTCCTTCT	GTTGCCATCA	2580
CAGAGACACA	GGCATTTAAA	TATTTAACTT	ATTTATTTAA	CAAAGTAGAA	GGGAATCCAT	2640
TGCTAGCTTT	TCTGTGTTGG	TGTCTAATAT	TTGGGTAGGG	TGGGGGATCC	CCAACAATCA	2700
GGTCCCCTGA	GATAGCTGGT	CATTGGGCTG	ATCATTGCCA	GAATCTTCTT	CTCCTGGGGT	2760
CTGGCCCCCC	AAAATGCCTA	ACCCAGGACC	TTGGAAATTC	TACTCATCCC	AAATGATAAT	2820
TCCAAATGCT	GTTACCCAAG	GTTAGGGTGT	TGAAGGAAGG	TAGAGGGTGG	GGCTTCAGGT	2880
CTCAACGGCT	TCCCTAACCA	CCCCTCTTCT	CTTGGCCCCAG	CCTGGTTCCC	CCCCTTCCA	2940
CTCCCCCTCTA	CTCTCTCTAG	GACTGGGCTG	ATGAAGGCAC	TGCCCCAAAT	TTCCCCCTACC	3000
CCCAACTTTC	CCCTACCCCC	AACTTTCCCC	ACCAGCTCCA	CAACCCTGTT	TGGAGCTACT	3060
GCAGGACCAG	AAGCACAAAG	TGCGGTTTCC	CAAGCCTTTG	TCCATCTCAG	CCCCCAGAGT	3120
ATATCTGTGC	TTGGGGAATC	TCACACAGAA	ACTCAGGAGC	ACCCCTGCC	TGAGCTAAGG	3180
GAGGTCTTAT	CTCTCAGGGG	GGGTTTAAGT	GCCGTTTCCA	ATAATGTCGT	CTTATTTATT	3240
TAGCGGGGTG	AATATTTTAT	ACTGTAAGTG	AGCAATCAGA	GTATAATGTT	TATGGTGACA	3300
AAATTAAAGG	CTTTCTTATA	TGTTTAAAAA	AAAAAAAAAA	AAAAAAAAAA	AAAAAAAAAA	3360
AAAAAAAAARA	AAAAAAAAAA	AAAAAAAAAA	AAAAAATAA	AAAAAAAAAA		3410

(2) INFORMATION FOR SEQ ID NO:111:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1289 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:111:

AGCCAGGCGT	CCCTCTGCCT	GCCCACTCAG	TGGCAACACC	CGGGAGCTGT	TTTGTCCCTTT	60
GTGGAGCCTC	AGCAGTTCCC	TCTTTCAGAA	CTCACTGCCA	AGAGCCCTGA	ACAGGAGCCA	120
CCATGCAGTG	CTTCAGCTTC	ATTAAGACCA	TGATGATCCT	CTTCAATTTG	CTCATCTTTC	180
TGTGTGGTGC	AGCCCTGTTG	GCAGTGGGCA	TCTGGGTGTC	AATCGATGGG	GCATCCTTTC	240
TGAAGATCTT	CGGGCCACTG	TGCTCCAGTG	CCATGCAGTT	TGTCAACGTG	GGCTACTTCC	300
TCATCGCAGC	CGGCGTTGTG	GTCTTTGCTC	TTGGTTTCCCT	GGGCTGCTAT	GGTGCTAAGA	360
CTGAGAGCAA	GTGTGCCCTC	GTGACGTTCT	TCTTCATCCT	CCTCCTCATC	TTCATTGCTG	420
AGGTTGCAGC	TGCTGTGGTC	GCCTTGGTGT	ACACCACAAT	GGCTGAGCAC	TTCCTGACGT	480
TGCTGGTAGT	GCCTGCCATC	AAGAAAGATT	ATGGTTCCCA	GGAAGACTTC	ACTCAAGTGT	540
GGAACACCAC	CATGAAAGGG	CTCAAGTGCT	GTGGCTTCAC	CAACTATACG	GATTTTGGAG	600
ACTCACCTTA	CTTCAAAGAG	AACAGTGCCT	TTCCCCCATT	CTGTTGCAAT	GACAACGTCA	660
CCAACACAGC	CAATGAAACC	TGCACCAAGC	AAAAGGCTCA	CGACCAAAAA	GTAGAGGGTT	720
GCTTCAATCA	GCTTTTGTAT	GACATCCGAA	CTAATGCACT	CACCGTGGGT	GGTGTGGCAG	780
CTGGAATTGG	GGGCCTCGAG	CTGGCTGCCA	TGATTGTGTC	CATGTATCTG	TACTGCAATC	840
TACAATAAGT	CCACTTCTGC	CTCTGCCACT	ACTGCTGCCA	CATGGGAACT	GTGAAGAGGG	900
ACCCTGGCAA	GCAGCAGTGA	TTGGGGGAGG	GGACAGGATC	TAACAATGTC	ACTTGGGCCA	960
GAATGGACCT	GCCCTTTCTG	CTCCAGACTT	GGGGCTAGAT	AGGGACCACT	CCTTTTAGCG	1020
ATGCCTGACT	TTCTTTCCAT	TGGTGGGTGG	ATGGGTGGGG	GGCATTCCAG	AGCCTCTAAG	1080
GTAGCCAGTT	CTGTTGCCCA	TTCCCCAGT	CTATTAAACC	CTTGATATGC	CCCCTAGGCT	1140
TAGTGGTGT	CCCAGTGCTC	TACTGGGGGA	TGAGAGAAAG	GCATTTTATA	GCCTGGGCAT	1200
AAGTGAAATC	AGCAGAGCCT	CTGGGTGGAT	GTGTAGAAGG	CACTTCAAAA	TGCATAAACC	1260
TGTTACAATG	TTAAAAAAA	AAAAAAA				1289

(2) INFORMATION FOR SEQ ID NO:112:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 315 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:112:

```

Met Val Phe Thr Val Arg Leu Leu His Ile Phe Thr Val Asn Lys Gln
1           5           10           15

Leu Gly Pro Lys Ile Val Ile Val Ser Lys Met Met Lys Asp Val Phe
20           25           30

Phe Phe Leu Phe Phe Leu Gly Val Trp Leu Val Ala Tyr Gly Val Ala
35           40           45

Thr Glu Gly Leu Leu Arg Pro Arg Asp Ser Asp Phe Pro Ser Ile Leu
50           55           60

Arg Arg Val Phe Tyr Arg Pro Tyr Leu Gln Ile Phe Gly Gln Ile Pro
65           70           75           80

Gln Glu Asp Met Asp Val Ala Leu Met Glu His Ser Asn Cys Ser Ser
85           90           95

Glu Pro Gly Phe Trp Ala His Pro Pro Gly Ala Gln Ala Gly Thr Cys
100          105          110

Val Ser Gln Tyr Ala Asn Trp Leu Val Val Leu Leu Leu Val Ile Phe
115          120          125

Leu Leu Val Ala Asn Ile Leu Leu Val Asn Leu Leu Ile Ala Met Phe
130          135          140

Ser Tyr Thr Phe Gly Lys Val Gln Gly Asn Ser Asp Leu Tyr Trp Lys
145          150          155          160

Ala Gln Arg Tyr Arg Leu Ile Arg Glu Phe His Ser Arg Pro Ala Leu
165          170          175

Ala Pro Pro Phe Ile Val Ile Ser His Leu Arg Leu Leu Leu Arg Gln
180          185          190

Leu Cys Arg Arg Pro Arg Ser Pro Gln Pro Ser Ser Pro Ala Leu Glu
195          200          205

His Phe Arg Val Tyr Leu Ser Lys Glu Ala Glu Arg Lys Leu Leu Thr

```

87

210	215	220
Trp Glu Ser Val His Lys Glu Asn Phe Leu Leu Ala Arg Ala Arg Asp		
225	230	235 240
Lys Arg Glu Ser Asp Ser Glu Arg Leu Lys Arg Thr Ser Gln Lys Val		
	245	250 255
Asp Leu Ala Leu Lys Gln Leu Gly His Ile Arg Glu Tyr Gln Gln Arg		
	260	265 270
Leu Lys Val Leu Glu Arg Glu Val Gln Gln Cys Ser Arg Val Leu Gly		
	275	280 285
Trp Val Ala Glu Ala Leu Ser Arg Ser Ala Leu Leu Pro Pro Gly Gly		
	290	295 300
Pro Pro Pro Pro Asp Leu Pro Gly Ser Lys Asp		
305	310	315

(2) INFORMATION FOR SEQ ID NO:113:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 553 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:113:

Met Val Gln Arg Leu Trp Val Ser Arg Leu Leu Arg His Arg Lys Ala		
1	5	10 15
Gln Leu Leu Leu Val Asn Leu Leu Thr Phe Gly Leu Glu Val Cys Leu		
	20	25 30
Ala Ala Gly Ile Thr Tyr Val Pro Pro Leu Leu Leu Glu Val Gly Val		
	35	40 45
Glu Glu Lys Phe Met Thr Met Val Leu Gly Ile Gly Pro Val Leu Gly		
	50	55 60
Leu Val Cys Val Pro Leu Leu Gly Ser Ala Ser Asp His Trp Arg Gly		
	65	70 75 80
Arg Tyr Gly Arg Arg Arg Pro Phe Ile Trp Ala Leu Ser Leu Gly Ile		
	85	90 95
Leu Leu Ser Leu Phe Leu Ile Pro Arg Ala Gly Trp Leu Ala Gly Leu		
	100	105 110

88

Leu Cys Pro Asp Pro Arg Pro Leu Glu Leu Ala Leu Leu Ile Leu Gly
 115 120 125
 Val Gly Leu Leu Asp Phe Cys Gly Gln Val Cys Phe Thr Pro Leu Glu
 130 135 140
 Ala Leu Leu Ser Asp Leu Phe Arg Asp Pro Asp His Cys Arg Gln Ala
 145 150 155 160
 Tyr Ser Val Tyr Ala Phe Met Ile Ser Leu Gly Gly Cys Leu Gly Tyr
 165 170 175
 Leu Leu Pro Ala Ile Asp Trp Asp Thr Ser Ala Leu Ala Pro Tyr Leu
 180 185 190
 Gly Thr Gln Glu Glu Cys Leu Phe Gly Leu Leu Thr Leu Ile Phe Leu
 195 200 205
 Thr Cys Val Ala Ala Thr Leu Leu Val Ala Glu Glu Ala Ala Leu Gly
 210 215 220
 Pro Thr Glu Pro Ala Glu Gly Leu Ser Ala Pro Ser Leu Ser Pro His
 225 230 235 240
 Cys Cys Pro Cys Arg Ala Arg Leu Ala Phe Arg Asn Leu Gly Ala Leu
 245 250 255
 Leu Pro Arg Leu His Gln Leu Cys Cys Arg Met Pro Arg Thr Leu Arg
 260 265 270
 Arg Leu Phe Val Ala Glu Leu Cys Ser Trp Met Ala Leu Met Thr Phe
 275 280 285
 Thr Leu Phe Tyr Thr Asp Phe Val Gly Glu Gly Leu Tyr Gln Gly Val
 290 295 300
 Pro Arg Ala Glu Pro Gly Thr Glu Ala Arg Arg His Tyr Asp Glu Gly
 305 310 315 320
 Val Arg Met Gly Ser Leu Gly Leu Phe Leu Gln Cys Ala Ile Ser Leu
 325 330 335
 Val Phe Ser Leu Val Met Asp Arg Leu Val Gln Arg Phe Gly Thr Arg
 340 345 350
 Ala Val Tyr Leu Ala Ser Val Ala Ala Phe Pro Val Ala Ala Gly Ala
 355 360 365
 Thr Cys Leu Ser His Ser Val Ala Val Val Thr Ala Ser Ala Ala Leu
 370 375 380
 Thr Gly Phe Thr Phe Ser Ala Leu Gln Ile Leu Pro Tyr Thr Leu Ala
 385 390 395 400

Ser Leu Tyr His Arg Glu Lys Gln Val Phe Leu Pro Lys Tyr Arg Gly
 405 410 415
 Asp Thr Gly Gly Ala Ser Ser Glu Asp Ser Leu Met Thr Ser Phe Leu
 420 425 430
 Pro Gly Pro Lys Pro Gly Ala Pro Phe Pro Asn Gly His Val Gly Ala
 435 440 445
 Gly Gly Ser Gly Leu Leu Pro Pro Pro Pro Ala Leu Cys Gly Ala Ser
 450 455 460
 Ala Cys Asp Val Ser Val Arg Val Val Val Gly Glu Pro Thr Glu Ala
 465 470 475 480
 Arg Val Val Pro Gly Arg Gly Ile Cys Leu Asp Leu Ala Ile Leu Asp
 485 490 495
 Ser Ala Phe Leu Leu Ser Gln Val Ala Pro Ser Leu Phe Met Gly Ser
 500 505 510
 Ile Val Gln Leu Ser Gln Ser Val Thr Ala Tyr Met Val Ser Ala Ala
 515 520 525
 Gly Leu Gly Leu Val Ala Ile Tyr Phe Ala Thr Gln Val Val Phe Asp
 530 535 540
 Lys Ser Asp Leu Ala Lys Tyr Ser Ala
 545 550

(2) INFORMATION FOR SEQ ID NO:114:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 241 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:114:

Met Gln Cys Phe Ser Phe Ile Lys Thr Met Met Ile Leu Phe Asn Leu
 1 5 10 15
 Leu Ile Phe Leu Cys Gly Ala Ala Leu Leu Ala Val Gly Ile Trp Val
 20 25 30
 Ser Ile Asp Gly Ala Ser Phe Leu Lys Ile Phe Gly Pro Leu Ser Ser
 35 40 45
 Ser Ala Met Gln Phe Val Asn Val Gly Tyr Phe Leu Ile Ala Ala Gly

90

50		55		60
Val Val Val Phe Ala Leu Gly Phe Leu Gly Cys Tyr Gly Ala Lys Thr				
65		70		75
				80
Glu Ser Lys Cys Ala Leu Val Thr Phe Phe Phe Ile Leu Leu Leu Ile				
	85		90	
				95
Phe Ile Ala Glu Val Ala Ala Ala Val Val Ala Leu Val Tyr Thr Thr				
	100		105	
				110
Met Ala Glu His Phe Leu Thr Leu Leu Val Val Pro Ala Ile Lys Lys				
	115		120	
				125
Asp Tyr Gly Ser Gln Glu Asp Phe Thr Gln Val Trp Asn Thr Thr Met				
	130		135	
				140
Lys Gly Leu Lys Cys Cys Gly Phe Thr Asn Tyr Thr Asp Phe Glu Asp				
	145		150	
				155
				160
Ser Pro Tyr Phe Lys Glu Asn Ser Ala Phe Pro Pro Phe Cys Cys Asn				
	165		170	
				175
Asp Asn Val Thr Asn Thr Ala Asn Glu Thr Cys Thr Lys Gln Lys Ala				
	180		185	
				190
His Asp Gln Lys Val Glu Gly Cys Phe Asn Gln Leu Leu Tyr Asp Ile				
	195		200	
				205
Arg Thr Asn Ala Val Thr Val Gly Gly Val Ala Ala Gly Ile Gly Gly				
	210		215	
				220
Leu Glu Leu Ala Ala Met Ile Val Ser Met Tyr Leu Tyr Cys Asn Leu				
	225		230	
				235
				240
Gln				

(2) INFORMATION FOR SEQ ID NO:115:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 366 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:115:

GCTCTTTCTC	TCCCCCTCCTC	TGAATTTAAT	TCTTTCAACT	TGCAATTTGC	AAGGATTACA	60
CATTTCACTG	TGATGTATAT	TGTGTTGCAA	AAAAAAAAAA	GTGTCTTTGT	TTAAAATTAC	120
TTGGTTTGTG	AATCCATCTT	GCTTTTTCCC	CATTGGAAC	AGTCATTAAC	CCATCTCTGA	180
ACTGGTAGAA	AAACATCTGA	AGAGCTAGTC	TATCAGCATC	TGACAGGTGA	ATTGGATGGT	240
TCTCAGAACC	ATTTACCCCA	GACAGCCTGT	TTCTATCCTG	TTTAATAAAT	TAGTTTGGGT	300
TCTCTACATG	CATAACAAAC	CCTGCTCCAA	TCTGTACAT	AAAAGTCTGT	GACTTGAAGT	360
TTAGTC						366

(2) INFORMATION FOR SEQ ID NO:116:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 282 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:116:

ACAAAGATGA	ACCATTTCTT	ATATTATAGC	AAAATTAAAA	TCTACCCGTA	TTCTAATATT	60
GAGAAATGAG	ATNAAACACA	ATNTTATAAA	GTCTACTTAG	AGAAGATCAA	GTGACCTCAA	120
AGACTTTACT	ATTTTCATAT	TTTAAGACAC	ATGATTTATC	CTATTTTAGT	AACCTGGTTC	180
ATACGTTAAA	CAAAGGATAA	TGTGAACAGC	AGAGAGGATT	TGTTGCCAGA	AAATCTATGT	240
TCAATCTNGA	ACTATCTANA	TCACAGACAT	TTCTATTCTT	TT		282

(2) INFORMATION FOR SEQ ID NO:117:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 305 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:117:

ACACATGTGG	CTTCACTGCC	TTCTTAGATG	CTTCTGGTCA	ACATANAGGA	ACAGGGACCA	60
TATTTATCCT	CCCTCCTGAA	ACAATTGCAA	AATAANACAA	AATATATGAA	ACAATTGCAA	120
AATAAGGCAA	AATATATGAA	ACAACAGGTC	TCGAGATATT	GGAAATCAGT	CAATGAAGGA	180
TACTGATCCC	TGATCACTGT	CCTAATGCAG	GATGTGGGAA	ACAGATGAGG	TCACCTCTGT	240
GACTGCCCCA	GCTTACTGCC	TGTAGAGAGT	TTCTANGCTG	CAGTTCAGAC	AGGGAGAAAT	300
TGGGT						305

(2) INFORMATION FOR SEQ ID NO:118:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 71 base pairs

92

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:118:

ACCAAGGTGT NTGAATCTCT GACGTGGGGA TCTCTGATTC CCGCACAATC TGAGTGGAAA	60
AANTCCTGGG T	71

(2) INFORMATION FOR SEQ ID NO:119:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 212 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:119:

ACTCCGGTTG GTGTCAGCAG CACGTGGCAT TGAACATNGC AATGTGGAGC CCAAACCACA	60
CAAAATGGGG TGAAATTGGC CAACTTTCTA TNAACTTATG TTGGCAANTT TGCCACCAAC	120
AGTAAGCTGG CCCTTCTAAT AAAAGAAAAT TGAAAGGTTT CTCCTAANC GGAATTAANT	180
AATGGANTCA AGANACTCCC AGGCCTCAGC GT	212

(2) INFORMATION FOR SEQ ID NO:120:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 90 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:120:

ACTCGTTGCA NATCAGGGGC CCCCAGAGT CACCGTTGCA GGACTCCTTC TGGTCTTGCC	60
CTCCGCCGGC GCAGAACATG CTGGGGTGGT	90

(2) INFORMATION FOR SEQ ID NO:121:

(i) SEQUENCE CHARACTERISTICS:

93

- (A) LENGTH: 218 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:121:

TGTANCGTGA	ANACGACAGA	NAGGGTTGTC	AAAAATGGAG	AANCCTTGAA	GTCATTTTGA	60
GAATAAGATT	TGCTAAAAGA	TTGGGGGCTA	AAACATGGTT	ATTGGGAGAC	ATTTCTGAAG	120
ATATNCANGT	AAATTANGGA	ATGAATTCAT	GTTTCTTTTG	CGAATTCCTT	TACGATNGCC	180
AGCATANACT	TCATGTGGGG	ATANCAGCTA	CCCTTGTA			218

(2) INFORMATION FOR SEQ ID NO:122:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 171 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:122:

TAGGGGTGTA	TGCAACTGTA	AGGACAAAAA	TTGAGACTCA	ACTGGCTTAA	CCAATAAAGG	60
CATTTGTTAG	CTCATGGAAC	AGGAAGTCGG	ATGGTGGGGC	ATCTTCAGTG	CTGCATGAGT	120
CACCACCCCG	GCGGGGTCAT	CTGTGCCACA	GGTCCCTGTT	GACAGTGCGG	T	171

(2) INFORMATION FOR SEQ ID NO:123:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 76 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:123:

TGTAGCGTGA	AGACNACAGA	ATGGTGTGTG	CTGTGCTATC	CAGGAACACA	TTTATTATCA	60
TTATCAANTA	TTGTGT					76

(2) INFORMATION FOR SEQ ID NO:124:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 131 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:124:

```

ACCTTTCCCC AAGGCCAATG TCCTGTGTGC TAACTGGCCG GCTGCAGGAC AGCTGCAATT      60
CAATGTGCTG GGTCAATATG AGGGGAGGAG ACTGTAAAT AGCCAATTTT ATTCTCTTGG      120
TTAAGATTTG T                                     131

```

(2) INFORMATION FOR SEQ ID NO:125:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 432 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:125:

```

ACTTTATCTA CTGGCTATGA AATAGATGGT GGAAAATTGC GTTACCAACT ATACCACTGG      60
CTTGAAAAAG AGGTGATAGC TCTTCAGAGG ACTTGTGACT TTTGCTCAGA TGCTGAAGAA      120
CTACAGTCTG CATTTGCCAG AAATCAAGAT GAATTTGGAT TAAATGAGGA TGCTGAAGAT      180
TTGCCTCACC AAACAAAAGT GAAACAACCTG AGAGAAAATT TTCAGGAAAA AAGACAGTGG      240
CTCTTGAAGT ATCAGTCACT TTTGAGAATG TTTCTTAGTT ACTGCATACT TCATGGATCC      300
CATGGTGGGG GTCTTGCAAT TGTAAGAATG GAATTGATTT TGCTTTTGCA AGAATCTCAG      360
CAGGAAACAT CAGAACCACT ATTTTCTAGC CCTCTGTCAG AGCAAACCTC AGTGCCCTCTC      420
CTCTTTGCTT GT                                     432

```

(2) INFORMATION FOR SEQ ID NO:126:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 112 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:126:

ACACAACCTTG AATAGTAAAA TAGAACTGA GCTGAAATTT CTAATTCACCT TTCTAACCAT 60
AGTAAGAATG ATATTTCCCC CCAGGGATCA CCAAATATTT ATAAAAATTT GT 112

(2) INFORMATION FOR SEQ ID NO:127:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 54 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:127:

ACCACGAAAC CACAAACAAG ATGGAAGCAT CAATCCACTT GCCAAGCACA GCAG 54

(2) INFORMATION FOR SEQ ID NO:128:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 323 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:128:

ACCTCATTAG TAATTGTTTT GTTGTTTCAT TTTTTTCTAA TGTCTCCCCT CTACCAGCTC 60
ACCTGAGATA ACAGAATGAA AATGGAAGGA CAGCCAGATT TCTCCTTTGC TCTCTGCTCA 120
TTCTCTCTGA AGTCTAGGTT ACCCATTTTG GGGACCCATT ATAGGCAATA AACACAGTTC 180
CCAAAGCATT TGGACAGTTT CTTGTTGTGT TTTAGAATGG TTTTCCTTTT TCTTAGCCTT 240
TTCCTGCAAA AGGCTCACTC AGTCCCTTGC TTGCTCAGTG GACTGGGCTC CCCAGGGCCT 300
AGGCTGCCTT CTTTTCATG TCC 323

(2) INFORMATION FOR SEQ ID NO:129:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 192 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:129:

ACATACATGT GTGTATATTT TTAAATATCA CTTTTGTATC ACTCTGACTT TTTAGCATACT	60
TGAAAACACA CTAACATAAT TTNTGTGAAC CATGATCAGA TACAACCCAA ATCATTTCATC	120
TAGCACATTC ATCTGTGATA NAAAGATAGG TGAGTTTCAT TTCCTTCACG TTGGCCAATG	180
GATAAACAAA GT	192

(2) INFORMATION FOR SEQ ID NO:130:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 362 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:130:

CCCTTTTTTA TGAATGAGT AGACTGTATG TTTGAANATT TANCCACAAC CTCTTTGACA	60
TATAATGACG CAACAAAAAG GTGCTGTTTA GTCCTATGGT TCAGTTTATG CCCCTGACAA	120
GTTTCCATTG TGTTTTGCCG ATCTTCTGGC TAATCGTGGT ATCCTCCATG TTATTAGTAA	180
TTCTGTATTC CATTTTGTTA ACGCCTGGTA GATGTAACCT GCTANGAGGC TAACTTTATA	240
CTTATTTAAA AGCTCTTATT TTGTGTCAT TAAAAAGGCA ATTTATGTGC AGCACTTTAT	300
TGCAGCAGGA AGCAGTGTG GGTGTTGT AAAGCTCTT GCTAATCTTA AAAAGTAATG	360
GG	362

(2) INFORMATION FOR SEQ ID NO:131:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 332 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:131:

CTTTTTGAAA GATCGTGTCC ACTCCTGTGG ACATCTTSTT TTAATGGAGT TTCCCATGCA	60
GTANGACTGG TATGSTTGCA GCTGTCCAGA TAAAAACATT TGAAGAGCTC CAAAATGAGA	120
GTTCTCCAG GTTCCCTCTG CTGCTCCAAG TCTCAGCAGC AGCCTCTTTT AGGAGGATC	180
TTCTGAACTA GATTAAGGCA GTTGTAAT ETGATGTGAT TTGGTTTATT ATCCAACTAA	240
CTTCCATCTG TTATCACTG AGAAAGCCCA GACTCCCAAN GACNGGTACG GATTGTGGGC	300
ATANAAGGAT TGGGTGAAGC TGGCGTTGTG GT	332

(2) INFORMATION FOR SEQ ID NO:132:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 322 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:132:

ACTTTTGCCA TTTTGTATAT ATAAACAATC TTGGGACATT CTCCIGAAAA CTAGGTGTCC	60
AGTGGCTAAG AGAACTCGAT TTCAAGCAAT TCTGAAAGGA AAACCAGCAT GACACAGAAT	120
CTCAAATTCC CAAACAGGGG CTCTGTGGGA AAAATGAGGG AGGACCTTTG TATCTCGGGT	180
TTAGCAAGT TAAAATGAAN ATGACAGGAA AGGCTTATTT ATCAACAAAG AGAAGAGTTG	240
GGATGCTTCT AAAAAAACT TTGGTAGAGA AAATAGGAAT GCTNAATCCT AGGGAAGCCT	300
GTAACAATCT ACAATTGGTC CA	322

(2) INFORMATION FOR SEQ ID NO:133:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 278 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:133:

ACAAGCCTTC ACAAGTTTAA CTAAATTGGG ATTAATCTTT CTGTANTTAT CTGCATAATT	60
CTTGTTTTTC TTTCCATCTG GCTCCTGGGT TGACAATTTG TGGAAACAAC TCTATTGCTA	120
CTATTTAAAA AAAATCACAA ATCTTTCCCT TTAAGCTATG TTNAATTCAA ACTATTCCTG	180
CTATTCCTGT TTTGTCAAAG AAATTATATT TTTCAAAATA TGTNTATTTG TTTGATGGGT	240
CCCACGAAAC ACTAATAAAA ACCACAGAGA CCAGCCTG	278

(2) INFORMATION FOR SEQ ID NO:134:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 121 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:134:

GTTTANAAAA	CTTGTTTASC	TCCATAGAGG	AAAGAATGTT	AAACTTTGTA	TTTAAAAACA	60
TGATTCTCTG	AGGTTAAACT	TGTTTTTCAA	ATGTTATTTT	TACTTGTATT	TTGCTTTTGG	120
T						121

(2) INFORMATION FOR SEQ ID NO:135:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 350 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:135:

ACTTANAACC	ATGCCTAGCA	CATCAGAATC	CCTCAAAGAA	CATCAGTATA	ATCCTATACC	60
ATANCAAGTG	GTGACTGGTT	AAGCGTGCGA	CAAAGGTCAG	CTGGCACATT	ACTTGTGTGC	120
AAACTTGATA	CTTTTGTTCT	AAGTAGGAAC	TAGTATACAG	TNCCTAGGAN	TGGTACTCCA	180
GGGTGCCCCC	CAACTCCTGC	AGCCGCTCCT	CTGTGCCAGN	CCCTGNAAGG	AACTTTTCGCT	240
CCACCTCAAT	CAAGCCCTGG	GCCATGCTAC	CTGCAATTGG	CTGAACAAAC	GTTTGCTGAG	300
TTCCCAAGGA	TGCAAAGCCT	GGTGCTCAAC	TCCTGGGGCG	TCAACTCAGT		350

(2) INFORMATION FOR SEQ ID NO:136:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 399 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:136:

TGTACCGTGA	AGACGACAGA	AGTTGCATGG	CAGGGACAGG	GCAGGGCCGA	GGCCAGGGTT	60
GCTGTGATTG	TATCCGAATA	NTCCTCGTGA	GAAAAGATAA	TGAGATGACG	TGAGCAGCCT	120
GCAGACTTGT	GTCTGCCTTC	AANAAGCCAG	ACAGGAAGGC	CCTGCCTGCC	TTGGCTCTGA	180
CCTGGCGGCC	AGCCAGCCAG	CCACAGGTGG	GCTTCTTCCT	TTTGTGGTGA	CAACNCCAAG	240
AAAAGTGCAG	AGGCCAGGG	TCAGGTGTNA	GTGGGTANGT	GACCATAAAA	CACCAAGGTGC	300
TCCCAGGAAC	CCGGGCAAAAG	GCCATCCECA	CCTACAGCCA	GCATGCCCCAC	TGGCGTGATG	360
GGTGCAAGAG	GATGAASCAG	CCAGNTTCTC	TGCTGTGGT			399

(2) INFORMATION FOR SEQ ID NO:137:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 165 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:137:

ACTGGTGTGG TNGGGGGTGA TGCTGGTGGT ANAAGTTGAN GTGACTTCAN GATGGTGTGT	60
GGAGGAACTG TGTGAACGTA GGGATGTAGA NGTTTTGGCC GTGCTAAATG AGCTTCGGGA	120
TTGGCTGGTC CCACTGGTGG TCACTGTCAT TGGTGGGGTT CCTGT	165

(2) INFORMATION FOR SEQ ID NO:138:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 338 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:138:

ACTCACTGGA ATGCCACATT CACAACAGAA TCAGAGGTCT GTGAAAACAT TAATGGCTCC	60
TTAACTTCTC CAGTAAGAAT CAGGGACTTG AAATGGAAAC GTTAACAGCC ACATGCCCAA	120
TGCTGGGCAG TCTCCCATGC CTTCCACAGT GAAAGGGCTT GAGAAAAATC ACATCCAATG	180
TCATGTGTTT CCAGCCACAC CAAAAGGTGC TTGGGGTGGG GGGCTGGGGG CATANANGGT	240
CANGCCTCAG GAAGCCTCAA GTTCCATTCA GCTTTGCCAC TGTACATTCC CCATNTTTAA	300
AAAAACTGAT GCCTTTTTTT TTTTTTTTTT TAAAATTC	338

(2) INFORMATION FOR SEQ ID NO:139:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 382 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:139:

GGGAATCTTG GTTTTTGGCA TCTGGTTTGC CTATAGCCGA GGCCACTTTG ACAGAACAAA	60
GAAAGGGACT TCGASTAAGA AGGTGATTTA CAGCCAGCCT AGTGCCCGAA GTGAAGGAGA	120
ATTCAAACAG ACCTCGTCAT TCCTGGTGTG AGCCTGGTCG CCTCACCACC TATCATCTCC	180

100

ATTTGCCTTA CTCAGGTGCT ACCGGACTCT GSCCCCTGAT GTCTGTAGTT TCACAGGATG	240
CCTTATTTCT CTCTACAGC CCACAGGGCT CCTACTTCT TCGGATCTCT TTTTAATAAT	300
CTCAGCTATG TGCCCCATCC TCCTTCATGC CCTCCCTCCG TTTCCCTACCA CTGCTGAGTG	360
CCCTGGAAC T3TTTAAAT GT	382

(2) INFORMATION FOR SEQ ID NO:140:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 200 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:140:

ACCAAANCTT CTTTCTGTTG TGTTNGATTT TACTATAGGG GTTTNGCTTN TTCTAAANAT	60
ACTTTTCATT TAACANCTTT TGTTAAGTGT CAGGCTGCAC TTGCTCCAT ANAATTATTG	120
TTTTACATT TCAACTGTG TGTGTTTGT TCTTANAGCA TTGGTGAAAT CACATATTTT	180
ATATTCAGCA TAAAGGAGAA	200

(2) INFORMATION FOR SEQ ID NO:141:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 335 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:141:

ACTTTATTTT CAAAACACTC ATATGTTGCA AAAAACACAT AGAAAAATAA AGTTTGGTGG	60
GGGTGCTGAC TAAACTTCAA GTCACAGACT TTTATGTGAC AGATTGGAGC AGGGTTTGT	120
ATGCATGTAG AGAACCCAAA CTAATTTATT AACACAGGATA GAAACAGGCT GTCTGGGTGA	180
AATGGTTCTG AGAACCATCC AATTCACCTG TCAGATGCTG ATANACTAGC TCTTCAGATG	240
TTTTTCTACC AGTTCAGAGA TNGGTTAATG ACTANTTCCA ATGGGGAAAA AGCAAGATGG	300
ATTCACAAAC CAAGTAATTT TAAACAAAGA CACTT	335

(2) INFORMATION FOR SEQ ID NO:142:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 459 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

101

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:142:

ACCAGGTAA	TATTGCCACA	TATATCCTTT	CCAATTGCGG	GCTAAACAGA	CGTGTATTTA	60
GGGTTGTTTA	AAGACAACCC	AGCTTAATAT	CAAGAGAAAT	TGTGACCTTT	CATGCAGTAT	120
CTGATGGAGA	AAACACTGAG	TTTGTACAAA	TCTTATTTTA	TTCAGATAGC	AGTCTGATCA	180
CACATGGTCC	AACAACACTC	AAATAATAAA	TCAAATATNA	TCAGATGTTA	AAGATTGGTC	240
TTCAAACATC	ATAGCCAATG	ATGCCCCGCT	TGCCTATAAT	CTCTCCGACA	TAAAACCACA	300
TCAACACCTC	AGTGGCCACC	AAACCATTCA	GCACAGCTTC	CTTAAGTGTG	AGCTGTTTGA	360
AGCTACCACT	CTGAGCACTA	TTGACTATNT	TTTTCANGCT	CTGAATAGCT	CTAGGGATCT	420
CAGCANGGGT	GGGAGGAACC	AGCTCAACCT	TGGCGTANT			459

(2) INFORMATION FOR SEQ ID NO:143:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 140 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:143:

ACATTTTCCTT	CCACCAAGTC	AGGACTCCTG	GCTTCTGTGG	GAGTTCTTAT	CACCTGAGGG	60
AAATCCAAAC	AGTCTCTCCT	AGAAAGGAAT	AGTGTACCA	ACCCACCCA	TCTCCCTGAG	120
ACCATCCGAC	TTCCCTGTGT					140

(2) INFORMATION FOR SEQ ID NO:144:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 164 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:144:

ACTTCAGTAA	CAACATACAA	TAACAACATT	AAGTGTATAT	TGCCATCTTT	GTCATTTTCT	60
ATCTATACCA	CTCTCCCTTC	TGAAAACAA	N AATCACTANC	CAATCACTTA	TACAAATTTG	120
AGGCAATTAA	TCCATATTTG	TTTTCAATAA	GGAAAAAAG	ATGT		164

(2) INFORMATION FOR SEQ ID NO:145:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 303 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:145:

```

ACGTAGACCA TCCAACTTTG TATTTGTAAT GGCAAACATC CAGNAGCAAT TCCTAAACAA      60
ACTGGAGGGT ATTTATACCC AATTATCCCA TTCATTAACA TGCCCTCCTC CTCAGGCTAT      120
GCAGGACAGC TATCATAAGT CGGCCUAGG ATCCAGATAC TACCATTTGT ATAAACTTCA      180
GTAGGGGAGT CCATCCAAGT GACAGGTCTA ATCAAAGGAG GAAATGGAAC ATAAGCCCAG      240
TAGTAAATN TTGTTAGCT GAAACAGCCA CAAAAGACTT ACCGCCGTGG TGATTACCAT      300
CAA                                         303

```

(2) INFORMATION FOR SEQ ID NO:146:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 327 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:146:

```

ACTGCAGCTC AATTAGAAGT GGTCTCTGAC TTTTCATCANC TTCTCCCTGG GCTCCATGAC      60
ACTGGCCTGG AGTGACTCAT TGCTCTGGTT GGTGAGAGA GTCCTTTGC CAACAGGCCT      120
CCAAGTCAGG GCTGGGATTT GTTTCCTTTC CACATTCTAG CAACAATATG CTGGCCACTT      180
CCTGAACAGG GAGGGTGGGA GGAGCCAGCA TGAACAAGC TGCCACTTTC TAAAGTAGCC      240
AGACTTGCCC CTGGGCCTGT CACACCTACT GATGACCTTC TGTGCCTGCA GGATGGAATG      300
TAGGGGTGAG CTGTGTGACT CTATGGT                                         327

```

(2) INFORMATION FOR SEQ ID NO:147:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 173 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:147:

ACATTGTTTT	TTTGAGATAA	AGCATTGANA	GAGCTCTCCT	TAACGTGACA	CAATGGAAGG	60
ACTGGAACAC	ATACCCACAT	CTTTGTTCTG	AGGGATAATT	TTCTGATAAA	GTCTTGCTGT	120
ATATTCAAGC	ACATATGTTA	TATATTATTC	AGTTCATGT	TTATAGCCTA	GTT	173

(2) INFORMATION FOR SEQ ID NO:148:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 477 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:148:

ACAACCACTT	TATCTCATCG	AATTTTTTAAC	CCAAACTCAC	TCACTGTGCC	TTTCTATCCT	60
ATGGGATATA	TTATTTGATG	CTCCATTTCA	TCACACATAT	ATGAATAATA	CACTCATACT	120
GCCCTACTAC	CTGCTGCAAT	AATCACATT	CCTTCCTGTC	CTGACCCTGA	AGCCATTGGG	180
GTGGTCCTAG	TGGCCATCAG	TCCANGCCTG	CACCTTGAGC	CCTTGAGCTC	CATTGCTCAC	240
NCCANCCAC	CTCACCGACC	CCATCCTCTT	ACACAGCTAC	CTCCTTGCTC	TCTAACCCCA	300
TAGATTATNT	CCAAATTCAG	TCAATTAAGT	TACTATTAAC	ACTCTACCCG	ACATGTCCAG	360
CACCACTGGT	AAGCCTTCTC	CAGCCAACAC	ACACACACAC	ACACNCACAC	ACACACATAT	420
CCAGGCACAG	GCTACCTCAT	CTTCACAATC	ACCCCTTTAA	TTACCATGCT	ATGGTGG	477

(2) INFORMATION FOR SEQ ID NO:149:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 207 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:149:

ACAGTTGTAT	TATAATATCA	AGAAATAAAC	TTGCAATGAG	AGCATTTAAG	AGGGAAGAAC	60
TAACGTATTT	TAGAGAGCCA	AGGAAGGTTT	CTGTGGGGAG	TGGGATGTAA	GGTGGGGCCT	120
GATGATAAAT	AAGAGTCAGC	CAGGTAAGTG	GGTGGTGTGG	TATGGGCACA	GTGAAGAACA	180
TTTCAGGCAG	AGGGAACAGC	AGTGAAA				207

(2) INFORMATION FOR SEQ ID NO:150:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 111 base pairs

104

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:150:

ACCTTGATTT CATTGCTGCT CTGATGGAAA CCCAACTATC TAATTTAGCT AAAACATGGG	60
CAGTTAAATG TGGTCAGTGT TTGGACTTGT TAACTANTGG CATCTTTGGG T	111

(2) INFORMATION FOR SEQ ID NO:151:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 196 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:151:

AGCGCGGCAG GTCATATTGA ACATTCCAGA TACCTATCAT TACTCGATGC TGTTGATAAC	60
AGCAAGATGG CTTTGAAGTC AGGGTCACCA CCAGCTATTG GACCTTACTA TGAAAACCAT	120
GGATACCAAC CGGAAAACCC CTATCCCGCA CAGCCCACTG TGGTCCCCAC TGTCTACGAG	180
GTGCATCCGG CTCAGT	196

(2) INFORMATION FOR SEQ ID NO:152:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 132 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:152:

ACAGCACTTT CACATGTAAG AAGGGAGAAA TTCCTAAATG TAGGAGAAAAG ATAACAGAAC	60
CTTCCCCTTT TCATCTAGTG GTGGAAACCT GATGCTTTAT GTTGACAGGA ATAGAACCAG	120
GAGGGAGTTT GT	132

(2) INFORMATION FOR SEQ ID NO:153:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 285 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:153:

ACAAACCCCA NGANAGGCCA CTGGCCGTGG TGTCATGGCC TCCAAACATG AAAGTGTGAG	60
CTTCTGCTCT TATGTCTCTA TCTGACAACT CTTTACCATT TTTATCCTCG CTCAGCAGGA	120
GCACATCAAT AAAGTCCAAA CTCTTGACT TGGCCTTGGC TTGGAGGAAG TCATCAACAC	180
CCTGGCTAGT GAGGGTGCGG CGCCGCTCCT GGATGACGGC ATCTGTGAAG TCGTGCACCA	240
GTCTGCAGGC CCTGTGGAAG CGCCGTCCAC ACGGACTNAG GAATT	285

(2) INFORMATION FOR SEQ ID NO:154:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 333 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:154:

ACCACAGTCC TGTTGGGCCA GGGCTTCATG ACCCTTTCTG TGAAAAGCCA TATTATCACC	60
ACCCCAAATT TTTCTTAAA TATCTTTAAC TGAAGGGGTC AGCCTCTTGA CTGCAAAGAC	120
CCTAAGCCGG TTACACAGCT AACTCCCACT GGCCCTGATT TGTGAAATTG CTGCTGCCTG	180
ATTGGCACAG GAGTCGAAGG TGTTCACTC CCCTCCTCCG TGGAAACGAGA CTCTGATTTG	240
AGTTTCACAA ATTCTCGGGC CACCTCGTCA TTGCTCCTCT GAAATAAAAT CCGGAGAATG	300
GTCAGGCCTG TCTCATCCAT ATGGATCTTC CGG	333

(2) INFORMATION FOR SEQ ID NO:155:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 308 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:155:

106

ACTGGAAATA ATAAAACCCA CATCACAGTC TTGTGTCAAA GATCATCAGG GCATGGATGG	60
GAAAGTGCTT TGGGAAGTGT AAAGTGCCTA ACACATGATC GATGATTTT GTTATAATAT	120
TTGAATCAGG GTGCATAAA ACTCTCCTGC CTGCTCCTCC TGGGCCCCAG CCCCAGCCCC	180
ATCACAGCTC ACTGCTCTGT TCATCCAGGC CCAGCATGTA GTGGCTGATT CTTCTTGGCT	240
GCTTTTAGCC TCCANAAGTT TCTCTGAAGC CAACCAAACC TCTANGTGTA AGGCATGCTG	300
GCCCTGGT	308

(2) INFORMATION FOR SEQ ID NO:156:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 295 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:156:

ACCTTGCTCG GTGCTTGGAA CATATTAGGA ACTCAAAATA TGAGATGATA ACAGTGCCTA	60
TTATTGATTA CTGAGAGAAC TGTTAGACAT TTAGTTGAAG ATTTTCTACA CAGGAACTGA	120
GAATAGGAGA TTATGTTTGG CCCTCATATT CTCTCCTATC CTCCTTGCCT CATTCTATGT	180
CTAATATATT CTCAATCAAA TAAGGTTAGC ATAATCAGGA AATCGACCAA ATACCAATAT	240
AAAACCAGAT GTCTATCCTT AAGATTTTCA AATAGAAAAC AAATTAACAG ACTAT	295

(2) INFORMATION FOR SEQ ID NO:157:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 126 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:157:

ACAAGTTTAA ATAGTGCTGT CACTGTGCAT GTGCTGAAAT GTGAAATCCA CCACATTTCT	60
GAAGAGCAAA ACAAATTCTG TCATGTAATC TCTATCTTGG GTCGTGGGTA TATCTGTCCC	120
CTTAGT	126

(2) INFORMATION FOR SEQ ID NO:158:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 442 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:158:

ACCCACTGGT CTTGGAAACA CCCATCCTTA ATACGATGAT TTTTCTGTCT TGTGAAAATG	60
AANCCAGCAG GCTGCCCCCTA GTCAGTCCTT CCTTCCAGAG AAAAAGAGAT TTGAGAAAGT	120
GCCTGGGTAA TTCACCATTA ATTTCTCTCC CCAAACCTCTC TGAGTCTTCC CTTAATATTT	180
CTGGTGGTTC TGACCAAAGC AGGTCATGGT TTGTTGAGCA ITTGGGATCC CAGTGAAGTA	240
NATGTTTGTA GCCTTGCATA CTTAGCCCTT CCCACGCACA AACGGAGTGG CAGAGTGGTG	300
CCAACCCTGT TTTCCAGTC CACGTAGACA GATTACAGT GCGGAATTCT GGAAGCTGGA	360
NACAGACGGG CTCTTTGCAG AGCCGGGACT CTGAGANGGA CATGAGGGCC TCTGCCTCTG	420
TGTTCAATCT CTGATGTCCT GT	442

(2) INFORMATION FOR SEQ ID NO:159:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 498 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:159:

ACTTCCAGGT AACGTTGTTG TTTCCGTTGA GCCTGAACTG ATGGGTGACG TTGTAGGTTT	60
TCCAACAAGA ACTGAGGTTG CAGAGCGGGT AGGGAAGAGT GCTGTTCCAG TTGCACCTGG	120
GCTGCTGTGG ACTGTTGTTG ATTCCTCACT ACGGCCCAAG GTTGTGGAAC TGGCANAAAG	180
GTGTGTTGTT GGANTTGAGC TCGGGCGGCT GTGGTAGGTT GTGGGCTCTT CAACAGGGGC	240
TGCTGTGGTG CCGGGANGTG AANGTGTGTT GTCACTTGAG CTTGGCCAGC TCTGGAAAGT	300
ANTANATTCT TCCTGAAGGC CAGCGCTTGT GGAGCTGGCA NGGGTCANTG TTGTGTGTAA	360
CGAACCAGTG CTGCTGTGGG TGGGTGTANA TCCTCCACAA AGCCTGAAGT TATGGTGTGN	420
TCAGGTAANA ATGTGGTTTC AGTGTCCCTG GGCNGCTGTG GAAGGTTGTA NATTGTCAAC	480
AAGGGAATAA GCTGTGGT	498

(2) INFORMATION FOR SEQ ID NO:160:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 380 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:160:

ACCTGCATCC	AGCTTCCCTG	CCAAACTCAC	AAGGAGACAT	CAACCTCTAG	ACAGGGAAAC	60
AGCTTCAGGA	TACTTCCAGG	AGACAGAGCC	ACCAGCAGCA	AAACAAATAT	TCCCATGCCT	120
GGAGCATGGC	ATAGAGGAAG	CTGANAAATG	TGGGGTCTGA	GGAAGCCATT	TGAGTCTGGC	180
CACTAGACAT	CTCATCAGCT	ACTTGTGTGA	AGAGATGCCC	CATGACCCCA	GATGCCTCTC	240
CCACCCTTAC	CTCCATCTCA	CACACTTGAG	CTTTCCACTC	TGTATAATTC	TAACATCCTG	300
GAGAAAAATG	GCAGTTTGAC	CGAACCTGTT	CACAACGGTA	GAGGCTGATT	TCTAACGAAA	360
CTGTAGAAT	GAAGCCTGGA					380

(2) INFORMATION FOR SEQ ID NO:161:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 114 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:161:

ACTCCACATC	CCCTCTGAGC	AGGCGGTTGT	CGTTCAAGGT	GTATTTGGCC	TTGCCTGTCA	60
CACTGTCCAC	TGGCCCCCTA	TCCACTTGGT	GCTTAATCCC	TCGAAAGAGC	ATGT	114

(2) INFORMATION FOR SEQ ID NO:162:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 177 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:162:

ACTTTCTGAA	TCGAATCAAA	TGATACTTAG	TGTAGTTTTA	ATATCCTCAT	ATATATCAAA	60
GTTTTACTAC	TCTGATAATT	TTGTAAACCA	GGTAACCAGA	ACATCCAGTC	ATACAGCTTT	120
TGGTGATATA	TAACTTGGCA	ATAACCCAGT	CTGGTGATAC	ATAAACTAC	TCACTGT	177

(2) INFORMATION FOR SEQ ID NO:163:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 137 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

109

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:163:

CATTTATACA GACAGGCGTG AAGACATTCA CGACAAAAAC GCGAAATTCT ATCCCGTGAC	60
CANAGAAGGC AGCTACGGCT ACTCCTACAT CCTGGCGTGG GTGGCCTTCC CCTGCACCTT	120
CATCAGCGGC ATGATGT	137

(2) INFORMATION FOR SEQ ID NO:164:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 469 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:164:

CTTATCACAA TGAATGTTCT CCTGGGCAGC GTTGTGATCT TTGCCACCTT CGTGACTTTA	60
TGCAATGCAT CATGCTATTT CATACTAAT GAGGGAGTTC CAGGAGATTC AACCAGGAAA	120
TGCATGGATC TCAAAGGAAA CAAACACCCA ATAAACTCGG AGTGGCAGAC TGACAACTGT	180
GAGACATGCA CTTGCTACGA AACAGAAATT TCATGTTGCA CCCTTGTTTC TACACCTGTG	240
GGTTATGACA AAGACAAC TG CCAAAGAATC TTCAAGAAGG AGGACTGCAA GTATATCGTG	300
GTGGAGAAGA AGGACCCAAA AAAGACCTGT TCTGTCAAGT AATGGATAAT CTAATGTGCT	360
TCTAGTAGGC ACAGGGCTCC CAGGCCAGGC CTCATTCTCC TCTGGCCTCT AATAGTCAAT	420
GATTGTGTAG CCATGCCTAT CAGTAAAAAG ATNTTTGACC AAACACTTT	469

(2) INFORMATION FOR SEQ ID NO:165:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 195 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:165:

ACAGTTTTTT ATANATATCG ACATTGCCGG CACTTGTGTT CAGTTTCATA AAGCTGGTGG	60
ATCCGCTGTC ATCCACTATT COTTGGCTAG AGTAAAAATT ATTCTTATAG CCCATGTCCC	120
TGCAGGCCGC CCGCCGCTAG TTCTCGTTCC AGTCGTCTTG GCACACAGGG TGCCAGGACT	180
TCCTCTGAGA TGAGT	195

(2) INFORMATION FOR SEQ ID NO:166:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 383 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:166:

```

ACATCTTACT AGTGTGGBAC ATCAGGGGGG CATCAGGGTC ACASTCACTC ATAGCCTCCT      60
CGAGGTGGGA GTTCACACCA CCGGTGTAGG TGTGCTCAAT CTGGGCTTG GCGCCACCT      120
TTGGAGAAGG GATATGCTGC ACACACATGT CCACAAAGCC TGTGAAGTOS CCAAAGAATT      180
TTTGACAGAC AGCTGTAGCA AGGGGCGGAT GTTCAGCTTC AGCTCCTCCT TCCTCAGGTG      240
GATGCCAACC TCGTCTAMGG TCCGTGGGAA GCTGGTGTCC ACNTCACCTA CAACCTGGGG      300
GANGATCTTA TAAAGAGGCT CCNAGATAAA CTCCACGAAA CTTCTCTGGG AGCTGCTAGT      360
NGGGGCCCTT TTGGTGAACT TTC                                     383

```

(2) INFORMATION FOR SEQ ID NO:167:

- (i) SEQUENCE CHARACTERISTICS
 (A) LENGTH: 247 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:167:

```

ACAGAGCCAG ACCTTGGCCA TAAATGAANC AGAGATTAAG ACTAAACCCC AAGTCGANAT      60
TGGAGCAGAA ACTGGAGCAA GAAGTGGGCC TGGGGCTGAA GTAGAGACCA AGGCCACTGC      120
TATANCCATA CACAGAGCCA ACTCTCAGGC CAAGGCNATG GTTGGGGCAG ANCCAGAGAC      180
TCAATCTGAN TCCAAAGTGG TGGCTGGAAC ACTGGTCATG ACANAGGCA'S TGACTCTGAC      240
TGANGTC                                     247

```

(2) INFORMATION FOR SEQ ID NO:168

- (i) SEQUENCE CHARACTERISTICS.
 (A) LENGTH: 273 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (vi) ORIGINAL SOURCE:

111

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:168:

ACTTCTAAGT TTTCTAGAAG TGGAAGGATT GTANTCATCC TGAAAATGGG TTTACTTCAA	60
AATCCCTCAN CCTTGTTCTT CACNACTGTC TATACTGANA GTGTCATGTT TCCACAAAGG	120
GCTGACACCT GAGCCTGNAT TTTCACTCAT CCCTGAGAAG CCCTTTCCAG TAGGGTGGGC	180
AAATCCCAAC TTCCTTGCCA CAAGCTTCCC AGGCTTTCTC CCCTGGAAAA CTCCAGCTTG	240
AGTCCCAGAT AACTCATGG GCTGCCCTGG GCA	273

(2) INFORMATION FOR SEQ ID NO:169:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 431 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:169:

ACAGCCTTGG CTTCCCCAAA CTCCACAGTC TCAGTGCAGA AAGATCATCT TCCAGCAGTC	60
AGCTCAGACC AGGGTCAAAG GATGTGACAT CAACAGTTTC TGGTTTCAGA ACAGGTTCTA	120
CTACTGTCAA ATGACCCCCC ATACTTCCTC AAAGGCTGTG GTAAGTTTGG CACAGGTGAG	180
GGCAGCAGAA AGGGGGTANT TACTGATGGA CACCATCTTC TCTGTATACT CCACACTGAC	240
CTTGCCATGG GCAAAGGCCC CTACCACAAA AACAATAGGA TCACTGCTGG GCACCAGCTC	300
ACGCACATCA CTGACAACCG GGATGGAAAA AGAANTGCCA ACTTTCATAC ATCCAAGTGG	360
AAAGTGATCT GATACTGGAT TCTTAATTAC CTTCAAAAGC TTCTGGGGGC CATCAGCTGC	420
TCGAACACTG A	431

(2) INFORMATION FOR SEQ ID NO:170:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 266 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:170:

ACCTGTGGGC TGGGCTGTTA TGCCTGTGCC GGCCTGCTGAA AGGGAGTTCA GAGGTGGAGC	60
TCAAGGAGCT CTGCAGGCAT TTTGCCAANC CTCTCCANAG CANAGGGAGC AACCTACACT	120
CCCCGCTAGA AAGACACCAG ATTGGAGTCC TGGGAGGGGG AGTTGGGGTG GGCATTTGAT	180
GTATACTTGT CACCTGAATG AANGAGCCAG AGAGGAANGA GACGAANATG ANATTGGCCT	240
TCAAAGCTAG GGGTCTGGCA GGTGGA	266

112

(2) INFORMATION FOR SEQ ID NO:171:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1248 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:171:

```

GGCAGCCAAA TCATAAACGG AGAGGACTTG AGCCCGCAGT GGCAGCCCTG GCAGGCGGCA      60
CTGGTCATGG AAAACGAATT GTTCTGCTGG GCGGTGCTGG TGCATCCGCA CTGGGTGCTG      120
TCAGCCGCAC ACTGTTTCCA GAAGTGAGTG CAGAGTTCCT ACACCATCGG GCTGGGCGCTG      180
CACAGTCTTG AGGCGGACCA AGAGCCAGGG AGCCAGATGG TGGAGGGCCAG CCTCTCCGTA      240
CGGCACCCAG AGTACAACAG ACCCTTGCTC GCTAACGACC TCATGCTCAT CAAGTTGGAC      300
GAATCCGTGT CCGAGTCTGA CACCATCCGG AGCATCAGCA TTGCTTCGCA GTGCCCTACC      360
GCGGGGAACCT CTTGCCCTCT TTCTGGCTGG GGTCTGCTGG CGAACGGCAG AATGCCCTAC      420
GTGCTGCAGT GCGTGAACGT GTCGGTGGTG TCTGAGGAGG TCTGCAGTAA GCTCTATGAC      480
CCGCTGTACC ACCCCAGCAT GTTCTGCGCC GCGGGAGGGC AAGACCAGAA GGACTCCTGC      540
AACGGTGACT CTGGGGGGCC CCTGATCTGC AACGGGTACT TGCAGGGCCT TGTGTCTTTC      600
GGAAAAGCCC CGTGTGGCCA AGTTGGCGTG CCAGGTGTCT ACACCAACCT CTGCAAATTC      660
ACTGAGTGGA TAGAGAAAAC CGTCCAGGCC AGTTAACTCT GGGGACTGGG AACCCTATGAA      720
ATTGACCCCC AAATACATCC TGCGGAAGGA ATTCAGGAAT ATCTGTTCCC AGCCCTCCTT      780
CCCTCAGGCC CAGGAGTCCA GGCCCCCAGC CCCTCCTCCC TCAAACCAAG GGTACAGATC      840
CCCAGCCCCC CTCTCCTCAG ACCCAGGAGT CCAGACCCCC CAGCCCCCTCC TCCCTCAGAC      900
CCAGGAGTCC AGCCCTCCT CCCTCAGAAC CAGGAGTCCA GACCCCCCAG CCCCTCCTCC      960
CTCAGACCCA GGGGTCCAGG CCCCCAACCC CTCCTCCCTC AGACTCAGAG GTCCAAGCCC      1020
CCAACCCNTC ATTCCCAGA CCCAGAGGTC CAGGTCCCAG CCCCTCNTCC CTCAGACCCA      1080
GCGGTCCAAT GCCACCTAGA CTNTCCCTGT ACACAGTGCC CCCTTGTTGGC ACGTTGACCC      1140
AACCTTACCA GTTGSTTTTT CATTTTNGT CCCTTTCCCC TAGATCCAGA AATAAAGTTT      1200
AAGAGAAGNG CAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAA      1248

```

(2) INFORMATION FOR SEQ ID NO:172:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 159 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:172:

```

Met Val Glu Ala Ser Leu Ser Val Arg His Pro Glu Tyr Asn Arg Pro
1           5           10           15

```

113

Leu Leu Ala Asn Asp Leu Met Leu Ile Lys Leu Asp Glu Ser Val Ser
 20 25 30
 Glu Ser Asp Thr Ile Arg Ser Ile Ser Ile Ala Ser Gln Cys Pro Thr
 35 40 45
 Ala Gly Asn Ser Cys Leu Val Ser Gly Trp Gly Leu Leu Ala Asn Gly
 50 55 60
 Arg Met Pro Thr Val Leu Gln Cys Val Asn Val Ser Val Val Ser Glu
 65 70 75 80
 Glu Val Cys Ser Lys Leu Tyr Asp Pro Leu Tyr His Pro Ser Met Phe
 85 90 95
 Cys Ala Gly Gly Gly Gln Xaa Gln Xaa Asp Ser Cys Asn Gly Asp Ser
 100 105 110
 Gly Gly Pro Leu Ile Cys Asn Gly Tyr Leu Gln Gly Leu Val Ser Phe
 115 120 125
 Gly Lys Ala Pro Cys Gly Gln Val Gly Val Pro Gly Val Tyr Thr Asn
 130 135 140
 Leu Cys Lys Phe Thr Glu Trp Ile Glu Lys Thr Val Gln Ala Ser
 145 150 155

(2) INFORMATION FOR SEQ ID NO:173:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1265 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:173:

GGCAGCCCGC	ACTCGCAGCC	CTGGCAGGCG	GCACTGGTCA	TGGAAAACGA	ATTGTTCTGC	60
TCGGGCGTCC	TGGTGCATCC	GCAGTGGGTG	CTGTCAGCCG	CACACTGTTT	CCAGAACTCC	120
TACACCATCG	GGCTGGGCCT	GCACAGTCTT	GAGGCCGACC	AAGAGCCAGG	GAGCCAGATG	180
GTGGAGGCCA	GCCTCTCCGT	ACGGCACCCA	GAGTACAACA	GACCTTGCT	CGCTAACGAC	240
CTCATGCTCA	TCAAGTTGGA	CGAATCCGTG	TCCGAGTCTG	ACACCATCCG	GAGCATCAGC	300
ATTGCTTCGC	AGTGCCCTAC	CGCGGGGAAC	TCTTGCTCTG	TTTCTGGCTG	GGGTCTGCTG	360
GCGAACGGTG	AGCTACGGG	TGTGTGTCTG	CCCTCTTCAA	GGAGGTCTCT	TGCCCAGTCG	420
CGGGGGCTGA	CCCAGAGCTC	TGCGTCCCAG	GCAGAATGCC	TACCGTGCTG	CAGTGCGTGA	480
ACGTGTCTGGT	GGTGTCTGAG	GAGGTCTGCA	GTAAGCTCTA	TGACCCGCTG	TACCAACCCA	540
GCATGTTCTG	CGCCGGCGGA	GGGCAAGACC	AGAAGGACTC	CTGCAACGGT	GACTCTGGGG	600
GGCCCCTGAT	CTGCAACGGG	TACTTGACAG	GCCTTGCTGC	TTTCGGAAAA	GCCCCGTGTG	660
GCCAAGTTGG	CGTGCCAGGT	GTCTACACCA	ACCTCTGCAA	ATTCAGTGAG	TGGATAGAGA	720

AAACCGTCCA	GGCCAGTTAA	CTCTGGGGAC	TGGGAACCCA	TSAAATTBAC	CCCCAAATAC	780
ATCCTGCCGA	AGGAATTCAG	GAATATCTGT	TCCCAGCCTT	TCTTCCCTCA	GGCCCAAGGAG	840
TCCAGGCCCTC	CAGCCCCTCC	TCCCTCAAAAC	CAAGGGTACA	BATCCCCAGC	CCCTCCTCCC	900
TCAGACCCAG	GAGTCCAGAC	CCCCCAGCCC	CTCTTCCCTC	AGACCCAGGA	CTCCAGCCCC	960
TCTTCCNTCA	GACCCAGGAG	TCCAGACCCC	CCAGCCCCTC	CTCCCTCAGA	CCCAGGGGTT	1020
GAGGCCCCCA	ACCCCTCCTC	CTTCAGAGTC	AGAGGTCCAA	CCCCCAACG	CTCTGTTCCC	1080
CAGACCCAGA	GGTNAGGTC	CCAGCCCCTC	TTCCNTCAGA	CCCAGNGGTC	CAATGCCACC	1140
TAGATTTTCC	CTGNACACAG	TGCCCCCTTG	TGGNANGTTG	ACCCAACCTT	ACCAGTTGGT	1200
TTTTCATTTT	TNGTCCCTTT	CCCCTAGATC	CAGAAATAAA	GTTTAAGAGA	NGNGCAAAAA	1260
AAAAA						1265

(2) INFORMATION FOR SEQ ID NO:174:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1459 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:174:

GGTCAGCCGC	ACACTGTTTC	CAGAAGTGAG	TGCAGAGCTC	CTACACCATC	GGGCTGGGCC	50
TGCACAGTCT	TGAGGCCGAC	CAAGAGCCAG	GGAGCCAGAT	GGTGGAGGCC	AGCCTCTCCG	120
TACGGCACCC	AGAGTACAAC	AGACCCTTGC	TCGCTAACGA	CCTCATGCTC	ATCAAGTTGG	180
ACGAATCCGT	GTCCGAGTCT	GACACCATCC	GGAGCATCAG	CATTGCTTCG	CAGTGCCCTA	240
CCGCGGGGAA	CTCTTGCCCT	GTTTCTGGCT	GGGGTCTGCT	GGCGAACGGT	GAGCTCACGG	300
GTGTGTGTCT	GCCCTCTTCA	AGGAGGTCCT	CTGCCCAGTC	GCGGGGGCTG	ACCCACAGCT	360
CTGCGTCCCA	GGCAGAATGC	CTACCGTGCT	GCAGTGCGTG	AACGTGTCCG	TGGTGTCTGA	420
NGAGGTCTGC	ANTAAGCTCT	ATGACCCGCT	GTACCACCCC	ANCATGTTCT	GCGCCGGCGG	480
AGGGCAAGAC	CAGAAGGACT	CCTGCAACGT	GAGAGAGGGG	AAAGGGGAGG	GCAGGCGACT	540
CAGGGAAGGG	TGGAGAAGGG	GGAGACAGAG	ACACACAGGG	CCGCATGGCG	AGATGCAGAG	600
ATGGAGAGAC	ACACAGGGAG	ACAGTGACAA	CTAGAGAGAG	AAACTGAGAG	AAACAGAGAA	660
ATAAACACAG	GAATAAAGAG	AAGCAAAGGA	AGAGAGAAAC	AGAAACAGAC	ATGGGGAGGC	720
AGAAACACAC	ACACATAGAA	ATGCAGTTGA	CCTTCCAACA	GCATGGGGCC	TGAGGGCGGT	780
GACCTCCACC	CAATAGAAAA	TCCTCTTATA	ACTTTTGACT	CCCCAAAAAC	CTGACTAGAA	840
ATAGCCTACT	GTTGACGGGG	AGCCTTACCA	ATAACATAAA	TAGTCGATTT	ATGCATACGT	900
TTTATGCATT	CATGATATAC	CTTTSTTGGA	ATTTTTTGAT	ATTTCTAAGC	TACACAGTTC	960
GTCTGTGAAT	TTTTTTAAAT	TGTTSCAACT	CTCCTAAAAAT	TTTTCTGATG	TGTTTATTGA	1020
AAAAATCCAA	GTATAAGTGG	ACTTGTGCAT	TCAAACCAGG	GTTGTTCAAG	GSTCAACTGT	1080
GTACCCAGAG	GGAAACAGTG	ACACAGATTC	ATAGAGGTGA	AACACGAAGA	GAAACAGGAA	1140
AAATCAAGAC	TCTACAAAGA	GGCTGGGCAG	GTTGGCTCAT	GCCTGTAATC	CCAGCACTTT	1200
GGGAGGCGAG	GCAGGCAGAT	CACTTGAGGT	AAGGAGTTCA	AGACCAGCCT	GCCCAAAATG	1260
GTGAAATCCT	GTCTGTACTA	AAAATACAAA	AATTAGCTGC	ATATGGTGGC	AAGGCGCTGT	1320
AATCCCAGCT	ACTTGGGAGG	CTGAGGCAGG	ACAATTGCTT	GAATATGGGA	GACAGAGGTT	1380
GAAGTGAGTT	GAGATCACAC	CACTATACTC	CAGCTGGGGC	AACAGAGTAA	GACTCTGTCT	1440
CAAAAAAAAA	AAAAAAAAAA					1459

(2) INFORMATION FOR SEQ ID NO:175:

(i) SEQUENCE CHARACTERISTICS:

115

- (A) LENGTH: 1167 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:175.

```

GCGCAGCCCT GGCAGGCGGC ACTGGTCATG GAAAACGAAT TGTTCCTGCTC GGGCGTCCTG      60
GTGCATCCGC AGTGGGTGCT GTCAGCCGCA CACTGTTTCC AGAACTCCTA CACCATCGGG      120
CTGGGCCTGC ACAGTCTTGA GGCCGACCAA GAGCCAGGGA GCCAGATGGT GGAGGCCAGC      180
CTCTCCGTAC GGCACCCAGA CTACAACAGA CTCTTGCTCG CTAACGACCT CATCCTCATC      240
AAGTTGGACG AATCCGTGTC CGAGTCTGAC ACCATCCGGA GCATCAGCAT TGCTTCGCAG      300
TGCCCTACCG CGGGGAACCTC TTCCCTCGTN TCTGGCTGGG GTCTGCTGGC GAACGGCAGA      360
ATGCCTACCG TGCTGCACTG CGTGAACGTG TCGGTGGTGT CTGAGGANGT CTGCAGTAAG      420
CTCTATGACC CGCTGTACCA CCCCAGCATG TTCTGCGCCG GCGGAGGGCA AGACCAGAAG      480
GACTCCTGCA ACGGTGACTC TGGGGGGCCC CTGATCTGCA ACGGGTACTT GCAGGGCCTT      540
GTGTCTTTTC GAAAAGCCCC GTGTGGCCAA CTTGGCGTGC CAGGTGTCTA CACCAACCTC      600
TGCAAATTCA CTGAGTGGAT AGAGAAAACC GTCCAGNCCA GTTAACTCTG GGGACTGGGA      660
ACCCATGAAA TTGACCCCCA AATACATCCT GCGGAANGAA TTCAGGAATA TCTGTTCCCA      720
GCCCCCTCCT CCTCAGGCCC AGGAGTCCAG GCCCCAGCC CCTCCTCCCT CAAACCAAGG      780
GTACAGATCC CCAGCCCCTC CTCCCTCAGA CCCAGGAGTC CAGACCCCCC AGCCCCTCNT      840
CCNTCAGACC CAGGAGTCCA GCCCCCTCCT CNTCAGACGC AGGAGTCCAG ACCCCCCAGC      900
CCNTCNTCCG TCAGACCCAG GGGTGCAGGC CCCCACCCC TCNTCCNTCA GAGTCAGAGG      960
TCCAAGCCCC CAACCCCTCG TTCCCCAGAC CCAGAGGTNC AGGTCCCAGC CCCTCCTCCC     1020
TCAGACCCAG CGGTCCAATG CCACCTAGAN TNTCCCTGTA CACAGTGCCC CCTTGTTGGCA     1080
NGTTGACCCA ACCTTACCAG TTGGTTTTTC ATTTTTTGTC CCTTTCCCTT AGATCCAGAA     1140
ATAAAGTNTA AGAGAAGCGC AAAAAAA
                                                                                   1167
  
```

(2) INFORMATION FOR SEQ ID NO:176:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 205 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:176:

```

Met Glu Asn Glu Leu Phe Cys Ser Gly Val Leu Val His Pro Gln Trp
1           5           10           15

Val Leu Ser Ala Ala His Cys Phe Gln Asn Ser Tyr Thr Ile Gly Leu
20           25           30

Gly Leu His Ser Leu Glu Ala Asp Gln Glu Pro Gly Ser Gln Met Val
  
```

116

35	40	45
Glu Ala Ser Leu Ser Val Arg His Pro Glu Tyr Asn Arg Leu Leu Leu		
50	55	60
Ala Asn Asp Leu Met Leu Ile Lys Leu Asp Glu Ser Val Ser Glu Ser		
65	70	75 80
Asp Thr Ile Arg Ser Ile Ser Ile Ala Ser Gln Cys Pro Thr Ala Gly		
85	90	95
Asn Ser Cys Leu Val Ser Gly Trp Gly Leu Leu Ala Asn Gly Arg Met		
100	105	110
Pro Thr Val Leu His Cys Val Asn Val Ser Val Val Ser Glu Xaa Val		
115	120	125
Cys Ser Lys Leu Tyr Asp Pro Leu Tyr His Pro Ser Met Phe Cys Ala		
130	135	140
Gly Gly Gly Gln Asp Gln Lys Asp Ser Cys Asn Gly Asp Ser Gly Gly		
145	150	155 160
Pro Leu Ile Cys Asn Gly Tyr Leu Gln Gly Leu Val Ser Phe Gly Lys		
165	170	175
Ala Pro Cys Gly Gln Leu Gly Val Pro Gly Val Tyr Thr Asn Leu Cys		
180	185	190
Lys Phe Thr Glu Trp Ile Glu Lys Thr Val Gln Xaa Ser		
195	200	205

(2) INFORMATION FOR SEQ ID NO:177:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1119 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:177:

GCGCACTCGC AGCCCTGGCA GCGGGCACTG GTCATGGAAA ACGAATTGTT CTGCTCGGGC	60
GTCCTGGTGC ATCCGCAGTG GGTGCTGTCA GCCGCACACT GTTCCAGAA CTCCTACACC	120
ATCGGGCTGG GCCTGCACAG TCTTGAGGCC GAACAAGAGC CAGGGAGCCA CAGCTGCSAG	180
GCCAGCCTCT CCGTACGGCA CCCAGAGTAC AACAGACCCT TGCTCGCTAA CGACCTCATG	240
CTCATCAAGT TGGACGAATC CGTGTCGAG TCTGACACCA TCCGAGCAT CAGCATTGCT	300
TGCGAGTGCC CTACCGCGGG GAACTCTTGC CTCGTTTCTG GCTGGGGTCT GCTGGCGAAG	360
GATGCTGTGA TTGCCATCCA GTCCCAGACT GTGGGAGGCT GGGAGTGTGA GAACCTTTCC	420
CAACCCTGBC AGGTTTGTAC CATTTGGCA ACTTCCAGTG CAGGACGTC CTGCTGCATC	480

```

CTCACTGGGT GCTCACTACT GCTCACTGCA TCACCCGGAA CACTGTGATC AACTAGCCAG      540
CACCATAGTT CTCCGAAGTC AGACTATCAT GATTACTGTG TTGACTGIGC TGTCTATTGT      600
ACTAACCATG CCGATGTTTA GGTGAAATTA GCGTCACTTG GCCTCAACCA TCTTGGTATC      660
CAGTTATCCT CACTGAATTC AGATTTCCTC CTTCAAGTGC AGCCATTCCC ACATAATTTT      720
TGACCTACAG AGGTGAGGGA TCATATAGCT CTTCAAGGAT GCTGGTACTC CCCTCACAAA      780
TTCATTTCTC CTGTTGTAGT GAAAGGTGCG CCCTCTGGAG CCTCCCAGGG TGGGTGTGCA      840
GGTCACAATG ATGAATGTAT GATCGTGTTC CCATTACCCA AAGCCTTTAA ATCCCTCATG      900
CTCAGTACAC CAGGGCAGGT CTAGCATTTC TTCATTTAGT GTATGCTGTC CATTTCATGCA      960
ACCACCTCAG GACTCCIGGA TTCTCTGCCT AGTTGAGCTC CTGCATGCTG CCTCCTTGGG     1020
GAGGTGAGGG AGAGGGCCCA TGTTTCAATG GGATCTGTGC AGTTGTAACA CATTAGGTGC     1080
TTAATAAACA GAAGCTGTGA TGTAAAAAAA AAAAAAAA     1119

```

(2) INFORMATION FOR SEQ ID NO:178:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 164 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:178:

```

Met Glu Asn Glu Leu Phe Cys Ser Gly Val Leu Val His Pro Gln Trp
1           5           10           15

Val Leu Ser Ala Ala His Cys Phe Gln Asn Ser Tyr Thr Ile Gly Leu
20           25           30

Gly Leu His Ser Leu Glu Ala Asp Gln Glu Pro Gly Ser Gln Met Val
35           40           45

Glu Ala Ser Leu Ser Val Arg His Pro Glu Tyr Asn Arg Pro Leu Leu
50           55           60

Ala Asn Asp Leu Met Leu Ile Lys Leu Asp Glu Ser Val Ser Glu Ser
65           70           75           80

Asp Thr Ile Arg Ser Ile Ser Ile Ala Ser Gln Cys Pro Thr Ala Gly
85           90           95

Asn Ser Cys Leu Val Ser Gly Trp Gly Leu Leu Ala Asn Asp Ala Val
100          105          110

Ile Ala Ile Gln Ser Xaa Thr Val Gly Gly Trp Glu Cys Glu Lys Leu
115          120          125

Ser Gln Pro Trp Gln Gly Cys Thr Ile Ser Ala Thr Ser Ser Ala Arg
130          135          140

Thr Ser Cys Cys Ile Leu Thr Gly Cys Ser Leu Leu Leu Thr Ala Ser

```

145

150

155

160

Pro Gly Thr Leu

(2) INFORMATION FOR SEQ ID NO:179:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 250 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:179:

CTGGAGTGCC TTGGTGTTC AAGCCCCTGC AGGAAGCAGA ATGCACCTTC TGAGGCACCT	60
CCAGCTGCCC CCGGCGGGG GATGCGAGGC TCGGAGCACC CTTGCCCCGC TGTGATTGCT	120
GCCAGGCACT GTTCATCTCA GCTTTTCTGT CCCTTTGCTC CCGGCAAGCG CTTCTGCTGA	180
AAGTTTCATAT CTGGAGCCTG ATGTCTTAAC GAATAAAGGT CCCATGCTCC ACCCGAAAAA	240
AAAAAAAAAA	250

(2) INFORMATION FOR SEQ ID NO:180:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 202 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:180:

ACTAGTCCAG TGTGGTGGAA TTCCATTGTG TTGGGCCCCA CACAATGGCT ACCTTTAACA	60
TCACCCAGAC CCCGCCCCCTG CCCGTGCCCC ACGCTGCTGC TAACGACAGT ATGATGCTTA	120
CTCTGCTACT CGGAACTAT TTTTATGTAA TTAATGTATG CTTTCTTGTT TATAAATGCC	180
TGATTTAAAA AAAAAAAAAA AA	202

(2) INFORMATION FOR SEQ ID NO:181:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 558 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:181:

TCCYTTTGKT NAGGTTTKHG AGACAMCCCK AGACCTWAAN CTGTGTCACA GACTTCYNGG	60
AATGTTTAGG CAGTCTAGT AATTCYTCG TAATGATTCT GTTATTACTT TCCTNATTCT	120
TTATTCCTCT TTCTCTGAA GATTAATGAA GTTGAAAATT GAGGTGGATA AATACAAAAA	180
GGTAGTGTGA TAGTATAAGT ATCTAAGTGC AGATGAAAGT GTGTTATATA TATCCATTCA	240
AAATTATGCA AGTTAGTAAT TACTCAGGGT TAACTAAATT ACTTTAATAT GCTGTTGAAC	300
CTACTCTGTT CCTGGCTAG AAAAAATTAT AAACAGGACT TTGTTAGTTT GGGGAAGCCAA	360
ATTGATAATA TTCTATGTTT TAAAAGTTGG GCTATACATA AATTATTAAG AATATGGAW	420

TTTTATTCCC	AGGAATATGG	KGTTCAATTT	ATGAATATTA	CSCRGGATAG	AWGTWTGAGT	480
AAAAYCACTT	TTGGTWAATA	YGTWAATATG	TCMTAAATAA	ACAARKGCTTT	GACTTATTTT	540
CAAAAAAAAA	AAAAAAAA					558

(2) INFORMATION FOR SEQ ID NO:182:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 479 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:182:

ACAGGGWTTK	GRGGATGCTA	AGSCCCCRGA	RWTYGTTTGA	TCCAACCCCTG	GCTTWTTTTC	60
AGAGGGGAAA	ATGGGGCCTA	GAAGTTACAG	MSCATYTAGY	TGGTGCGMTG	GCACCCCTGG	120
CSTCACACAG	ASTCCCGAGT	AOCTGGGACT	ACAGGCACAC	AGTCACTGAA	GCAGGCCCTG	180
TTWGCAATTC	ACGTTGCCAC	CTCCAACCTA	AACATTCTTC	ATATGTGATG	TCCTTAGTCA	240
CTAAGGTAA	ACTTTCCAC	CCAGAAAAGG	CAACTTAGAT	AAAATCTTAG	AGTACTTCA	300
TACTMTCTA	AGTCCTCTTC	CAGCCTCACT	KKGAGTCCTM	CYTGGGGGTT	GATAGGAANT	360
NTCTCTGGC	TTTCTCAATA	AARTCTCTAT	YCATCTCATG	TTTAATTTGG	TACGCATARA	420
AWTGSTGARA	AAATTAAAT	GTTCTGGTTY	MACTTTAAAA	ARAAAAAAAA	AAAAAAAA	479

(2) INFORMATION FOR SEQ ID NO:183:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:183:

AGCGGGGAGC	AGAAGCTAAA	GCCAAAGCCC	AAGAAGAGTG	GCAGTGCCAG	CACTGGTGCC	60
AGTACCAGTA	CCAATAACAG	TGCCAGTGCC	AGTGCCAGCA	CCAGTGGTGG	CTTCAGTGCT	120
GGTGCCAGCC	TGACCGCCAC	TCTCACATTT	GGGCTCTTCG	CTGGCCTTGG	TGGAGCTGGT	180
GCCAGCACCA	GTGGCAGCTC	TGGTGCTGT	GGTTTCTCCT	ACAAGTGAGA	TTTTAGATAT	240
TGTTAATCCT	GCCAGTCTTT	CTCTTCAAGC	CAGGGTGCAT	CCTCAGAAAC	CTACTCAACA	300
CAGCACTCTA	GGCAGCCACT	ATCAATCAAT	TGAAGTTGAC	ACTCTGCATT	ARATCTATTT	360
GCCATTTCAA	AAAAAAAAAA	AAAA				384

(2) INFORMATION FOR SEQ ID NO:184:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 496 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:184:

ACCGAATTGG	GACCGCTGGC	TTATAAGCGA	TCATGTYINT	CCRGATATKAC	CTCAACGAGC	60
------------	------------	------------	------------	-------------	------------	----

120

AGGGGATCG	AGTGTATACG	CTGAAGAAAT	TTGACTTGAT	GGGACAACAG	ACCTGCTCAG	120
CCCATCTCTG	TCGCTTCTCG	CCAGATGACA	AATACTCTSG	ACACCGAATC	ACCATCAAGA	180
AACGCTTCAA	GGTGTCTCAT	AACCAGCAAC	CGCGGCTTBT	CTTCTGAGGG	TCCCTTAAAC	240
TGATGTCTTT	TCTGCCACCT	GTTACCCCTC	CGAGACTCG	TAACCAAACT	CTTCGGACTG	300
TGAGCCCTGA	TGCCTTTTTG	CCAGCCATAC	TCTTTGGCAT	CGAGTCTCTC	GTGGCGATTG	360
ATTATGCTTG	TGTGAGGCAA	TCATGCTGGC	ATCACCTATA	AAGGGAACAC	ATTTGACTTT	420
TTTTTCTCAT	ATTTTAAATT	ACTACMAGAW	TATTWMAGAW	WAAATGAWTT	GAAAACTST	480
TAAAAA	AAAAAA					496

(2) INFORMATION FOR SEQ ID NO:185:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:185:

GCTGGTAGCC	TATGGCGKGG	CCCACGGAGG	GGCTCCTGAG	CCCACGGRAC	AGTGAATTCC	60
CAAGTATCYT	GGCGSGCGTC	TTCTACCGTC	CCTACTTGCA	GATCTTCGGG	CAGATTCCCC	120
AGGAGGACAT	GGACGTGGCC	CTCATGGAGC	ACAGCAACTG	YTCGTGGGAG	CCCGGCTTCT	180
GGGCACACCC	TCCTGGGGCC	CAGGCGGGCA	CCTGCGTCTC	CCAGTATGCC	AACTGGCTGG	240
TGGTGCTGCT	CCTCGTCATC	TTCTGCTCTG	TGGCCAAACAT	CCTGCTGCTC	AACTTGCTCA	300
TTGCCATGTT	CAGTTACACA	TTCGGCAAAAG	TACAGGGCAA	CAGCGATCTC	TACTGGGAAG	360
GCGCAGCGTT	ACCGCCTCAT	CCGG				384

(2) INFORMATION FOR SEQ ID NO:186:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 577 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:186:

GAGTTAGCTC	CTCCACAACC	TTGATGAGGT	CGTCTGCACT	GCCCTCTCGC	TTCATACCGC	60
TNCCATCGTC	ATACTGTAGG	TTTGCCACCA	CYTCTGGCA	TCTTGGGGCG	GCNTAATATT	120
CCAGGAAACT	CTCAATCAAG	TCACCGTCGA	TGAAACCTGT	GGGCTGGTTC	TGTCTTCCGC	180
TCGGTGTGAA	AGGATCTCCC	AGAAGGAGTG	CTCGATCTTC	CCCACACTTT	TGATGACTTT	240
ATTGAGTCGA	TTCTGCATGT	CCAGCAGGAG	GTTGTACCAG	CTCTCTGACA	GTGAGGTCAC	300
CAGCCCTATC	ATGCCGTTGA	MCCTGCCGAA	GARCACCGAG	CCTTGTGTGG	GGGKKGAAGT	360
CTCACCAGAG	TTCTGCATTA	CCAGAGAGCC	GTGGCAAAAG	ACATTGACAA	ACTCGCCACG	420
GTGGAAAAAG	AMCAMCTCCT	GGARGTGCTN	GCCGCTCCTC	GTCMGTGGT	GGCAGCGCTW	480
TCCTTTTGAC	ACACAAACAA	GTTAAAGGCA	TTTTAGCCCC	CCAGAAANTT	GTCATCATCC	540
AAGATNTCGC	ACAGCACTNA	TCCAGTTGGG	ATTAAAT			577

(2) INFORMATION FOR SEQ ID NO:187:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 534 base pairs
- (B) TYPE: nucleic acid

121

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:187:

AACATCTTCC	TGTATAATGC	TGTGTAATAT	CGATCCGATN	TTGTCTGSTG	AGAATYCATW	60
ACTKGGAAAA	GMAACATTAA	AGCCTGGACA	CTGGTATTAA	AATTCACAAT	ATGCAACACT	120
TTAAACAGTG	TGTCAATCTG	CTCCCYYNAC	TTTGTTCATCA	CCAGTCTGGG	AAKAAGGGTA	180
TGCCCTATTG	ACACCTGTTA	AAAGGGCGCT	AAGCATTTTT	GATTCAACAT	CTTTTTTTTT	240
GACACAAGTC	CGAAAAAAGC	AAAAGTAAAC	AGTTATYAAT	TTGTTAGCCA	ATTCACCTTC	300
TTCATGGGAC	AGAGCCATYT	GATTTAAAAA	GCAAATTGCA	TAATATTGAG	CTTYGGGAGC	360
TGATATTTGA	GCGGAAGAGT	AGCCTTTCTA	CTTCACCAGA	CACAACTCCC	TTTCATATTG	420
GGATGTTNAC	NAAAGTWATG	TCTCTWACAG	ATGGGATGCT	TTTGTGGCAA	TTCTGTTCTG	480
AGGATCTCCC	AGTTTATTTA	CCACTTGCAC	AAGAAGGCGT	TTTCTTCCTC	AGGC	534

(2) INFORMATION FOR SEQ ID NO:188:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 761 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:188:

AGAAACCAGT	ATCTCTNAAA	ACAACCTCTC	ATACCTTGTG	GACCTAATTT	TGTGTGCGTG	60
TGTGTGTGCG	CGCATATTAT	ATAGACAGGC	ACATCTTTTT	TACTTTTGTA	AAAGCTTATG	120
CCTCTTTGGT	ATCTATATCT	GTGAAAGTTT	TAATGATCTG	CCATAATGTC	TTGGGGACCT	180
TTGTCTTCTG	TGTAAATGGT	ACTAGAGAAA	ACACCTATNT	TATGAGTCAA	TCTAGTTNGT	240
TTTATTGAC	ATGAAGGAAA	TTTCCAGATN	ACAACACTNA	CAAACCTCTC	CTKGACKARG	300
GGGGACAAAG	AAAAGCAAAA	CTGAMCATAA	RAACAATWA	CCTGGTGAGA	ARTTGCATAA	360
ACAGAAATWR	GGTAGTATAT	TGAARNACAC	CATCATTAAA	RMGTTWTKTT	WTTCTCCCTT	420
GCAAAAAACA	TGTACNGACT	TCCCGTTGAG	TAATGCCAAG	TTGTTTTTTT	TATNATAAAA	480
CTTGCCCTTC	ATTACATGTT	TNAAAGTGGT	GTGGTGGGCC	AAAATATTGA	AATGATGGAA	540
CTGACTGATA	AAGCTGTACA	AATAAGCAGT	GTGCCTAACA	AGCAACACAG	TAATGTTGAC	600
ATGCTTAATT	CACAAATGCT	AATTTCATT	TAAATGTTTG	CTAAAATACA	CTTTGAACCTA	660
TTTTTCTGTN	TTCCAGAGC	TGAGATNTTA	GATTTTATGT	AGTATNAAGT	GAAAAANTAC	720
GAAAATAATA	ACATTGAAGA	AAAANANAAA	AAANAAAAAA	A		761

(2) INFORMATION FOR SEQ ID NO:189:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 482 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:189:

TTTTTTTTTT	TTTGCCGATN	CTACTATTTT	ATTGCAGGAN	GTGGGGGTGT	ATGCACCGCA	60
CACCGGGGCT	ATNAGAAGCA	AGAAGGAAGG	AGGGAGGGCA	CAGCCCCTTG	CTGAGCAACA	120
AAGCCGCGCT	CTGCCTTCTC	TGTCTGTCTC	CTGGTGCAGG	CACATGGGGA	GACCTTCCCC	180

122

AAGGCAG3GG	CCACCAGTCC	AGGGGTG3GA	ATACAGG3GG	TGGGANGTGT	GCATAAGAAG	240
TGATAG3CAC	AGCCCACCCC	BTACAGACCC	CTCCGCTCCT	GACAG3TN3A	TTTCGACCAG	300
GTCATT3TGC	CCT3CCCCAG	CACAGCGTAN	ATCTGGAAAA	GACAGAAT3C	TTTCCTTTTC	360
AAATTT3GCT	NGT3ATNGAA	N3GGCANTTT	TCCAAATTNG	GCTNG3TCTT	GGTACNCTTG	420
GTTCCGG3CCA	GCT3CNC3TC	CAAAAANTAT	TCAC3CENNCT	CCNAATT3GT	TGCNGGNCCC	480
CC						482

(2) INFORMATION FOR SEQ ID NO:190:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 471 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:190:

TTTTTTTTTT	TTTTAAAACA	GTTTTTCACA	ACAAAATTTA	TTAGAAGAAT	AGTGGTTTTG	60
AAAACCTCTG	CATCCAGTGA	GAACCTACCAT	ACACCACATT	ACAGCTNG3A	ATGTNCTCCA	120
AATGTCTGGT	CAAATGATAC	AATGGAACCA	TTCAATCTTA	CACATGCA3G	AAAGAACAAG	180
CGCTTTTGGC	ATACAATGCA	CAAAAAAAAA	AGGGGGGGGG	GACCACAT3G	ATTTAAATTT	240
TAAGTACTCA	TCACATACAT	TAAGACACAG	TTCTAGTCCA	GTGNAAAATC	AGAACTGCNT	300
TGAAAAATTT	CATGTATGCA	ATCCAACCAA	AGAACTTNAT	TGGTGATCAT	GANTNCTCTA	360
CTACATCNAC	CTTGATCATT	GCCAGGAACN	AAAAGTTNAA	ANCACNCNGT	ACAAAAANAA	420
TCTGTAATTN	ANTTCAACCT	CCGTACNGAA	AAATNTTNTT	TATACACTCC	C	471

(2) INFORMATION FOR SEQ ID NO:191:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 402 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:191:

GAGGGATTGA	AGGTCTGTTC	TASTGTGCGM	CTGTTTCAGCC	ACCAACTCTA	ACAAGTTGCT	60
GTCTTCCACT	CACGTCTGT	AAGCTTTTTA	ACCCAGACWG	TATCTTCATA	AATAGAACAA	120
ATTCTTCACC	AGTCACATCT	TCTAGGACCT	TTTTGGATTG	AGTTAGTATA	AGCTCTTCCA	180
CTTCCTTTGT	TAAGACTTCA	TCTGGTAAAG	TCTTAAGTTT	TGTAGAAAGG	AATTYAATTG	240
CTCGTTCTCT	AACAATGTCC	TCTCCTTGAA	GTATTTGGCT	GAACAACCCA	CCTAAAGTCC	300
CTTTGTGCAT	CCATTTTAAA	TATACTTAAT	AGGGCATTGK	TNCACTAGGT	TAAATTCTGC	360
AAGAGTCATC	TGTCTGCAAA	AGTTGCGTTA	GTATATCTGC	CA		402

(2) INFORMATION FOR SEQ ID NO:192:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 601 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:192:

GAGCTCGGAT	CCAATAATCT	TTGTCTGAGG	GCAGCACACA	TATNCAGTGC	CATGGNAACT	60
GGTCTACCCC	ACATGGGAGC	AGCATGCCGT	AGNTATATAA	GGTCATTCCC	TGAGTCAGAC	120
ATGCTYTTTT	GAYTACCGTG	TGCCAAGTGC	TGGTGATTCT	YAACACACYT	CCATCCCGYT	180
CTTTTGTGGA	AAAACCTGGCA	CTTKTCTGGA	ACTAGCARGA	CATCACTTAC	AAATTCACCC	240
ACGAGACACT	TGAAAGGTGT	AACAAAGCGA	YTCTTGCAAT	GCTTTTTGTC	CCFCCGGCAC	300
CAGTTGTCAA	TACTAACCCG	CTGGTTTGCC	TCCATCACAT	TTGTGATCTG	TAGCTCTGGA	360
TACATCTCCT	GACAGTACTG	AAGAACTTCT	TCTTTTGTTC	CAAAAGCARG	TCTTGCTGCC	420
IGTTGGATCA	GCTTCCCATT	TCCCAGTCYG	AATGTTTACA	TGGCATATTT	WACTTCCCAC	480
AAAACATTGC	GATTTGAGGC	TCAGCAACAG	CAAATCCTGT	TCCGGCATTG	GCTGCAAGAG	540
CCTCGATGTA	GCCGGCCAGC	GCCAAGGCAG	GCGCCGTGAG	CCCCACCAGC	AGCAGAAGCA	600
G						601

(2) INFORMATION FOR SEQ ID NO:193:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 608 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:193:

ATACAGCCCA	NATCCCACCA	CGAAGATGCG	CTTGTTGACT	GAGAACCTGA	TGCGGTCACT	60
GGTCCCCTG	TAGCCCCAGC	GACTCTCCAC	CTGCTGGAAG	CGGTTGATGC	TGCACTCYTT	120
CCCAACGCAG	GCAGMAGCGG	GSCCGGTCAA	TGAACTCCAY	TCGTGGCTTG	GGGTKGACGJ	180
TKAAGTGCG	GAAGAGGCTG	ACCACCTCGC	GGTCCACCAG	GATGCCCCGAC	TGTGCGGGAC	240
CTGCAGCGAA	ACTCCTCGAT	GGTCATGAGC	GGGAAGCGAA	TGAGGCCCCAG	GGCCTTGCCC	300
AGAACCTTCC	GCCTGTTCTC	TGGCGTCACC	TGCAGCTGCT	GCCGCTGACA	CTCGGCCTCG	360
GACCAGCGGA	CAAACGGCRT	TGAACAGCCG	CACCTCACGG	ATGCCCAGTG	TGTCGCGCTC	420
CAGGAMMGSC	ACCAGCGTGT	CCAGGTCAAT	GTCGGTGAAG	CCCTCCGCGG	GTRATGGCGT	480
CTGCACTGTT	TTTGTCGATG	TTCTCCAGGC	ACAGGCTGGC	CAGCTGCGGT	TCATCGAAGA	540
GTCGCGCCTG	CGTGAGCAGC	ATGAAGGCGT	TGTCGGCTCG	CAGTTCTTCT	TCAGGAACTC	600
CACGCAAT						608

(2) INFORMATION FOR SEQ ID NO:194:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 392 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:194:

GAACGGCTGG	ACCTTGCCCTC	GCATTGTGCT	TGCTGGCAGG	GAATACCTTG	GCAAGCAGYT	60
CCAGTCCGAG	CAGCCCCAGA	CCGCTGCCGC	CCGAAGCTAA	GCCTGCCTCT	GGCCTTCCCC	120
TCCGCCTCAA	TGCAGAACCA	GTAGTGGGAG	CACTGTGTTT	AGAGTTAAGA	GTGAACACTG	180
TTTGATTTTA	CTTGGGAATT	TCCTCTGTTA	TATAGCTTTT	CCCAATGCTA	ATTTCCAAAC	240
AACAACAACA	AAATAACATG	TTTGCTGTG	AAGTTGTATA	AAAGTAGGTG	ATTCTGTATT	300
TAAAGAAAAAT	ATTACTGTTA	CATATACTGC	TTGCAATTTT	TGTATTTATT	GKTNCTSTGJ	360
AAATAAATAT	AGTTATTAAA	GCTTGTTCANT	CC			392

(2) INFORMATION FOR SEQ ID NO:195:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 502 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:195:

```

CCSTTKGAGG GGTKAGGKYC CAGTTYCCGA GTGGAAGAAA CAGGCCAGGA GAAGTGCGTG      60
CCGAGCTGAG GCAGATGTTT CCACAGTGAC CCCCAGAGCC STGGGSTATATA GTYTCTGACC      120
CCTCNCAAGG AAAGACCACS TTCTGGGGAC ATGGGCTGGA GGGCAGGACC TAGAGGCACC      180
AAGGGAAGGC CCCATTCCGG G3ST3TTCCG CGAGGAGGAA G3GAAGGGGGC TCTGTGTGCC      240
CCCCASGAGG AAGAGGCCCT GAGTCTGGG ATCAGACACC CCTTCACGTG TATCCCCACA      300
CAAATGCAAG CTCACCAAGG TCCCTCTCA GTCCCTTCC STACACCCTG AMCGGCCACT      360
GSCSCACACC CACCCAGAGG ACGCCACCCG CCATGGGGAR TGTGCTCAAG GARTCGCNGG      420
GCARCGTGGA CATCTNGTCC CAGAAGGGGG CAGAATCTCC AATAGANGGA CTGARCMSTT      480
GCTNANAAAA AAAAAANAAAA AA

```

(2) INFORMATION FOR SEQ ID NO:196:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 665 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:196:

```

GCTTACTTGG TTTCATTGCC ACCACTTAGT GGATGTCATT TAGAACCATT TTGTCTGCTC      60
CCTCTGGAAG CCTTGCGCAG AGCGGACTTT GTAATTGTTG GAGAATAACT GCTGAATTTT      120
WAGCTGTTTK GAGTTGATTS GCACCACTGC ACCCACAAC TCAATATGAA AACYAWTTGA      180
ACTWATTTAT TATCTTGTA AAAGTATAAC AATGAAAATT TTGTTCATAC TGTATTKATC      240
AAGTATGATG AAAAGCAAWA GATATATATT CTTTTATTAT GTTAAATTAT GATTGCCATT      300
ATTAATCGGC AAAATGTGGA GTGTATGTTT TTTTCACAGT AATATATGCC TTTTGTAAC T      360
TCACTTGGTT ATTTTATTGT AAATGARTTA CAAAATTCTT AATTTAAGAR AATGGTATGT      420
WATATTTATT TCATTAATTT CTTTCCTKGT TTACGTWAAT TTTGAAAAGA WTGCATGATT      480
TCTTGACAGA AATCGATCTT GATGCTGTGG AAGTAGTTTG ACCCACATCC CTATGAGTTT      540
TTCTTAGAAT GTATAAAGGT TGTAGCCCAT CNAACTTCAA AGAAAAAAT GACCACATAC      600
TTTGCAATCA GGCTGAAATG TGGCATGCTN TTCTAATTCC AACTTTATAA ACTAGCAAAN      660
AAGTG

```

(2) INFORMATION FOR SEQ ID NO:197:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 492 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:197:

TTTTNTTTTT	TTTTTTTTTGC	AGGAAGGATT	CCATTTATTG	TGGATGCATT	TTCACAATAT	60
ATGTTTATTG	GAGCGATCCA	TTATCAGTGA	AAAGTATCAA	GTGTTTATAA	NATTTTTAGG	120
AAGGCAGATT	CACAGAACAT	GCTNGTCNGC	TTGCAGTTTT	ACCTCGTANA	GATNACAGAG	180
AATTATAGTC	NAACCAGTAA	ACNAGGAATT	TACTTTTCAA	AAGATTAAAT	CCAAACTGAA	240
CAAAATTCTA	CCCTGAAACT	TACTCCATCC	AAATATTGGA	ATAANAGTCA	GCAGTGATAC	300
ATTCTCTTCT	GAACCTTAGA	TTTTCTAGAA	AAATATGTAA	TAGTGATCAG	GAAGAGCTCT	360
TGTTCAAAAG	TACAAACNAAG	CAATGTTCCC	TTACCATAGG	CCTTAATTCA	AACCTTGATC	420
CATTTCACTC	CCATCACGGG	AGTCAATGCT	ACCTGGGACA	CTTGTATTTT	GTTTCATNCTG	480
ANCNTGGCTT	AA					492

(2) INFORMATION FOR SEQ ID NO:198:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 478 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:198:

TTTNTTTTTGN	ATTTTCANTCT	GTANNAANTA	TTTTCATTAT	GTTTATTANA	AAAATATNAA	60
TGTNTCCACN	ACAAATCATN	TTACNTNACT	AAGAGGCCAN	CTACATTGTA	CAACATACAC	120
TGAGTATATT	TTGAAAAGGA	CAAGTTTAAA	GTANACNCAT	ATTGCCCGANC	ATANCACATT	180
TATACATGGC	TTGATTGATA	TTTAGCACAG	CANAAACTGA	GTGAGTTACC	AGAAAANAAT	240
NATATATGTC	AATCNGATTT	AAGATACAAA	ACAGATCCTA	TGGTACATAN	CATCNIGTAG	300
GAGTTGTGGC	TTTATGTTTA	CTGAAAGTCA	ATGCAGTTCC	TGTACAAAGA	GATGGCCGTA	360
AGCATTCTAG	TACCTCTACT	CCATGGTTAA	GAATCGTACA	CTTATGTTTA	CATATGTNCA	420
GGGTAAGAAT	TGTGTTAAGT	NAANTTATGG	AGAGGTCCAN	GAGAAAAAAT	TGATNCAA	478

(2) INFORMATION FOR SEQ ID NO:199:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 482 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:199:

AGTGACTTGT	CCTCCAACAA	AACCCCTTGA	TCAAGTTTGT	GGCACTGACA	ATCAGACCTA	60
TGCTAGTTCC	TGTCATCTAT	TCGCTACTAA	ATGCAGACTG	GAGGGGACCA	AAAAGGGGCA	120
TCAACTCCAG	CTGGATTATT	TTGGAGCCTG	CAAATCTATT	CCTACTTGTA	CGGACTTTGA	180
AGTGATTGAG	TTTCCTCTAC	GGATGAGAGA	CTGGCTCAAG	AATATCCTCA	TGCAGCTTTA	240
TGAAGCCNAC	TCTGAACACG	CTGGTTATCT	NAGATGAGAA	NCAGAGAAAT	AAAGTCNAGA	300
AAATTTACCT	GGANGAAAAG	AGGCTTTNGG	CTGGGGACCA	TCCCATTGAA	CCTTCTCTTA	360
ANGGACTTTA	AGAANAAACT	ACCACATGTN	TGTNGTATCC	TGGTGCCNGG	CCGTTTANTG	420
AACNTNGACN	NCACCCTTNT	GGAATANANT	CTTGACNGCN	TCCTGAACTT	GCTCCTCTGC	480
GA						482

(2) INFORMATION FOR SEQ ID NO:200:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 270 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:200:

CGGCCGCAAG	TGCAACTCCA	GCTGGGGCCG	TGCGGACGAA	GATTCTGCCA	GCAGTTGGTC	60
CGACTGCGAC	GACGGCGGCG	GCGACAGTCG	CAGGTGCAGC	GCGGGCGCCT	GGGGTCTTGC	120
AAGGCTGAGC	TGACGCCGCA	GAGGTGCTGT	CACGTCCAC	GACCTTGACG	CCGTGCGGGA	180
CAGCCGGAAC	AGAGCCCGGT	GAANGCGGGA	GGCCTCGGGG	AGCCCCCTCGG	GAAGGGCGGC	240
CCGAGAGATA	CGCAGGTGCA	GGTGGCCGCC				270

(2) INFORMATION FOR SEQ ID NO:201:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 419 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:201:

TTTTTTTTTT	TTTTGGAATC	TACTGCGAGC	ACAGCAGGTC	AGCAACAAGT	TTATTTTGCA	60
GCTAGCAAGG	TAACAGGGTA	GGGCATGGTT	ACATGTTTAC	GTCAACTTCC	TTTGTCGTCC	120
TTGATTGGTT	TGTCTTTATG	GGGGCGGGGT	GGGGTAGGGG	AAANCGAAGC	ANAANTAACA	180
TGGAGTGGGT	GCACCCTCCC	TGTAGAACCT	GGTTACNAAA	GCTTGGGGCA	GTTTACCTCC	240
TCTGTGACCG	TCATTTTCTT	GACATCAATG	TTATTAGAAG	TCAGGATATC	TTTTAGAGAG	300
TCCACTGTNT	CTGGAGGGAG	ATTAGGGTTT	CTTGCCAANA	TCCAANCAAA	ATCCACNTGA	360
AAAAGTTGGA	TGATNCANGT	ACNGAATACC	GANGGCATAN	TTCTCATANT	CGGTGGCCA	419

(2) INFORMATION FOR SEQ ID NO:202:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 509 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:202:

TTTNTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	60
TGGCACTTAA	TCCATTTTAA	TTTCAAAATG	TCTACAAANT	TTNAATNCNC	CATTATACNG	120
GTNATTTTNC	AAAATCTAAA	NNTTATTCAA	ATNTNAGCCA	AANTCCTTAC	NCAAATNNAA	180
TACNCCNAAA	AATCAAAAAT	ATACNTNTCT	TTCAGCAAAC	TTNGTTACAT	AAATTAATAA	240
AATATATACG	GCTGGTGTCT	TCAAAGTACA	ATTATCTTAA	CACTGCAAAC	ATNTTTNNAA	300
GGAACATAAA	TAAAAAATAA	CACTNCCGCA	AAGGTATAAG	GGAACAACAA	ATTCTTTTAA	360
CAACANNCNC	NATTATAAAA	ATCATATCTC	AAATCTTAGG	GGAATATATA	CTTCACACNG	420
GGATCTTAAC	TTTACTNCA	CTTTGTTTAT	TTTTTTANAA	CCATTGTNTT	GGCCCCAACA	480
CAATGCGAAT	NCCNCCNCNC	TGGACTAGT				509

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:197:

TTTTNTTTTT	TTTTTTTTTGC	AGGAAGGATT	CCATTTATTG	TGGATGCATT	TTCACAATAT	60
ATGTTTATTG	GAGCGATCCA	TTATCAGTGA	AAAGTATCAA	GTGTTTATAA	NATTTTTTAGG	120
AAGGCAGATT	CACAGAACAT	GCTNGTCNGC	TTGCAGTTTT	ACCTCGTANA	GATNACAGAG	180
AATTATACTC	NAACCAGTAA	ACNAGGAATT	TACTTTTCAA	AAGATTAAAT	CCAAAC'TGAA	240
CAAAATTCTA	CCCTGAAACT	TACTCCATCC	AAATATTGGA	ATAANAGTCA	GCAGTGATAC	300
ATTCTCTTCT	GAAC'TT'TAGA	TTTTCTAGAA	AAATATGTAA	TAGTGATCAG	GAAGAGCTCT	360
TGTTCAAAAG	TACAACNAAG	CAATGTTCCC	TTACCATAGG	CCTTAATTCA	AAC'TTTGATC	420
CATTTCACTC	CCATCACGGG	AGTCAATGCT	ACCTGGGACA	CTTGTATTTT	GTTCA'TNCTG	480
ANCNTGGCTT	AA					492

(2) INFORMATION FOR SEQ ID NO:198:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 478 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:198:

TTTNTTTTTGN	ATTTCA'NTCT	GTANNAANTA	TTTTCATTAT	GTTTATTANA	AAAATATNAA	60
TGTNTCCACN	ACAAATCATN	TTACNTNAGT	AAGAGGCCAN	CTACATTGTA	CAACATACAC	120
TGAGTATATT	TTGAAAAGGA	CAAGTTTAAA	GTANACNCAT	ATTGCCGANC	ATANCACATT	180
TATACATGGC	TTGATTGATA	TTTAGCACAG	CANAAACTGA	GTGAGTTACC	AGAAAANAAT	240
NATATATGTC	AATCNGATTT	AAGATACAAA	ACAGATCCTA	TGGTACATAN	CATCNTGTAG	300
GAGTTGTGGC	TTTATGTTTA	CTGAAAGTCA	ATGCAGTTCC	TGTACAAAGA	GATGGCCCGTA	360
AGCATTCTAG	TACCTCTACT	CCATGGTTAA	GAATCGTACA	CTTATGTTTA	CATATGTNCA	420
GGGTAAGAAT	TGTGTTAAGT	NAANTTATGG	AGAGGTCCAN	GAGAAAAATT	TGATNCAA	478

(2) INFORMATION FOR SEQ ID NO:199:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 482 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:199:

AGTGACTTGT	CCTCCAACAA	AACCCCTTGA	TCAAGTTTGT	GGCACTGACA	ATCAGACCTA	60
IGCTAGTTCC	TGTCATCTAT	TCGCTACTAA	ATGCAGACTG	GAGGGGACCA	AAAAGGGGCA	120
TCAACTCCAG	CTGGATTATT	TTGGAGCCTG	CAAATCTATT	CCTACTTGTA	CGGACTTTGA	180
AGTGATTGAG	TTTCCTCTAC	GGATGAGAGA	CTGGCTCAAG	AATATCCTCA	TGCAGCTTTA	240
TGAAGCCNAC	TCTGAACACG	CTGGTTATCT	NAGATGAGAA	NCAGAGAAAT	AAAGTCNAGA	300
AAATTTACCT	GGANGAAAAG	AGGCTTTNGG	CTGGGGACCA	TCCCATTGAA	CCTTCTCTTA	360
ANGGACTTTA	AGAANA'AACT	ACCACATGTN	TGTNGTATCC	TGGTGCCNGG	CCGTTTANTG	420
AACNTNGACN	NCACCCTTNT	GGAATANANT	CTTGACNGCN	TCCTGAACTT	GCTCCTCTGC	480
GA						482

(2) INFORMATION FOR SEQ ID NO:200:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 270 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:200:

CGGCCGCAAG	TGCAACTCCA	GCTGGGGCCG	TGCGGACGAA	GATTCTGCCA	GCAGTTGGTC	60
CGACTGCGAC	GACGGCGGCG	GCGACAGTCG	CAGGTGCAGC	GCGGGCGCCT	GGGGTCTTGC	120
AAGGCTGAGC	TGACGCCGCA	GAGGTCTGT	CACGTCCCAC	GACCTTGACG	CCGTCGGGGA	180
CAGCCGGAAC	AGAGCCCGGT	GAANGCGGGA	GGCCTCGGGG	AGCCCCCTCG	GAAGGGCGGC	240
CCGAGAGATA	CGCAGGTGCA	GGTGGCCGCC				270

(2) INFORMATION FOR SEQ ID NO:201:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 419 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:201:

TTTTTTTTTT	TTTTGGAATC	TACTGCGAGC	ACAGCAGGTC	AGCAACAAGT	TTATTTTGCA	60
GCTAGCAAGG	TAACAGGGTA	GGGCATGGTT	ACATGTTTCC	GTCAACTTCC	TTTGTCGTGG	120
TTGATTGGTT	TGTCTTTATG	GGGGCGGGGT	GGGGTAGGGG	AAANCGAAGC	ANAANTAACA	180
TGGAGTGGGT	GCACCCTCCC	TGTAGAACCT	GGTTACNAAA	GCTTGGGGCA	GTTTACCTGG	240
TCTGTGACCG	TCATTTTCTT	GACATCAATG	TTATTAGAAG	TCAGGATATC	TTTTAGAGAG	300
TCCACTGTNT	CTGGAGGGAG	ATTAGGTTT	CTTGCCAANA	TCCAANCAAA	ATCCACNTGA	360
AAAAGTTGGA	TGATNCANGT	ACNGAATACC	GANGGCATAN	TTCTCATANT	CGGTGGCCA	419

(2) INFORMATION FOR SEQ ID NO:202:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 509 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:202:

TTTNTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	60
TGGCACTTAA	TCCATTTTTA	TTTCAAAATG	TCTACAAANT	TTNAATNCNC	CATTATACNG	120
GTNATTTTNC	AAAATCTAAA	NNTTATTCAA	ATNTNAGCCA	AANTCCTTAC	NCAAATNNAA	180
TACNCNCAAA	AATCAAAAAT	ATACNTNTCT	TTCAGCAAAC	TTNGTTACAT	AAATTAATAAA	240
AATATATACG	GCTGGTGTTT	TCAAAGTACA	ATTATCTTAA	CACTGCAAAC	ATNTTTNNAA	300
GGAACATAAA	TAAAAAATAA	CACTNCCGCA	AAGGTTAAAG	GGAACAACAA	ATTCTTTTAA	360
CAACANCNNC	NATTATAAAA	ATCATATCTC	AAATCTTAGG	GGAATATATA	CTTCACACNG	420
GGATCTTAAC	TTTTACTNCA	CTTTGTTTAT	TTTTTTANAA	CCATTGTNTT	GGGCCCCAACA	480
CAATGGNAAT	NCCNCCNCNC	TGGACTAGT				509

(2) INFORMATION FOR SEQ ID NO:203:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 583 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:203:

TTTTTTTTTT	TTTTTTTTTGA	CCCCCTCTT	ATAAAAAACA	AGTTACCATT	TTATTTTACT	60
TACACATATT	TATTTTATAA	TTGGTATTAG	ATATTCAAAA	GGCAGCTTTT	AAAATCAAAC	120
TAAATGGAAA	CTGCCTTAGA	TACATAATTC	TTAGGAATTA	GCTTAAAATC	TGCCTAAAGT	180
GAAAATCTTC	TCTAGCTCTT	TTGACTGTAA	ATTTTGTACT	CTTGTA AAC	ATCCAAATTC	240
ATTTTCTTG	TCTTTAAAT	TATCTAATCT	TTCCATTTT	TCCCTATTCC	AAGTCAATTT	300
GCTTCTCTAG	CCTCATTTCC	TAGCTCTTAT	CTACTATTAG	TAAGTGGCTT	TTTCCTAAA	360
AGGGAAAACA	GGAAGAGANA	ATGGCACACA	AAACAAACAT	TTATATTCA	TATTTCTACC	420
TACGTTAATA	AAATAGCATT	TTGTGAAGCC	AGCTCAAAAG	AAGGCTTAGA	TCCTTTTATG	480
TCCATTTTAG	TCACTAAACG	ATATCNAAAG	TGCCAGAATG	CAAAAGGTTT	GTGAACATTT	540
ATTCAAAAGC	TAATATAAGA	TATTTACAT	ACTCATCTTT	CTG		583

(2) INFORMATION FOR SEQ ID NO:204:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 589 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:204:

TTTTTTTTNT	TTTTTTTTTT	TTTTTNTCTC	TTCTTTTTTT	TTGANAATGA	GGATCGAGTT	60
TTTCACTCTC	TAGATAGGGC	ATGAAGAAAA	CTCATCTTTC	CAGCTTTTAA	ATAACAATCA	120
AATCTCTTAT	GCTATATCAT	ATTTTAAGTT	AAACTAATGA	GTCAGTGGCT	TATCTTCTCC	180
TGAAGGAAAT	CTGTTTCATTC	TTCTCATTCA	TATAGTTATA	TCAAGTACTA	CCTTGCATAT	240
TGAGAGGTTT	TTCTTCTCTA	TTTACACATA	TATTTCCATG	TGAATTTGTA	TCAAACCTTT	300
ATTTTCATGC	AAACTAGAAA	ATAATGTNTT	CTTTTGCATA	AGAGAAGAGA	ACAATATNAG	360
CATTACAAAA	CTGCTCAAAT	TGTTTGTTAA	GNTTATCCAT	TATAATTAGT	TNGGCAGGAG	420
CTAATACAAA	TCACATTTAC	NGACNAGCAA	TAATAAAACT	GAAGTACCAG	TTAAATATCC	480
AAAATAATTA	AAGGAACATT	TTTAGCCTGG	GTATAATTAG	CTAATTCAT	TTACAAGCAT	540
TTATTNAGAA	TGAATTCACA	TGTTATTATT	CCNTAGCCCA	ACACAATGG		589

(2) INFORMATION FOR SEQ ID NO:205:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 545 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:205:

128

```

TTTTTNTTTT TTTTTTCACT AATAATCAGA ACAATATTTA TTTTATATT TAAAATTCAT      60
AGAAAAGTGC CTTACATTTA ATAAAAGTTT GTTCTCTCAA GTGATCAGAG GAATTAGATA      120
TNGTCTTGAA CACCAATATT AATTTGAGGA AAATACACCA AAATACATTA AGTAAATTAT      180
TTAAGATCAT AGAGCTTGTA AGTGAAAAGA TAAAATTTGA CCTCAGAAAC TCTGAGCATT      240
AAAATCCAC TATTAGCAAA TAAATTACTA TGGACTTCTT GCTTTAATTT TGTGATGAAT      300
ATGGGGTGTG ACTGGTAAAC CAACACATTC TGAAGGATAC ATTACTTAGT GATAGATTCT      360
TATGTACTTT GCTANATNAC GTGGATATGA GTTGACAAGT TTCTCTTTCT TCAATCTTTT      420
AAGGGGCGNGA NGAAATGAGG AAGAAAAGAA AAGGATTACG CATACTGTTT TTTCTATNGG      480
AAGGATTAGA TATGTTTCTT TTGCCAATAT TAAAAAATA ATAATGTTTA CTACTAGTGA      540
AACC

```

(2) INFORMATION FOR SEQ ID NO:206:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 487 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:206:

```

TTTTTTTTTT TTTTTTAGTC AAGTTTCTNA TTTTATTAT AATTAAAGTC TTGGTCATTT      60
CATTTATTAG CTCTGCAACT TACATATTTA AATTAAAGAA ACGTTNNTAG ACAACTCTNA      120
CAATTTATAA ATGTAAGGTG CCATTATTGA GTANATATAT TCCTCCAAGA GTGGATGTGT      180
CCCTTCTCCC ACCAACTAAT GAANCAGCAA CATTAGTTTA ATTTTATTAG TAGATNATAC      240
ACTGCTGCAA ACGCTAATTC TCTTCTCCAT CCCCATGTNG ATATTGTGTA TATGTGTGAG      300
TTGGTNAGAA TGCATCANCA ATCTNACAAT CAACAGCAAG ATGAAGCTAG GCNTGGGCTT      360
TCGGTGAAAA TAGACTGTGT CTGTCTGAAT CAAATGATCT GACCTATCCT CGGTGGCAAG      420
AACTCTTCGA ACCGCTTCCT CAAAGGCNGC TGCCACATTT GTGGCNTCTN TTGCACTTGT      480
TTCAAAA

```

(2) INFORMATION FOR SEQ ID NO:207:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 332 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:207:

```

TGAATTGGCT AAAAGACTGC ATTTTTANAA CTAGCAACTC TTATTTCTTT CTTTTAAAAA      60
TACATAGCAT TAAATCCCAA ATCCTATTTA AAGACCTGAC AGCTTGAGAA GGCTACTACT      120
GCATTTATAG GACCTTCTGG TGGTTCTGCT GTTACNTTTC AANTCTGACA ATCCTTGANA      180
ATCTTTGCAT GCAGAGGAGG TAAAAGGTAT TGGATTTTCA CAGAGGAANA ACACAGCGCA      240
GAAATGAAGG GGCCAGGCTT ACTGAGCTTG TCCACTGGAG GGCTCATGGG TGGGACATGG      300
AAAAGAAGGC AGCCTAGGCC CTGGGGAGCC CA

```

(2) INFORMATION FOR SEQ ID NO:208:

(i) SEQUENCE CHARACTERISTICS:

129

- (A) LENGTH: 524 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:208:

```

AGGGCGTGGT GCGGAGGGCG TTA CTGTTT GTCTCAGTAA CAATAAATAC AAAAAGACTG      60
GTTGTGTTCC GGCCCCATCC AACCACGAAG TTGATTTCTC TTGTGTGCAG AGTGACTGAT      120
TTTAAAGGAC ATGGAGCTTG TCACAATGTC ACAATGTCAC AGTGTGAAGG GCACACTCAC      180
TCCCGCGTGA TTCACATTTA GCAACCAACA ATAGTCATG AGTCCATACT TGTAAATACT      240
TTTGGCAGAA TACTTNTTGA AACTTGCAGA TGATAACTAA GATCCAAGAT ATTTCCCAAA      300
GTAAATAGAA GTGGGTCATA ATATTAATTA CCTGTTTACA TCAGCTTCCA TTTACAAGTC      360
ATGAGCCCAG AACTGACAT CAACTAAGC CCACTTAGAC TCCTCACCAC CAGTCTGTCC      420
TGTCATCAGA CAGGAGGCTG TCACCTTGAC CAAATTCTCA CCAGTCAATC ATCTATCCAA      480
AAACCATTAC CTGATCCACT TCCGTAATG CACCACCTTG GTGA                          524

```

(2) INFORMATION FOR SEQ ID NO:209:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 159 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:209:

```

GGGTGAGGAA ATCCAGAGTT GCCATGGAGA AAATTCCAGT GTCAGCATTC TTGCTCCTTG      60
TGGCCCTCTC CTACACTCTG GCCAGAGATA CCACAGTCAA ACCTGGAGCC AAAAAGGACA      120
CAAAGGACTC TCGACCCAAA CTGCCCCAGA CCCTCTCCA                          159

```

(2) INFORMATION FOR SEQ ID NO:210:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 256 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:210:

```

ACTCCCTGGC AGACAAAGGC AGAGGAGAGA GCTCTGTTAG TTCTGTGTTG TTGAACTGCC      60
ACTGAATTTT TTTCCACTTG GACTATTACA TGCCANTTGA GGGACTAATG GAAAAACGTA      120
TGGGGAGATT TTANCCAATT TANGTNTGTA AATGGGGAGA CTGGGGCAGG CGGGAGAGAT      180
TTGCAGGGTG NAAATGGGAN GGCTGGTTTG TTANATGAAC AGGGACATAG GAGGTAGGCA      240
CCAGGATGCT AAATCA                          256

```

(2) INFORMATION FOR SEQ ID NO:211:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 264 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:211:

ACATTGTTTT	TTTGAGATAA	AGCATTGAGA	GAGCTCTCCT	TAACGTGACA	CAATGGAAGG	60
ACTGGAACAC	ATACCCACAT	CTTTGTTCTG	AGGGATAATT	TTCTGATAAA	GTCTTGCTGT	120
ATATTCAAGC	ACATATGTTA	TATATTATTC	AGTTCCATGT	TTATAGCCTA	GTTAAGGAGA	180
GGGGAGATAC	ATTCNGAAAG	AGGACTGAAA	GAAATACTCA	AGTNGGAAAA	CAGAAAAAGA	240
AAAAAAGGAG	CAAATGAGAA	GCCT				264

(2) INFORMATION FOR SEQ ID NO:212:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 328 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:212:

ACCCAAAAAT	CCAATGCTGA	ATATTTGGCT	TCATTATTC	CANATTCTTT	GATTGTCAAA	60
GGATTTAATG	TTGTCCTCAG	TTGGGCACTT	CAGTTAGGAC	CTAAGGATGC	CAGCCGGCAG	120
GTTTATATAT	GCAGCAACAA	TATTCAAGCG	CGACAACAGG	TTATTGAACT	TGCCCCGCCAG	180
TTNAATTTTCA	TTCCCATTTGA	CTTGGGATCC	TTATCATCAG	CCAGAGAGAT	TGAAAATTTA	240
CCCCTACNAC	TCTTTACTCT	CTGGANAGGG	CCAGTGGTGG	TAGCTATAAG	CTTGGCCACA	300
TTTTTTTTTTC	CTTTATTCCT	TTGTCAGA				328

(2) INFORMATION FOR SEQ ID NO:213:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 250 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:213:

ACTTATGAGC	AGAGCGACAT	ATCCNAGTGT	AGACTGAATA	AAACTGAATT	CTCTCCAGTT	60
TAAAGCATTG	CTCACTGAAG	GGATAGAAGT	GACTGCCAGG	AGGGAAAAGT	AGCCAAGGCT	120
CATTATGCCA	AAGGANATAT	ACATTTCAAT	TCTCCAAACT	TCTTCCTCAT	TCCAAGAGTT	180
TTCAATATTT	GCATGAACCT	GCTGATAANC	CATGTTAANA	AACAAATATC	TCTCTNACCT	240
TCTCATCGGT						250

(2) INFORMATION FOR SEQ ID NO:214:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 444 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:214:

```

ACCCAGAATC CAATGCTGAA TATTTGGCTT CATTATTCCC AGATTCTTTG ATTGTCAAAG      60
GATTTAATGT TGTCTCAGCT TGGGCACTTC AGTTAGGACC TAAGGATGCC AGCCGGCAGG      120
TTTATATATG CAGCAACAAT ATTCAAGCGC GACAACAGGT TATTGAACTT GCCCGCCAGT      180
TGAATTTTCAT TCCCATTGAC TTGGGATCCT TATCATCAGC CANAGAGATT GAAAATTTAC      240
CCCTACGACT CTTTACTCTC TGGAGAGGGC CAGTGGTGGT AGCTATAAGC TTGGCCACAT      300
TTTTTTTTCC TTTATTCCTT TGTACAGAT GCGATTCATC CATATGCTAN AAACCAACAG      360
AGTGACTTTT ACAAAATTCC TATAGANATT GTGAATAAAA CCTTACCTAT AGTTGCCATT      420
ACTTTGCTCT CCCTAATATA CCTC                                     444

```

(2) INFORMATION FOR SEQ ID NO:215:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 366 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:215:

```

ACTTATGAGC AGAGCGACAT ATCCAAGTGT ANACTGAATA AACTGAATT CTCTCCAGTT      60
TAAAGCATTG CTCACTGAAG GGATAGAAGT GACTGCCAGG AGGGAAAGTA AGCCAAGGCT      120
CATTATGCCA AAGGANATAT ACATTTCAAT TCTCCAACT TCTTCCTCAT TCCAAGAGTT      180
TTCAATATTT GCATGAACCT GCTGATAAGC CATGTTGAGA AACAAATATC TCTCTGACCT      240
TCTCATCGGT AAGCAGAGGC TGTAGGCAAC ATGGACCATA GCGAANAAAA AACTTAGTAA      300
TCCAAGCTGT TTTCTACACT GTAACCAGGT TTCCAACCA GGTGGAAATC TCCTATACTT      360
GGTGCC                                     366

```

(2) INFORMATION FOR SEQ ID NO:216:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 260 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:216:

```

CTGTATAAAC AGAACTCCAC TGCANGAGGG AGGGCCGGGC CAGGAGAATC TCCGCTTGTC      60
CAAGACAGGG GCCTAAGGAG GGTCTCCACA CTGCTNNTAA GGGCTNTNC ATTTTTTAT      120
TAATAAAAAG TNNAAAAGGC CTCTTCTCAA CTTTTTCCC TTNGGCTGGA AAATTTAAAA      180
ATCAAAAATT TCCTNAAGTT NTCAAGCTAT CATATATACT NTATCCTGAA AAAGCAACAT      240

```

AATTCTTCCT TCCCTCCTTT

262

(2) INFORMATION FOR SEQ ID NO:217:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 262 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:217:

ACCTACGTGG	GTAAGTTTAN	AAATGTTATA	ATTTTCAGGAA	NAGGAACGCA	TATAATTGTA	60
TCTTCCCTAT	AATTTTCTAT	TTTAATAAGG	AAATAGCAAA	TTGGCGCTCGG	GGGAATCTAG	120
GGCATTCTAC	AGTTTGAGCA	AAATGCAATT	AAATGTGGAA	GCACAGCACT	GAAAAATTTT	180
ATGAATAATC	TGTATGATTA	TATGTCTCTA	GAGTAGATTT	ATAATTAGCC	ACTTACCCTA	240
ATATCCTTCA	TGCTTGTAAG	GT				262

(2) INFORMATION FOR SEQ ID NO:218:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 205 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:218:

ACCAAGGTGG	TGCATTACCG	GAANTGGATC	AANGACACCA	TCGTGGCCAA	CCCCTGAGCA	60
CCCCTATCAA	CTCCCTTTTG	TAGTAAACTT	GGAACCTTGG	AAATGACCAG	GCCAAGACTC	120
AGGCCTCCCC	AGTTCTACTG	ACCTTTGTCC	TTANGTNTNA	NGTCCAGGGT	TGCTAGGAAA	180
ANAAATCAGC	AGACACAGGT	GTAAA				205

(2) INFORMATION FOR SEQ ID NO:219:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 114 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:219:

TACTGTTTTG	TCTCAGTAAC	AATAAATACA	AAAAGACTGG	TTGTGTTCGG	GCCCCATCCA	60
ACCACGAAGT	TGATTTCTCT	TGTGTGCAGA	GTGACTGATT	TTAAAGGACA	TGGA	114

(2) INFORMATION FOR SEQ ID NO:220:

(i) SEQUENCE CHARACTERISTICS:

133

- (A) LENGTH: 93 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:220:

ACTAGCCAGC ACAAAGGCA GGGTAGCCTG AATTGCTTTC TGCTCTTTAC ATTTCTTTTA	60
AAATAAGCAT TTAGTGCTCA GTCCCTACTG AGT	93

(2) INFORMATION FOR SEQ ID NO:221:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 167 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:221:

ACTANGTGCA GGTGCGCACA AATATTTGTC GATATTCCTT TCATCTTGGA TTCCATCAGC	60
TCTTTTGCCC AGCCTGTGGC TCTACTGTAG TAAGTTTCTG CTGATGAGGA GCCAGNATGC	120
CCCCCACTAC CTTCCCTGAC GTCGCCANA AATCACCCAA CCTCTGT	167

(2) INFORMATION FOR SEQ ID NO:222:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 351 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:222:

AGGGCGTGGT GCGGAGGGCG GTACTGACCT CATTAGTAGG AGGATGCATT CTGGCACCCC	60
GTTCTTCACC TGTCCCCCAA TCCTTAAAAG GCCATACTGC ATAAAGTCAA CAACAGATAA	120
ATGTTTGCTG AATTAAAGGA TGGATGAAAA AAATTAATAA TGAATTTTGG CATAATCCAA	180
TTTTCTCTTT TATATTTCTA GAAGAAGTTT CTTTGAGCCT ATTAGATCCC GGGAATCTTT	240
TAGGTGAGCA TGATTAGAGA GCTTGTAGGT TGCTTTTACA TATATCTGGC ATATTTGAGT	300
CTCGTATCAA AACAATAGAT TGGTAAAGGT GGTATTATTG TATTGATAAG T	351

(2) INFORMATION FOR SEQ ID NO:223

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:223:

AAAACAAACA AACAAAAAAA ACAATTCTTC ATTCAGAAAA ATTATCTTAS	GGACTGATAT	60
TGGTAATTAT GGTCAATTTA ATWRTRTTKT GGGGCATTTT	CTTACATTGT CTTGACAAGA	120
TTAAAAATGTC TGTGCCAAAA TTTTGTAATTT TATTGGGAGA	CTTCTTATCA AAAGTAATGC	180
TGCCAAAGGA AGTCTAAGGA ATTAGTAGTG TTCCCMTCAC	TTGTTTGGAG TGTGCTATTC	240
TAAAAGATTT TGATTTCCCTG GAATGACAAT TATATTTTAA	CTTTGGTGGG GGAAANAGTT	300
ATAGGACCAC AGTCTTCACT TCTGATACTT GTAAATTAAT	CTTTTATTGC ACTTGTTTTC	360
ACCATTAAGC TATATGTTTA AAA		383

(2) INFORMATION FOR SEQ ID NO:224

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 320 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:224

CCCCTGAAGG CTTCTTGTTA GAAAATAGTA CAGTTACAAC	CAATAGGAAC AACAAAAAGA	60
AAAAGTTTGT GACATTGTAG TAGGGAGTGT GTACCCCTTA	CTCCCCATCA AAAAAAAAAT	120
GGATACATGG TTAAAGGATA RAAGGGCAAT ATTTTATCAT	ATGTTCTAAA AGAGAAGGAA	180
GAGAAAATAC TACTTTCTCR AAATGGAAGC CCTTAAAGGT	GCTTTGATAC TGAAGGACAC	240
AAATGTGGCC GTCCATCCTC CTTTARAGTT GCATGACTTG	GACACGGTAA CTGTTGCAGT	300
TTTARACTCM GCATTGTGAC		320

CLAIMS

1. A method for detecting prostate cancer in a patient, comprising:
 - (a) contacting a biological sample obtained from the patient with a binding agent which is capable of binding to a polypeptide, the polypeptide comprising an immunogenic portion of a prostate protein or a variant thereof, wherein said protein comprises an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences and variants of said nucleotide sequences; and
 - (b) detecting in the sample a protein or polypeptide that binds to the binding agent, thereby detecting prostate cancer in the patient.
2. The method of claim 1 wherein the binding agent is a monoclonal antibody.
3. The method of claim 2 wherein the binding agent is a polyclonal antibody.
4. A method for monitoring the progression of prostate cancer in a patient, comprising:
 - (a) contacting a biological sample obtained from the patient with a binding agent that is capable of binding to a polypeptide, said polypeptide comprising an immunogenic portion of a prostate protein or a variant thereof, wherein said protein comprises an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences and variants of said nucleotide sequences;
 - (b) determining in the sample an amount of a protein or polypeptide that binds to the binding agent;
 - (c) repeating steps (a) and (b); and

(d) comparing the amount of polypeptide detected in steps (b) and (c) to monitor the progression of prostate cancer in the patient.

5. A monoclonal antibody that binds to a polypeptide comprising an immunogenic portion of a prostate protein or a variant thereof, wherein said protein comprises an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 8-29, 41-45, 47-52, 54-65, 70, 73, 74, 79, 81, 87, 90, 92, 93, 97, 103, 104, 107, 109-111, 115-160, 171, 173-175, 177, 181, 188, 191, 193, 194, 198, 203, 204, 207, 209-211, 220, 222-224, the complements of said nucleotide sequences variants of said nucleotide sequences.

6. A method for inhibiting the development of prostate cancer in a patient, comprising administering to the patient a therapeutically effective amount of a monoclonal antibody according to claim 5.

7. The method of claim 6 wherein the monoclonal antibody is conjugated to a therapeutic agent.

8. A method for detecting prostate cancer in a patient comprising:

- (a) obtaining a biological sample from the patient;
- (b) contacting the sample with at least two oligonucleotide primers in a polymerase chain reaction, wherein at least one of the oligonucleotides is specific for a DNA molecule encoding a polypeptide comprising an immunogenic portion of a prostate protein or of a variant thereof, said protein comprising an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences variants of said nucleotide sequences; and
- (c) detecting in the sample a DNA sequence that amplifies in the presence of the oligonucleotide primers, thereby detecting prostate cancer.

9. The method of claim 8, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA molecule having a sequence selected from SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224.

10. A diagnostic kit comprising:

- (a) one or more monoclonal antibodies of claim 5; and
- (b) a detection reagent.

11. A diagnostic kit comprising:

- (a) one or more monoclonal antibodies that bind to a polypeptide encoded by a DNA molecule having a nucleotide sequence selected from the group consisting of SEQ ID Nos: 5-7, 30-40, 46, 53, 66-69, 71, 72, 75-78, 80, 82-86, 88, 89, 91, 94-96, 98-102, 105, 106, 161-170, 179, 180, 182-187, 189, 190, 192, 195-197, 199-202, 205, 206, 208, 212-219, 221, the complements of said sequences and variants of said nucleotide sequences; and
- (b) a detection reagent.

12. The kit of claims 10 or 11 wherein the monoclonal antibodies are immobilized on a solid support.

13. The kit of claim 12 wherein the solid support comprises nitrocellulose, latex or a plastic material.

14. The kit of claims 10 or 11 wherein the detection reagent comprises a reporter group conjugated to a binding agent.

15. The kit of claim 14 wherein the binding agent is selected from the group consisting of anti-immunoglobulins, Protein G, Protein A and lectins.

16. The kit of claim 14 wherein the reporter group is selected from the group consisting of radioisotopes, fluorescent groups, luminescent groups, enzymes, biotin and dye particles.

17. A diagnostic kit comprising at least two oligonucleotide primers, at least one of the oligonucleotide primers being specific for a DNA molecule encoding a polypeptide comprising an immunogenic portion of a prostate protein or a variant thereof, said protein comprising an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences and variants of said nucleotide sequences.

18. A diagnostic kit of claim 17 wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA molecule having a sequence selected from SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224.

19. A method for detecting prostate cancer in a patient, comprising:


- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with an oligonucleotide probe specific for a DNA molecule encoding a polypeptide comprising an immunogenic portion of a prostate protein or a variant thereof, said protein comprising an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences and variants of said nucleotide sequences; and
- (c) detecting in the sample a DNA sequence that hybridizes to the oligonucleotide probe, thereby detecting prostate cancer in the patient.

20. The method of claim 19 wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA molecule having a sequence selected from the group consisting of SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224.

21. A diagnostic kit comprising an oligonucleotide probe specific for a DNA molecule encoding a polypeptide comprising an immunogenic portion of a prostate

protein or a variant thereof, said protein comprising an amino acid sequence encoded by a DNA molecule having a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224, the complements of said nucleotide sequences variants of said nucleotide sequences.

22. The diagnostic kit of claim 21, wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA molecule having a sequence selected from the group consisting of SEQ ID Nos: 2-3, 5-107, 109-111, 115-171, 173-175, 177 and 179-224.





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶:G01N 33/574, 33/577, C07K 16/30, A61K
39/395, 47/48, C12Q 1/68, G01N 33/543

A2

(11) International Publication Number:

WO 98/37418

(43) International Publication Date:

27 August 1998 (27.08.98)

(21) International Application Number: PCT/US98/03690

(22) International Filing Date: 25 February 1998 (25.02.98)

(30) Priority Data:

08/806,596	25 February 1997 (25.02.97)	US
08/904,809	1 August 1997 (01.08.97)	US
09/020,747	9 February 1998 (09.02.98)	US

(71) Applicant: CORIXA CORPORATION [US/US]; Suite 200,
1124 Columbia Street, Seattle, WA 98104 (US).(72) Inventors: XU, Jiangchun; 15805 Southeast 43rd Place,
Bellevue, WA 98006 (US). DILLON, Davin. C.; 21607
N.E. 24th Street, Redmond, WA 98053 (US).(74) Agents: MAKI, David, J. et al.; Seed and Berry LLP,
6300 Columbia Center, 701 Fifth Avenue, Seattle, WA
98104-7092 (US).

(81) Designated States: AL, AM, AT, AU, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

Without international search report and to be republished upon receipt of that report.

(54) Title: COMPOUNDS FOR IMMUNODIAGNOSIS OF PROSTATE CANCER AND METHODS FOR THEIR USE

(57) Abstract

Compounds and methods for diagnosing prostate cancer are provided. The inventive compounds include polypeptides containing at least a portion of a prostate tumor protein. The inventive polypeptides may be used to generate antibodies useful for the diagnosis and monitoring of prostate cancer. Nucleic acid sequences for preparing probes, primers, and polypeptides are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/03690

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 6 and 7 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see FURTHER INFORMATION sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-10, 12-22 (all partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-10,12-22 (all partially)

Invention 1:

Methods and diagnostic kits for detecting and monitoring prostate cancer comprising either a binding agent capable of binding to a polypeptide encoded by a DNA molecule with Seq.ID 2 or oligonucleotide primers specific for a DNA molecule with Seq. ID 2 as well as an monoclonal antibody binding to a prostate protein encoded by a DNA molecule with Seq. ID 2 used in a method for inhibiting development of prostate cancer.

2. Claims: 1-10,12-22 (all partially)

Inventions 2-130:

Methods and diagnostic kits for detecting and monitoring prostate cancer comprising either a binding agent capable of binding to a polypeptide encoded by a DNA molecule with Seq.ID 3 or oligonucleotide primers specific for a DNA molecule with Seq. ID 3 as well as an monoclonal antibody binding to a prostate protein encoded by a DNA molecule with Seq. ID 3 used in a method for inhibiting development of prostate cancer.

...ibidem for Seq. ID 8-29,
41-45,47-52,54-65,70,73,74,79,81,87,90,92,93,97,103,104,107,115-160,171,181,188,191,193,194,198,203,204,207,209-211,220,222-224.

3. Claims: 1-4,8,9,11-22 (all partially)

Inventions 131-215:

Methods and diagnostic kits for detecting and monitoring prostate cancer comprising either a binding agent capable of binding to a polypeptide encoded by a DNA molecule with Seq.ID 5 or oligonucleotide primers specific for a DNA molecule with Seq. ID 5 as well as an monoclonal antibody binding to a prostate protein encoded by a DNA molecule with Seq. ID 5 used in a method for inhibiting development of prostate cancer.

...ibidem for each of Seq. ID
6,7,30-40,46,53,66-69,71,72,75-78,80,82-86,88,89,91,94-96,98-102,105,106,161-170,179,180,182-187,189,190,192,195-197,198-202,205,206,208,212-219.

INTERNATIONAL SEARCH REPORT

International Application No

PC., JS 98/03690

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 G01N33/574 G01N33/577 G01N33/543 C07K16/30 C12Q1/68
 A61K39/395 A61K47/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12Q C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 21671 A (UNIV COLUMBIA ; FISHER PAUL B (US); SHEN RUOQUAN (US)) 18 July 1996 see the whole document ---	1-10, 12-22
A	EL-SHIRBINY A M: "PROSTATIC SPECIFIC ANTIGEN" ADVANCES IN CLINICAL CHEMISTRY, vol. 31, 1994, pages 99-133, XP000617158 see the whole document ---	1-10, 12-22
A	WO 93 14775 A (WRIGHT GEORGE L JR) 5 August 1993 see the whole document ---	1-10, 12-22
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

31 August 1998

Date of mailing of the international search report

30 DEC 1998

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Hagenmaier, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT, JS 98/03690

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 09820 A (SLOAN KETTERING INST CANCER ;ISRAELI RON S (US); HESTON WARREN D W) 11 May 1994 see the whole document ---	1-10, 12-22
A	WO 95 04548 A (JENNER TECHNOLOGIES) 16 February 1995 see the whole document ---	1-10, 12-22
A	ROBSON C N ET AL: "IDENTIFICATION OF PROSTATIC ANDROGEN REGULATED GENES USING THE DIFFERENTIAL DISPLAY TECHNIQUE" PROCEEDINGS OF THE ANNUAL MEETING OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, TORONTO, MAR. 18 - 22, 1995, no. MEETING 86, 18 March 1995, AMERICAN ASSOCIATION FOR CANCER RESEARCH, page 266 XP002019344 see the whole document ---	1-10, 12-22
A	BLOK L J ET AL: "ISOLATION OF CDNAS THAT ARE DIFFERENTIALLY EXPRESSED BETWEEN ANDROGEN-DEPENDENT AND ANDROGEN-INDEPENDENT PROSTATE CARCINOMA CELLS USING DIFFERENTIAL DISPLAY PCR" PROSTATE, vol. 26, no. 4, April 1995, pages 213-224, XP000611577 see the whole document ---	1-10, 12-22
A	ALEXEYEV ET AL.: "IMPROVED ANTIBIOTIC-RESISTANCE GENE CASSETTES AND OMEGA ELEMENTS FOR E.COLI VECTOR CONSTRUCTION AND IN VITRO DELETION/INSERTION MUTAGENESIS" GENE, vol. 160, 1995, pages 63-67, XP002076033 & DATABASE EMBL AC: U35129, 1995 "pBSL141" see abstract ---	1-10, 12-22
P,A	DATABASE EMBL AC: AA453562, 11 June 1997 HILLIER ET AL.: "HOMO SAPIENS cDNA CLONE 788180" XP002075910 see abstract -----	1-10, 12-22

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT, JS 98/03690

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9621671 A	18-07-1996	AU 4751196 A	31-07-1996
		CA 2209941 A	18-07-1996
		EP 0804458 A	05-11-1997

WO 9314775 A	05-08-1993	CA 2106487 A	01-08-1993

WO 9409820 A	11-05-1994	CA 2147499 A	11-05-1994
		EP 0668777 A	30-08-1995
		JP 8506005 T	02-07-1996
		US 5538866 A	23-07-1996

WO 9504548 A	16-02-1995	AU 686660 B	12-02-1998
		AU 7631294 A	28-02-1995
		CA 2168952 A	16-02-1995
		EP 0721345 A	17-07-1996
		JP 9504000 T	22-04-1997

